

For assay of the promising substance, was chosen alkalimetry method in the medium of ethanol with a potentiometric fixation of the end point of titration ( $s = 1$ ).

**Results and discussion.** The development of quality control methodologies was carried out according to State Pharmacopoeia of Ukraine. Resynthesis of the substance was performed using reagents that were purified according to standard procedures. The developed quality control methods are acceptable for use in control and analytical laboratories.

**Conclusion.** The synthesized substance meets all the requirements of State Pharmacopoeia of Ukraine in quality.

## AMOXICILLIN AND METAL SALTS INTERACTION STUDY BY UV- SPECTROPHOTOMETRY METHODS

Dobrova A.O., Golovchenko O.S.

Scientific supervisor: prof. Georgiyants V.A.

National University of Pharmacy, Kharkiv, Ukraine

anna.dobrova08@gmail.com

**Introduction.** Amoxicillin is an antibiotic of the penicillin group, contains carboxyl and amino groups and can enter into complex formation reactions. Joint administration of amoxicillin with preparations containing transition metal salts can lead to the formation of chelate complexes of various structures, that can affect the effectiveness of therapy. Previously, we conducted a study of the amoxicillin interaction with metal salts in molar ratios. However, this method does not allow to say in what ratio these compounds can be formed. In an attempt to establish it, a Job spectrophotometric analysis was performed.

**Aim.** To study the complexation of amoxicillin with metal salts using Job's spectrophotometric method.

**Materials and methods.** As stated in the Job's method, initial solutions of amoxicillin and salts with equal molar ratios ( $1 \cdot 10^{-3}$  M) were prepared. Further, primary solutions were diluted and series of solutions with ratios of antibiotic and salt 1:9, 2:8, 3:7, 4:6, 5:5, 6:4, 7:3, 8:2, 9:1 were prepared. Then the absorbance was measured at a wavelength 200-400nm on spectrophotometer Evolution 60S. Samples of calcium chloride, magnesium sulfate heptahydrate, aluminium sulfate, iron (III) chloride, iron (II) sulfate salts were taken as investigated. All solutions were prepared in 0.1 M hydrochloric acid medium which corresponds to pH of stomach.

**Results and discussion.** The UV-spectrum of amoxicillin in 0.1M hydrochloric acid has three maxima at wavelengths 203nm, 230nm and 272 nm. To control, we took into account changes in absorbance at all maxima. In the experiment with salts of calcium, iron(II) and aluminum, a proportional increase in the absorbance in all three maxima was observed, respectively as the concentration of the antibiotic increased. While when adding iron (III) salt, a change in the character of the spectrum was observed at a ratio of 1: 9, 2: 8, 3: 7, 4: 6, 5: 5 due to which a characteristic's maximum blur occurred at a wavelength of 272 nm.

**Conclusions.** Based on the results of the study, iron (III) salts are most likely able to interact with the formation of different ratios complex compounds. The obtained results confirm the importance of further research of amoxicillin's complexation properties with the antacids and other metal salts containing medications.

## RESEARCH OF THE SQUALENE CONTAIN IN PUMPKIN SEEDS BY A THIN-LAYER CHROMATOGRAPHY

Kotov S. A., Kotova E. E.

Scientific supervisor: Doctor of Pharmacy Gontova T. M.

National University of Pharmacy, Kharkiv, Ukraine

drsnuff@gmail.com

**Introduction.** Cucurbita pepo seeds are a well-known traditional herbal drug that has been used throughout the ages in folk and formal medicine. Pumpkin seeds contain a various chemical