

O. V. Kutova, I. V. Kovalevska, N. V. Demchenko

National University of Pharmacy of the Ministry of Health of Ukraine

Development of a methodological approach to determine regression equations in the study of the technology for manufacturing tablets based on quercetin

When developing the technology of tablets and tablet masses one of the tasks is to determine the amounts of excipients required to obtain pharmaco-technological quality indicators that meet the requirements of the State Pharmacopoeia of Ukraine (SPHU).

Aim. To develop an algorithm for determining the type of three-factor mathematical models with dependent variables.

Materials and methods. The study object was experimental observations of the quantitative effect of excipients in the tablet composition based on a solid dispersion of quercetin on the pharmacopoeial characteristics of this dosage form, in particular on its flowability. The experimental data were processed by the planned experiment using Mathcad 15 and MS Excel software according to the algorithm proposed.

Results and discussion. It has been found that the identification of mathematical models in pharmaceutical studies with three dependent factors, which total value is determined by the quantitative composition of the dosage form and fixed at a certain level, is difficult to perform due to the difficulty of interpreting the multiple regression parameters as characteristics of factors in isolation through their correlability. It has been proven that the replacement of variables leads to the determination of a mathematical model that does not reveal the mechanism of the factors' action and is a static description of their overall impact on the indicator studied.

Conclusions. As a result of the research conducted, regression equations were found to determine the effect of the amount of these excipients on the tablet flowability. It has been found that the most influential role in determining the target in the factor space studied is played by the interaction of factors. The algorithm for determining the mathematical description of the dependence with three variables has been proposed. Based on the fact that the research is conditioned by strict conditions in the quantitative composition, the chosen mathematical model of the dependence of the quadratic equation with two factors, for which there are limitations, has no correlation. The model determined is characterized by the possibility of providing the graphical interpretation and simplification of analysis and can be used for forecasting and optimization.

Key words: *quantitative effect of excipients; three-factor mathematical model; dependent mixture factors; identification algorithm*

O. V. Кутова, І. В. Ковалевська, Н. В. Демченко

Національний фармацевтичний університет Міністерства охорони здоров'я України

Розроблення методологічного підходу до визначення регресійних рівнянь під час дослідження технології виготовлення таблеток на основі кверцетину

Під час розроблення технології таблеток і таблеткових мас одним із завдань є визначення кількості допоміжних речовин, необхідних для отримання фармакотехнологічних показників якості, які відповідають вимогам Державної фармакопеї України.

Метою роботи є розроблення алгоритму визначення виду трифакторних математичних моделей із залежними змінними.

Матеріали та методи. Об'єктом досліджень були експериментальні спостереження кількісного впливу допоміжних речовин у складі таблеток на основі твердої дисперсії кверцетину на фармакопейні характеристики зазначеної лікарської форми, зокрема на її сипкість. Експериментальні дані обробляли з використанням запланованого експерименту за допомогою комп'ютерних програм Mathcad 15 та MS Excel за запропонованим алгоритмом.

Результати та їх обговорення. Визначено, що ідентифікацію математичних моделей у фармацевтичних дослідженнях з трьома залежними факторами, сумарне значення яких визначається кількісним складом лікарської форми та фіксується на певному рівні, важко проводити внаслідок ускладнення інтерпретації параметрів множинної регресії як характеристик факторів у ізольованому вигляді через їх корельованість. Доведено, що заміна змінних призводить до визначення математичної моделі, яка не розкриває механізму дії факторів і є статичним описом їх загального впливу на досліджуваний показник.

Висновки. У результаті проведених досліджень визначено рівняння регресії для виявлення впливу кількості зазначених допоміжних речовин на сипкість таблетки. З'ясовано, що найбільш впливову роль у визначенні цільового показника в досліджуваному факторному просторі відіграє взаємодія факторів. Запропоновано алгоритм визначення математичного опису залежності з трьома змінними. З огляду на те, що дослідження мають відповідати жорстким умовам за кількісним складом, обрано математичну модель залежності у вигляді квадратичного рівняння з двома факторами, за якими є обмеження і між якими немає кореляції. Така модель характеризується можливістю надання графічної інтерпретації та спрощення аналізу й може бути використана для прогнозування і оптимізації.

Ключові слова: *кількісний вплив допоміжних речовин; трифакторна математична модель; залежні змінні фактори; алгоритм ідентифікації*

О. В. Кутовая, И. В. Ковалевская, Н. В. Демченко

Национальный фармацевтический университет Министерства здравоохранения Украины

Разработка методологического подхода к установлению регрессионных уравнений при исследовании технологии изготовления таблеток на основе кверцетина

При разработке технологии таблеток и таблеточных масс одной из задач является определение количества вспомогательных веществ, необходимых для получения фармакотехнологических показателей качества, соответствующих требованиям Государственной фармакопеи Украины.

Целью работы является разработка алгоритма установления вида трехфакторных математических моделей с зависимыми переменными.

Материалы и методы. Объектом исследований были экспериментальные наблюдения количественного влияния вспомогательных веществ в составе таблеток на основе твердой дисперсии кверцетина на фармакопейные характеристики указанной лекарственной формы, в частности на ее сыпучесть. Обработку экспериментальных данных проводили с использованием планируемого эксперимента с помощью компьютерных программ Mathcad 15 и MS Excel по предложенному алгоритму.

Результаты и их обсуждение. Определено, что идентификацию математических моделей в фармацевтических исследованиях с тремя зависимыми факторами, суммарное значение которых определяется количественным составом лекарственной формы и фиксируется на определенном уровне, трудно проводить вследствие осложнения интерпретации параметров множественной регрессии как характеристик факторов в изолированном виде из-за их коррелируемости. Доказано, что замена переменных приводит к установлению математической модели, которая не раскрывает механизма действия факторов и является статическим описанием их общего влияния на исследуемый показатель.

Выводы. В результате проведенных исследований определены уравнения регрессии для установления влияния количества указанных вспомогательных веществ на сыпучесть таблетки. Установлено, что наиболее влиятельную роль в определении целевого показателя в исследуемом факторном пространстве играет взаимодействие факторов. Предложен алгоритм определения математического описания зависимости с тремя переменными. Исходя из того, что исследования должны отвечать жестким условиям по количественному составу, выбрана математическая модель зависимости в виде квадратичного уравнения с двумя факторами, по которым есть ограничения и между которыми нет корреляции. Установленная модель характеризуется возможностью предоставления графической интерпретации и упрощения анализа и может быть использована для прогнозирования и оптимизации.

Ключевые слова: количественное влияние вспомогательных веществ; трехфакторная математическая модель; зависимые переменные факторы; алгоритм идентификации

Introduction. Quantitative research in the development of the tablet technology involves finding the optimal composition of a dosage form (fillers, disintegrants, binders, glidants and lubricants, etc.), parameters of technological modes determined by the type of a dosage form (compression pressure, compression rate, film-forming solution temperature), the drying time of the coating, etc.), as well as ensuring the values of pharmaco-technological quality indicators, in accordance with the requirements of the State Pharmacopoeia of Ukraine (SPhU).

It should be noted that there is often a need to find a significant amount of excipients when developing the technology of tablet drugs. Multifactoriality is one of the specific features of these studies at the experimental stage. In this study, the determination of regression relationships is associated with three quantitative factors, which total value is fixed at a certain level. Interpretation of multiple regression parameters in this case is complicated by their correlation.

Various modifications of the planned experiment are used to construct multifactor models [1-5]. In the study of three factors in the presence of certain restrictions on two of them, the third factor becomes dependent and changes under the conditions of the passive experiment. As a result, the levels of the dependent factor x_3 in the experiments are only recorded, but not specified. In this case, the researcher is not able to control the dependent factor and becomes a passive observer of the phenome-

na occurring. We have to deal with the values of the factors, which change unevenly. Due to the uneven change in the corresponding range of the dependent factor, the usual methods of finding empirical dependences are unsuitable, and there is a need to develop the effective one in the conditions studied [6].

The aim of the article was to develop an algorithm for determining the type of three-factor mathematical models with dependent variables.

Materials and methods. The algorithm for identifying the regression equation for three dependent quantitative factors is shown on the example of the development of the tablet technology based on solid dispersion of quercetin [7]. According to the standard procedure at the first stage as a result of studies, the effect of excipients on the pharmacological characteristics of tablets was determined by their qualitative composition. Three excipients significantly affecting the tablet mixture and tablets – microcrystalline cellulose (x_1 factor), nesulin (x_2 factor) and sodium starch glycolate (x_3 factor) were selected. At the second stage it was necessary to determine the quantitative content of excipients under the following conditions: $12 < x_1 < 14$, $12 < x_2 < 14$, $x_1 + x_2 + x_3 = 29$.

The tablet weight and finished tablets were monitored according to five pharmaco-technological indicators (reviews, target functions): y_1 – the Kara index, y_2 – the tablet mass flowability, y_3 – the Hausner coefficient, y_4 – the tablet strength, y_5 – the tablet disintegration time.

The factor experiment planning matrix 2^2 and the research results

No.	x1	x2	y1	y2	y3	y4	y5
1	14 (+)	12 (-)	21.42 ± 0.4	4.5 ± 0.35	0.85 ± 0.05	0.56 ± 0.04	10 ± 0.5
2	14 (+)	14 (+)	12.03 ± 0.6	7.92 ± 0.86	1.39 ± 0.07	0.58 ± 0.03	13 ± 0.7
3	12 (-)	14 (+)	30.23 ± 0.9	2.68 ± 0.4	1.31 ± 0.055	0.61 ± 0.04	9 ± 0.66
4	12 (-)	12 (-)	17.0 ± 0.3	4.83 ± 0.43	1.15 ± 0.03	0.52 ± 0.03	5 ± 0.45

In classical regression analysis, the structure considered is known before planning, but in pharmaceutical research it is not always so. In many cases, the purpose of building a regression model is to elucidate the relationship between independent factors and the dependent pharmacopoeial index. Determining this structure is one of the tasks in the regression analysis. It is usually formed as a search for an equation with the best estimates of static significance. This is not entirely true as the aim of the study is not the goal of mathematical statistics alone.

The strategy for constructing the equation determines the requirements for planning the experiment. In this case, the formulation of the problem does not ensure the independence of factors, and as a result, there will be no good conditionality of the matrix full of factor planning.

To achieve the factor independence, the variable x3 was excluded from consideration [8-10]. This allowed us to apply the experimental areas of the standard form 2^2 and 3^2 .

The mathematical processing of the results of the study was performed using Mathcad 15 software [11], the least squares method (LSM). The estimation of the significance of regression equations and their coefficients was performed using statistical functions of the MS Excel spreadsheet [12].

Results and discussion. For this study, the effect of the third excipient P – a loosening substance, which amount directly depends on the number of the first two excipients, is excluded from consideration as the most correlated factor. To propose a type of the regression

equation, we used a graphical approach based on the visual analysis of location of the points of the experimental surface obtained according to Plan 2^2 (Tab. 1) using Mathcad 15 software (Fig. 1a).

The assumption about the linearity of the mathematical description was refuted by the type of response surface. In addition, the coefficient of determination of the linear equation $y_2 = f(x_1, x_2)$ was equal to 0.28, the average relative error of calculations at base points exceeded the accepted 3 %. Based on the graphical interpretation of the experimental dependence and the experimental plan 2^2 , the regression equations with the interaction of factors were analyzed:

$$y_2(x_1, x_2) = a_0 + a_1x_1 + a_2x_2 + a_3x_1x_2. \quad (1)$$

The plan with the minimum number of experiments (saturated plan) allows the method of least squares to determine the coefficients of the equation (1):

$$y_2(x_1, x_2) = 230.23 - 17.55x_1 - 18.035x_2 + 1.405x_1x_2, \quad (2)$$

but the requirements for the quality of the evaluation of the function and its parameters are not met despite the maximum coefficient of determination ($R^2 = 1$), the absence of a relative calculation error at base points and satisfactory coincidence of experimental and theoretical response surfaces (Fig. 2).

To make a final decision on the determination of the mathematical model, despite a good assessment of the adequacy




Fig. 1. The experimental response surfaces: a) by complete factorial experiment 2^2 , b) by complete factorial experiment 3^2




Fig. 2. The graphical interpretation of the dependence $y_2 = f(x_1, x_2)$: 1) experimental, 2) obtained by the equation (1)

of two statistical indicators, it was proposed to analyze the separate influence of factors on the target indicator within the interval studied. In this study, the increase in the content of microcrystalline cellulose with different amounts of neusilin in different ways affects the target: if $x_2 = 12\%$, when the greater x_1 , the lower the value of y_2 ; if $x_2 = 14\%$, then an increase in x_1 leads to an increase in $y_2 = f(x_1)$ (Fig. 3a).

An increase in the content of neusilin at $x_1 = 12\%$ leads to a decrease in the value of the mass flowability, at $x_1 = 14\%$ – to an increase in this indicator (Fig. 3b).

The main effect of the factors is masked and gives the final increasing linear dependence, hiding the non-linear behavior of variables within the factor space. Thus, the available information indicates the impossibility of using the equation (1) as a mathematical description of this study despite a good assessment of its adequacy on two statistical indicators, and indicates the need to consider a second-order regression model.

Increasing the experimental base by adding the experiment in the center of the plan $2^2 - y_2(13.13) = 7.0$ allows obtaining statistical estimates of the equation (1): the coefficient of determination – $R^2 = 0.77$ shows a sufficient approximation accuracy; the reliability according to the level of significance of the Fisher criterion is

$$\begin{aligned} Y_2(14, 12) &= 4.15 & \frac{y_2^2_0 - 4.15}{y_2^2_0} &= 0 \\ Y_2(14, 14) &= 7.42 & \frac{y_2^2_1 - 7.42}{y_2^2_1} &= 0 \\ Y_2(12, 14) &= 3.18 & \frac{y_2^2_2 - 3.18}{y_2^2_2} &= 0 \\ Y_2(12, 12) &= 5.53 & \frac{y_2^2_3 - 5.53}{y_2^2_3} &= 0 \end{aligned}$$

0.58 – the model is not significant; the p-values for the coefficients of the equation – 0.35, and it proves their insignificance. The results obtained do not allow us to consider the regression equation (1) as an adequate mathematical model.

There is a need to plan an experiment for a quadratic model. It is advisable to build the optimal plan for the quadratic model in such a way that it includes the points of the optimal plan for the linear model, maintaining orthogonality (Tab. 2).

The graphical interpretation of the experimental response surface y_2 obtained from the results of the experimental matrix according to Plan 3^2 differs from option 2^2 (Fig. 1b).

Calculations by the 2nd order regression equation, which coefficients are determined using the *regress* function in Mathcad 15:

$$\begin{aligned} y_2(x_1, x_2) &= -93.3472 - 17.8683x_1 + \\ &+ 32.325x_2 + 1.405x_1x_2 + \\ &+ 0.0133x_1^2 - 1.9367x_2^2, \end{aligned} \quad (3)$$

give a slight average relative error at base points not exceeding 0.5%. Experimental and theoretical response surfaces, which are given in Fig. 4, match quite well.




Fig. 3. The scattering diagram for factors: a) $y_2 = f(x_1)_{x_2 = \text{const}}$, b) $y_2 = f(x_2)_{x_1 = \text{const}}$

Table 2

The factor experiment planning matrix 3^2 and the research results

No.	x1	x2	y1	y2	y3	y4	y5
1	14 (+)	12 (-)	21.42 ± 0.4	4.15 ± 0.35	0.85 ± 0.05	0.56 ± 0.04	10 ± 0.5
2	14 (+)	14 (+)	12.03 ± 0.6	7.42 ± 0.86	1.39 ± 0.07	0.58 ± 0.03	13 ± 0.7
3	12 (-)	14 (+)	30.23 ± 0.9	3.18 ± 0.4	1.31 ± 0.055	0.61 ± 0.04	9 ± 0.66
4	12 (-)	12 (-)	17.0 ± 0.3	5.53 ± 0.43	1.15 ± 0.03	0.52 ± 0.03	5 ± 0.45
5	14 (+)	13 (0)	16.7 ± 0.35	7.8 ± 0.8	1.17 ± 0.32	0.62 ± 0.04	10.2 ± 1.2
6	12 (-)	13 (0)	23.6 ± 0.62	6.2 ± 0.64	1.19 ± 0.3	0.61 ± 0.05	5.5 ± 0.6
7	13 (0)	14 (+)	21.1 ± 1.1	5.3 ± 0.52	1.35 ± 0.35	0.56 ± 0.05	10.1 ± 0.9
8	13 (0)	12 (-)	19.2 ± 0.8	4.8 ± 0.4	1.17 ± 0.4	0.57 ± 0.07	5.5 ± 0.6
9	13 (0)	13 (0)	20.15 ± 0.95	7.0 ± 0.58	1.26 ± 0.45	0.58 ± 0.04	6.4 ± 0.54

Statistical estimates of the equation (3) are as follows: the coefficient of determination – $R^2 = 0.999$ shows an excellent approximation accuracy; the reliability according to the level of significance of the Fisher criterion is 0.0000429, that is, the model is significant; the p-value for all coefficients of the equation is much lower than 0.05, except for the value of 0.77 for the coefficient at $x1^2$. The presence of alternating positive and negative deviations indirectly confirms the absence of a systematic error in the regression equation constructed. Confidence intervals of the function at base points are: $y_2(14,12) = 4.1586 \pm 3.82$; $y_2(14,14) = 7.4402 \pm 3.82$; $y_2(12,14) = 3.145 \pm 3.82$; $y_2(12,12) = 5.4836 \pm 3.82$; $y_2(14,13) = 7.736 \pm 3.17$; $y_2(12,13) = 6.25 \pm 3.17$; $y_2(13,14) = 5.2794 \pm 3.17$; $y_2(13,12) = 4.808 \pm 3.17$; $y_2(13,13) = 6.980 \pm 3.17$.

The results obtained allow us to consider the regression equation (3) as an adequate mathematical description.

By the equation (3) it is impossible to determine the true mechanism of influence of factors $x1, x2, x3$ on the resultant trait due to the reasons previously mentioned. Therefore, the real mechanism of action and interaction of factors can be proposed by comparing the possible mathematical structures of the equation (3), and using the substitution $x3 = 29 - x1 - x2$. The equation (5) is identical to three parametric dependences:

$$y_2(x_1, x_2) = a_0 + a_1x_1 + a_2x_2 + a_3x_3 + a_4x_1x_2 + a_5x_1x_3 + a_6x_2x_3. \tag{4}$$

Taking into account that the assessment of the influence of the coefficients of the equation (5) using MS Excel, shows no significance of the component $x1^2$ (p -value is equal to 0.77), it is possible not to consider also the interaction of variables at level $x1x3$. Thus, we can offer the form of the equation as a mathematical description:


$$y_2(x_1, x_2, x_3) = a_0 + a_1x_1 + a_2x_2 + a_3x_3 + a_4x_1x_2 + a_5x_2x_3 + a_6x_2^2, \tag{5}$$

with the coefficients determined:

$$y_2(x_1, x_2, x_3) = -603.72 + 17.522x_3 + 3.124x_1x_2 + 1.719x_2x_3 - 0.218x_2^2. \tag{6}$$

In the equation (8), all the coefficients are significant, the average calculation error is 0.6 %, the coefficient of determination – $R^2 = 0.9994$ shows an excellent approximation accuracy; the reliability according to the level of significance of the Fisher criterion is 0.0000562, the graphical interpretation of the given equation proves the coincidence of the response surfaces similar to Fig. 4.

Confidence intervals of the function at base points were determined using the Student's test with 95 % reliability: $y_2(14,12) = 4.15 \pm 3.82$; $y_2(14,14) = 7.45 \pm 3.82$; $y_2(12,14) = 3.15 \pm 3.82$; $y_2(12,12) = 5.49 \pm 3.82$; $y_2(14,13) = 7.74 \pm 3.17$; $y_2(12,13) = 6.256 \pm 3.17$; $y_2(13,14) = 5.3 \pm 3.17$; $y_2(13,12) = 4.826 \pm 3.17$; $y_2(13,13) = 6.99 \pm 3.17$.



$\frac{y_2^0 - 4.171}{y_2^0} = -5.06 \times 10^{-3}$	$\frac{y_2^4 - 7.743}{y_2^4} = 7.308 \times 10^{-3}$
$\frac{y_2^1 - 7.455}{y_2^1} = -4.717 \times 10^{-3}$	$\frac{y_2^5 - 6.257}{y_2^5} = -9.194 \times 10^{-3}$
$\frac{y_2^2 - 3.159}{y_2^2} = 6.604 \times 10^{-3}$	$\frac{y_2^6 - 5.287}{y_2^6} = 2.453 \times 10^{-3}$
$\frac{y_2^3 - 5.495}{y_2^3} = 6.329 \times 10^{-3}$	$\frac{y_2^7 - 4.813}{y_2^7} = -2.708 \times 10^{-3}$

Fig. 4. The graphical interpretation of equation (5) and determination of the calculation error using it

In the same way, regression equations were determined for all targets studied in order to further optimize the technology of manufacturing tablets [13].

Conclusions and prospects for further research

The essence of the methodological approach proposed to the determination of regression equations is reflected in the example of a mathematical description of one of the five pharmacopoeial characteristics of tablets studied – the flowability of the tablet mass based on quercetin.

To reduce the required number of experiments, taking into account the dependence of the variables affecting the target indicator, the mathematical processing of the research results has been performed using two factors, which content in the dosage form studied is strictly limited by the research conditions.

A satisfactory estimate of the regression equation determined by the fractional plan with two variables has been revised after the analysis of the isolated influence of factors. There is a need to expand the information base within the factor space in order to consider the regression equation of the 2nd order.

The result of the mathematical processing is the determination of a regression equation in the form of an incomplete algebraic polynomial of the 2nd degree, in which the constants determined do not reflect the actual degree of influence of the variables. After the mathematical transformation, the dependence of three variables with refined coefficients has been obtained, revealing the real mechanism of the influence of factors on the indicator studied. Thus, the presence of the excipients in combination with the active ingredients in the formulation of tablets developed has a significant effect on the flowability of the test dosage form. Increasing the amount of neusilin from 12 to 14 % can lead to a decrease in flowability, but in the mixture with the amount of microcrystalline cellulose from 12 to 14 % and sodium starch glycolate from 1 to 3 % the target value increases.

The methodological approach proposed is applied to all pharmacopoeial characteristics studied. The regression equations obtained can be used to optimize the technology of manufacturing tablets.

Conflict of interests: authors have no conflict of interests to declare.

REFERENCES

1. Лапач С. Н. Регрессионный анализ. Процессный подход. *Математичні машини і системи*. 2016. № 1. С. 129–138.
2. Математичне планування експерименту при проведенні наукових досліджень в фармації / Т. А. Грошовий та ін. Тернопіль: Укрмедкнига, 2008. 367 с.
3. Песков К. В. Математическое моделирование при разработке лекарств. *Вестник «Росздравнадзора»*. 2013. № 1. С. 57–60.
4. Zivorad R. Lazic. Design of experiments in chemical engineering. A practical guide. Weinheim: Wiley-VCH, 2004. 610 p.
5. SampleSizeR: calculate sample sizes within completely randomized design. URL: <http://shiny.math.uzh.ch/git/reinhard.furrer/SampleSizeR/>.
6. Свідоцтво про реєстрацію авторського права на твір № 47998 від 25.02.2013. Комп'ютерна програма для автоматичної побудови математичної моделі функціонування довільного об'єкту / А. О. Журавський та ін. *Авторське право і суміжні права. Офіційний бюлетень*. 2013. № 30. С. 365.
7. Optimization of the composition of solid dispersion of quercetin / I. Kovalevska et al. *Current Issues in Pharmacy and Medicine Science*. 2021. Vol. 34, №. 1. P. 1–4. DOI: <https://doi.org/10.2478/cipms-2021-0001>.
8. Множественный регрессионный анализ. URL: <https://math.semestr.ru/regress/mregres.php>.
9. Draper N., Smith H. Applied Regression Analysis. 3rd ed. New-York: Wiley, 2014. 912 p.
10. Сенин А. Методы отбора переменных в регрессионные модели. URL: <https://basegroup.ru/community/articles/feature-selection>.
11. Воскобойников Ю. Е. Регрессионный анализ данных в пакете MATHCAD. Санкт-Петербург : Лань, 2011. 224 с.
12. Воскобойников Ю. Е. Эконометрика в Excel. Новосибирск, 2011. 154 с.
13. Свідоцтво про реєстрацію авторського права на твір № 82707 від 25.01.2019. Методика визначення оптимальних параметрів технологічного процесу : метод. рек. / О. В. Кутова та ін. *Авторське право і суміжні права. Офіційний бюлетень*. 2019. № 51. С. 337.

REFERENCES

1. Lapach, S. N. (2016). *Matematychni mashyny i systemy, 1*, 129–138.
2. Hroshovi, T. A., Martseniuk, V. P., Kucherenko, L. I. et al. (2008). *Matematychni planuvannia eksperymentu pry provedenni naukovykh doslidzhen' v farmatsii*. Ternopil': Ukrmedknyha, 367.
3. Peskov, K. V. (2013). *Vestnik «Roszdravnadzora», 1*, 57–60.
4. Zivorad, R. Lazic. (2004). *Design of experiments in chemical engineering. A practical guide*. Weinheim: Wiley-VCH, 610.
5. SampleSizeR: calculate sample sizes within completely randomized design. Available at: <http://shiny.math.uzh.ch/git/reinhard.furrer/SampleSizeR/>.
6. Zhuravs'kyi, A. O., Torianyk, E. I., Siniaieva, O. V., Zavorodnii, O. I. (2013). Svidotstvo pro reiestratsiiu avtorskoho prava na tvir № 47998 vid 25.02.2013. Komp'uterna prohrama dlia avtomatychnoi pobudovy matematychnoi modeli funktsiiuvannia dovilnoho ob'iektu. *Avtorske pravo i sumizhni prava. Ofitsiyni biuleten*, 30, 365.
7. Kovalevska, I., Ruban, O., Kutova, O., Levachkova, O. (2021). Optimization of the composition of solid dispersion of quercetin. *Current Issues in Pharmacy and Medicine Science, 1* (34), 1–4. doi: <https://doi.org/10.2478/cipms-2021-0001>.
8. Mnozhestvennyi reghressionnyi analiz. Available at: <https://math.semestr.ru/regress/mregres.php>.
9. Draper, N., Smith, H. (2014). *Applied Regression Analysis* (3rd ed.). New-York: Wiley, 912.
10. Senin, A. Metody otbora peremennykh v reghressionnye modeli. *basegroup.ru*. Available at: <https://basegroup.ru/community/articles/feature-selection>.

11. Voskoboinikov Yu. Ye. (2011). *Rehresionnyi analiz danykh v pakete MATHCAD*. Saint Petersburg: Lan, 224.
12. Voskoboinikov, Yu. Ye. (2021). *Ekonometrika v Excel*. Novosibirsk, 154.
13. Kutova, O. V., Kovalevs'ka, I. V., Kutovyi, D. S., Shapovalov, O. V., Zhuravs'kyi, A. O. (2019). Svidotstvo pro reiestratsiiu avtorskoho prava na tvir № 82707 vid 25.01.2019. *Metodyka vyznachennia optymalnykh parametriv tekhnolohichnoho protsesu. Avtorske pravo i sumizhni prava. Ofitsiinyi biuletен, 51, 337.*

Information about authors:

Kutova O. V., Candidate of Engineering Science (Ph.D.), associate professor of the Department of Technologies of Pharmaceutical Preparations, National University of Pharmacy of the Ministry of Health of Ukraine. E-mail: paxtoxt@gmail.com. ORCID: <https://orcid.org/0000-0002-3761-2831>

Kovalevska I. V., Doctor of Pharmacy (Dr. habil.), associate professor of the Department of Industrial Technology of Drugs, National University of Pharmacy of the Ministry of Health of Ukraine. E-mail: inga.kovalevskaya@gmail.com. ORCID: <https://orcid.org/0000-0001-5610-8334>

Demchenko N. V., Candidate of Economics (Ph.D.), Associate Professor of the Organization and Economics of Pharmacy Department, National University of Pharmacy of the Ministry of Health of Ukraine. E-mail: demchenata@ukr.net. ORCID: <https://orcid.org/000-0001-5915-0087>

Відомості про авторів:

Кутова О. В., кандидатка техн. наук, доцентка кафедри технологій фармацевтичних препаратів, Національний фармацевтичний університет Міністерства охорони здоров'я України. E-mail: paxtoxt@gmail.com. ORCID: <https://orcid.org/0000-0002-3761-2831>

Ковалевська І. В., докторка фармац. наук, доцентка кафедри заводської технології ліків, Національний фармацевтичний університет Міністерства охорони здоров'я України. E-mail: inga.kovalevskaya@gmail.com. ORCID: <https://orcid.org/0000-0001-5610-8334>

Демченко Н. В., кандидатка екон. наук, доцентка кафедри організації та економіки фармації, Національний фармацевтичний університет Міністерства охорони здоров'я України. E-mail: demchenata@ukr.net. ORCID: <https://orcid.org/000-0001-5915-0087>

Сведения об авторах:

Кутовая О. В., кандидат техн. наук, доцент кафедры технологий фармацевтических препаратов, Национальный фармацевтический университет Министерства здравоохранения Украины. E-mail: paxtoxt@gmail.com. ORCID: <https://orcid.org/0000-0002-3761-2831>

Ковалевская И. В., доктор фармац. наук, доцент кафедры заводской технологии лекарств, Национальный фармацевтический университет Министерства здравоохранения Украины. E-mail: inga.kovalevskaya@gmail.com. ORCID: <https://orcid.org/0000-0001-5610-8334>

Демченко Н. В., кандидат экон. наук, доцент кафедры организации и экономики фармации, Национальный фармацевтический университет Министерства здравоохранения Украины. E-mail: demchenata@ukr.net. ORCID: <https://orcid.org/000-0001-5915-0087>

Надійшла до редакції 02.12.2021 р.