МІНІСТЕРСТВО ОХОРОНИ ЗДОРОВ'Я УКРАЇНИ
НАЦІОНАЛЬНИЙ ФАРМАЦЕВТИЧНИЙ УНІВЕРСИТЕТ
ІНСТИТУТ ПІДВИЩЕННЯ КВАЛІФІКАЦІЇ
СПЕЦІАЛІСТІВ ФАРМАЦІЇ
КАФЕДРА КЛІНІЧНОЇ ФАРМАКОЛОГІЇ

АКТУАЛЬНІ ПИТАННЯ КЛІНІЧНОЇ ФАРМАКОЛОГІЇ ТА КЛІНІЧНОЇ ФАРМАЦІЇ (TOPICAL ISSUES OF CLINICAL PHARMACOLOGY AND CLINICAL PHARMACY)

Матеріали науково-практичної

Internet-конференції з міжнародною участю



29-30 жовтня 2024 року

м. Харків

МІНІСТЕРСТВО ОХОРОНИ ЗДОРОВ'Я УКРАЇНИ
НАЦІОНАЛЬНИЙ ФАРМАЦЕВТИЧНИЙ УНІВЕРСИТЕТ
ІНСТИТУТ ПІДВИЩЕННЯ КВАЛІФІКАЦІЇ СПЕЦІАЛІСТІВ ФАРМАЦІЇ
КАФЕДРА КЛІНІЧНОЇ ФАРМАКОЛОГІЇ

АКТУАЛЬНІ ПИТАННЯ КЛІНІЧНОЇ ФАРМАКОЛОГІЇ ТА КЛІНІЧНОЇ ФАРМАЦІЇ
(TOPICAL ISSUES OF CLINICAL PHARMACOLOGY AND CLINICAL PHARMACY)

Матеріали науково-практичної
Internet-конференції з міжнародною участю

29-30 жовтня 2024 року
м. Харків

Харків
НФаУ
2024

УДК 615.1:615.03
А 43

Редакційна колегія та оргкомітет: проф. Котвіцька А. А. (голова), проф. Федосов А. І. (співголова), проф. Владимирова І. М. (співголова), доц. Суріков О. О. (співголова), проф. Бутко Я. О. (співголова), проф. Міщенко О. Я. (співголова, відповідальний секретар), проф. Бондарєв Є. В. (член оргкомітету), доц. Файзуллін О. В. (член оргкомітету), доц. Мороз С. Г. (член оргкомітету), доц. Халєєва О. Л. (член оргкомітету), доц. Березняков А. В. (член оргкомітету), проф. Ткачова О. В., проф. Кіреєв І. В., проф. Риженко І. М., проф. Подольський І. М., доц. Козар В. В., доц. Шаповалова О. В., асист. Тимченко Ю. В., асист. Андріяненков О. В., асист. Кузенков Р. В.

Реєстраційне посвідчення в Укр ІНТЕІ № 613 від 11 грудня 2023 р.

Актуальні питання клінічної фармакології та клінічної фармації (Topical issues of clinical pharmacology and clinical pharmacy): матеріали наук.-практ. internet-конф. з міжнар. участю (29-30 жовт. 2024 р., м. Харків) / редкол. : О. Я. Міщенко, Я. О. Бутко, Є. В. Бондарєв та ін. – Харків : НФаУ, 2024. – 388 с.

У збірнику опубліковані матеріали науково-практичної internet-конференції з міжнародною участю «Актуальні питання клінічної фармакології та клінічної фармації». У матеріалах конференції висвітлено клініко-фармакологічні аспекти застосування лікарських засобів; питання безпеки фармакотерапіїі та системи фармаконагляду; адміністрування антимікробних препаратів як мультидисциплінарного підходу протидії антибіотикорезистентності; доказової медицини в клінічній практиці; вікові, гендерні, хронофармакологічні, фармакогенетичні чинники раціонального застосування лікарських засобів; інформаційні технології в клінічній фармакології та клінічній фармації; організаційно-управлінські підходи забезпечення раціонального застосування лікарських засобів; оцінка технологій охорони здоров'я, фармакоекономічні дослідження та формулярна система як складові забезпечення раціональної фармакотерапії.

Наведено технологічні аспекти розробки, впровадження та використання лікарських засобів; доклінічні та клінічні дослідження лікарських засобів. Розглянуто сучасні методичні аспекти викладання клінічної фармакології та клінічної фармації в Україні та світі.

Видання представляє інтерес для наукових, педагогічних та практичних фармацевтичних та медичних працівників.

Матеріали подаються мовою оригіналу. За достовірність інформації відповідальність несуть автори.

УДК 615.1:615.03

© НФаУ, 2024

Розділ 1

КЛІНІКО-ФАРМАКОЛОГІЧНІ АСПЕКТИ ЗАСТОСУВАННЯ ЛІКАРСЬКИХ ЗАСОБІВ

КЛІНІЧНА ФАРМАКОЛОГІЯ ТА ПЕРСПЕКТИВНІ ТЕНДЕНЦІЇ РОЗРОБКИ ЛІКІВ

Міщенко О. Я., Бутко Я. О., Андріяненков О. В.
Національний фармацевтичний університет, м. Харків, Україна
Кафедра клінічної фармакології
Інституту підвищення кваліфікації спеціалістів фармації
clinpharmacol_ipksph@nuph.edu.ua

Вступ. Клінічна фармакологія – це вивчення ліків на людях, від перших досліджень на людях до рандомізованих контрольованих досліджень (РКД) і оцінки співвідношення користь/ризик у великих популяціях. Клінічна фармакологія відіграє важливу роль у прийнятті обґрунтованих рішень на етапі розробки ліків, оскільки вона зосереджена на впливі ліків на людину. Клінічна фармакологія з такими складовими як фармакокінетика, фармакодинаміка та фармакогеноміка, є базовими на всіх етапах розробки ліків, включаючи вибір оптимальних доз для досліджень фази I, II і III, оцінку біоеквівалентності та біоподібних досліджень і планування клінічних досліджень. Включення клінічної фармакології в дослідження, а також у вимоги регуляторних органів покращує процес розробки ліків. Клінічна фармакологія також застосовується для безпосереднього догляду за пацієнтами з метою персоналізації лікування. Для оптимізації дозування для пацієнтів на індивідуальному рівні використовуються такі інструменти, як терапевтичний моніторинг ліків, фармакогеноміка та точне дозування на основі моделі [1, 2].

Розробка ліків – це динамічний сектор, який постійно розвивається, у якому постійно з’являються передові та інноваційні тенденції. Фармацевтична галузь переживає фазу трансформації, що спричинена передовими технологіями та зміною потреб у сфері охорони здоров’я [3].

Мета. Метою дослідження був аналіз основних тенденцій розробки ліків та роль клінічної фармакології.

Матеріали та методи. Аналітичний та бібліосемантичний аналізи.

Результати дослідження. Сьогодні визначають декілька провідних тенденцій, які революціонізують розробку ліків: штучний інтелект, віртуальні клінічні випробування, потреба в глобальній вакцинації, цифрові технології охорони здоров’я та персоналізована медицина [3].

Штучний інтелект (ШІ) став найбільш революційною технологією, яка з’явилася за останні роки. Він має потенціал змінити форму величезних секторів світової економіки, і фармацевтична промисловість не є винятком. Штучний інтелект є корисним інструментом для оптимізації процесу розробки ліків і може допомогти компаніям-виробникам фармацевтичних препаратів заощадити час і гроші. Стандартний термін розробки ліків становить близько 12 років, і повторювані завдання, такі як відсіювання тисяч кандидатів у молекули ліків, забирають дорогоцінний час на дослідження. Використовуючи такі можливості штучного інтелекту, як міркування, пошук рішень і машинне навчання, вчені-фармацевтики можуть швидше звузити вибір ліків-кандидатів, тим самим ско-

рочуючи час і витрати на етапі відкриття ліків. Оскільки інвестиції в передові технології, такі як ШІ, стрімко зростають, багато фармацевтичних виробничих компаній уклали угоди з провідними компаніями штучного інтелекту. Наприклад, компанія фармацевтичної промисловості Sanofi співпрацює з багатонаціональною технологічною компанією Baidu для розробки ліків на основі ШІ. Не буде применшенням заявити, що штучний інтелект революціонізує процес розробки ліків, і враховуючи його потенціал для оптимізації процесів і економії коштів, ця тенденція не сповільнюється [3-5].

Сфера клінічних випробувань зазнала безпрецедентної трансформації під час пандемії COVID-19. На тлі прискорених досліджень вакцин і карантину по всьому світу фармацевтичні виробничі компанії змушені шукати нові та інноваційні способи проведення клінічних випробувань. Віртуальне клінічне випробування – це інноваційна дослідницька концепція, яка стосується випробувань, які збирають дані від учасників у нетрадиційних закладах охорони здоров'я. Цей цифровий підхід зменшує потребу в поїздках до клінічних центрів, а учасників можна націлити через сайти соціальних мереж і веб-платформи. Замість відвідування медичних закладів учасникам надсилають усі необхідні фармацевтичні субстанції та обладнання. Незважаючи на те, що клінічні дослідники відносно не використовують віртуальні клінічні випробування, популярність їх зростає.

Необхідність глобальної вакцинації. Швидке впровадження вакцин під час пандемії COVID-19 було неперевершеним науковим досягненням, але стало можливим лише завдяки співпраці регуляторів і фармацевтичних виробників. Усуваючи непотрібні договірні бар'єри, розширюючи та зміцнюючи ланцюжки постачання та враховуючи соціальні фактори, що стосуються коливань, фармацевтичні компанії можуть сприяти просуванню до глобальної справедливості щодо вакцин [3].

Цифрові технології охорони здоров'я стають все більш невід'ємною частиною процесу розробки ліків, особливо у віртуальних клінічних випробуваннях і дистанційному моніторингу пацієнтів. Передова технологія цифрової охорони здоров'я породила кілька інновацій, у тому числі переносні пристрої, мобільні програми для здоров'я та телемедицину. Багато людей уже використовують цифрові технології охорони здоров'я як частину свого повсякденного життя, будь-то для відстеження фізичної форми чи контролю сну. Можливості цифрових технологій охорони здоров'я дуже широкі: їх можна використовувати для збору точних персональних даних про здоров'я в режимі реального часу та розробки персоналізованої медицини. Хоча цифрові технології охорони здоров'я мають незліченну кількість переваг, є ряд недоліків, які фармацевтичні компанії повинні враховувати. Етичні дилеми щодо того, де зберігатимуться електронні медичні записи та особисті дані, викликають справжнє занепокоєння пацієнтів і дослідників, і деякі пацієнти можуть мати проблеми з використанням технологій загалом. Незважаючи на ці виклики, зростання цифрових

технологій охорони здоров'я є тенденцією, яка, ймовірно, продовжиться в найближчому майбутньому [3-7].

Персоналізована медицина (ПМ) передбачає, що ліки та медичні втручання мають бути адаптовані відповідно до індивідуальних потреб пацієнта. Потенційні переваги ПМ роблять її привабливим підходом до розробки ліків, наприклад, можливість призначати більш ефективні ліки, запобігати прогресуванню захворювання та прогнозувати сприйнятливість до захворювання. Існують різні способи, за допомогою яких фармацевтичні виробничі компанії можуть впроваджувати персоналізовану медицину в процеси розробки ліків. Наприклад, дослідник може захотіти перевірити, чи препарат, призначений для націлювання на конкретну генну мутацію, ефективний лише для пацієнтів, у яких є така мутація, і чи персоналізований лікарський підхід допоможе визначити пацієнтів, які, швидше за все, реагуватимуть на препарат. Його також можна використовувати для визначення оптимальної дози препарату для конкретного пацієнта та моніторингу відповіді пацієнта на терапію [3-7].

У сфері прецизійної медицини нові фенотипи та ендотипи хвороб, ймовірно, допоможуть визначити нові фармакологічні мішені, пацієнтів, які відповідають на терапію, і пацієнтів, які мають ризик побічних ефектів від лікарських засобів.

Висновки. Наведені тенденції означають рух до більш ефективного сектору розробки ліків, орієнтованого на пацієнта. Запровадження цих інновацій оптимізує роботу, сприятиме кращим показникам здоров'я та допоможе фармацевтичним компаніям адаптуватися до майбутніх викликів у сфері охорони здоров'я.

Список використаних джерел інформації:

1. What is Clinical Pharmacology? The American Society for Clinical Pharmacology and Therapeutics (ASCPT); [Available from: https://www.ascpt.org/Resources/Knowledge-Center/What-is-Clinical-Pharmacology
2. Liu Q., Ahadpour M., Rocca M., Huang S.M. Clinical Pharmacology Regulatory Sciences in Drug Development and Precision Medicine: Current Status and Emerging Trends. AAPS J. 2021.12;23(3): 54. doi: 10.1208/s12248-021-00563-3.
3. Top 5 Drug Development Trends of 2023. Назва з екрану: https://www.i-pharmconsulting.com/blog/top-drug-development-trends/
4. Boniolo F., Dorigatti E., Ohnmacht A.J. et al. Artificial intelligence in early drug discovery enabling precision medicine. Expert Opin Drug Discov. 2021;16(9):991-1007.
5. Nayarisseri A., Khandelwal R., Tanwar P. et al. Artificial Intelligence, Big Data and Machine Learning Approaches in Precision Medicine & Drug Discovery. Curr Drug Targets. 2021;22(6):631-655. doi: 10.2174/1389450122999210104205732.
6. Rongen G.A., Marquet P., van Gerven J.M.A.; EACPT research working group. The scientific basis of rational prescribing. A guide to precision clinical pharmacology based on the WHO 6-step method. Eur J Clin Pharmacol. 2021. 77(5): 677-683. doi: 10.1007/s00228-020-03044-2.
7. Cracowski J.L., Hulot J.S., Laporte S., Charvériat M., Roustit M., Deplanque D., Girodet P.O.; French Society of Pharmacology and Therapeutics (SFPT). Clinical pharmacology: Current innovations and future challenges. Fundam Clin Pharmacol. 2022. 36(3): 456-467. doi: 10.1111/fcp.12747.

STEREOCHEMICAL ASPECTS OF PHARMACODYNAMICS AND PHARMACOKINETICS OF MEDICINES

Podolskyi I. M., Podolskyi M. I.
National University of Pharmacy, Kharkiv, Ukraine
Department of Clinical Pharmacology
illya.podolsky@nuph.edu.ua

Introduction. From the point of view of stereochemistry, medicines can be divided into achiral, racemic, and pure enantiomers. Approximately 50 % of small molecules currently used in medicine are chiral, i.e. contain at least one center of asymmetry in their structure. Still, most of them are used as racemates, and only about 25 % as pure enantiomers. In the achiral environment, enantiomers have identical physical and chemical properties, but in the chiral environment, enantiomers with different pharmacological and pharmacokinetic properties can act as different medicines. Many chiral medicines are still used as racemates, although it has been proved that the pharmacokinetic and pharmacological profiles of enantiomers differ. Thus, it is known that the desired pharmacological effect is usually limited to only one of the enantiomers, called the eutomer, while the other enantiomer, called the distomer, has no effect on the pharmacological action or is even toxic. Thus, the strategy of "chiral switch", i.e. replacing a chiral medicine used as a racemate with its eutomer, has been considered a promising way to introduce safer and more effective medicines for some time.

Aim. To conduct a retrospective analysis of the phenomenon of "chiral switch" of medicines, focusing on successful recent examples.

Materials and methods. Analysis of literature sources on the subject, evaluation and systematization of the knowledge gained.

Results and their discussion. Living organisms consist of enantiomerically pure chiral substances, since in the living world all amino acids have an L-absolute configuration, and all carbohydrates have a D-absolute configuration. As a result, major physiological processes are stereoselective and use only one of all possible enantiomers. As a result, enantiomers can have different pharmacokinetic and pharmacological profiles, since the human body is a chiral environment (built of chiral structures: proteins, amino acids, enzymes, phospholipids), and the pharmacological activity of pharmaceuticals depends mainly on their interaction with various chiral targets, such as proteins (receptors, enzymes), nucleic acids (DNA, RNA) or biomembranes (phospholipids, glycolipids). One of the best-known examples of chiral switching is arylpropionic acid derivatives ("profenes") – anti-inflammatory drugs (ibuprofen, ketoprofen) that were introduced into medicine as racemates, except for naproxen, which is used in therapy as S-naproxen (R-naproxen is hepatotoxic). In the case of profenes, the S-enantiomers (dexibuprofen, dexketoprofen) are mainly responsible for the anti-inflammatory effect associated with cyclooxygenase-2 (COX-2) inhibition. However, *in vivo* metabolic bioconversion of the distomer (R-enantiomer) to the eutomer (S-enantiomer) occurs. Another important example of chiral switching is the

class of proton pump inhibitors (PPIs) (omeprazole, lansoprazole). PPIs were introduced into therapy as racemates, their chirality is the result of the presence of a chiral sulfur atom in their structure. Chiral switching of omeprazole to esomeprazole (S-enantiomer) was introduced based on the assumption that lower interindividual variability (slow and fast metabolizers) and average higher plasma levels would provide higher dose efficacy. In the case of the selective serotonin reuptake inhibitor (SSRI) antidepressant citalopram, the inhibition of serotonin reuptake is closely related to the activity of S-citalopram (escitalopram), which is more than 100 times more potent than R-citalopram. The use of S-ketamine (esketamine), which is a more effective analgesic and anesthetic due to antagonism with N-methyl-D-aspartate (NMDA) receptors, leads to shorter recovery after administration, better tolerability and reduced side effects (hallucinations and agitation). S-ofloxacin (levofloxacin) binds more efficiently to the enzyme DNA gyrase and topoisomerase IV than R-ofloxacin; S-ofloxacin has 2 times stronger antibacterial activity compared to racemate (ofloxacin), while R-ofloxacin is pharmacologically inactive. There are many more examples of successful chiral switching: bupivacaine-levobupivacaine, cetirizine-levocetirizine, albuterol-levalbuterol, amlodipine-levamlodipine, zopiclone-eszopiclone, and others.
However, there are also examples of unsuccessful attempts to introduce pure enantiomers (labetalol, satolol, fluoxetine). In these cases, as a rule, individual enantiomers have been shown to be less safe in clinical trials compared to the racemate.
Moreover, it was found that for many newly marketed single-enantiomer drugs, few randomized clinical trials (RCTs) were conducted that made direct comparisons with their racemic precursors for efficacy or safety endpoints. When single-enantiomer drugs were directly compared with their racemic precursors, most RCTs favored neither drug on the basis of efficacy or safety, and RCTs tended to be small and without well-defined endpoints. Although RCTs were identified favoring single-enantiomer drugs based on nonpredefined and secondary endpoints, results from predefined primary end points should take priority over other analyses. RCTs are typically powered to assess primary endpoints, and secondary analyses may be more susceptible to false positive findings. These findings raise concerns about the greater costs to the health care system incurred by chiral switching, without evidence to support benefit to patients, and the need for physicians and payers to encourage or require high-quality direct-comparison RCTs to inform the use of these more expensive therapeutics.

Conclusions. The potential therapeutic benefits of the chiral switching strategy are associated with an improvement in the therapeutic index due to increased selectivity and efficacy at receptors, reduced side effects, reduced interindividual variability in therapeutic response, reduced doses administered, improved pharmacokinetic profile, and reduced drug-drug interactions. However, today this approach is losing its relevance, which is primarily due to the current regulatory policy, which encourages exhaustive research of chiral molecules of new pharmaceuticals and allows the introduction of racemates only when it is proven that it has its advantages. Secondly, intensive research in this area has to some extent exhausted the issue of chiral switching of most “old” racemates. Patent conflicts are also an important aspect.

FACTORS CONTRIBUTING TO MEDICATION ADHERENCE IN MOROCCAN ADULT ASTHMA PATIENTS: CLINICAL AND PHARMACEUTICAL RESEARCH

Otrishko I. A., Bezugla N. P., Vietrova K. V., Mesbahi H.
National University of Pharmacy, Kharkiv, Ukraine
Department of pharmacology and clinical pharmacy
innaotrishko@gmail.com

Introduction. Asthma presents with coughing, breathlessness, wheezing, and tightness in the chest, is associated with airway inflammation and occasionally airflow obstruction. According the statistics of WHO, asthma affected an estimated 262 million people in 2019 and caused 455 000 deaths. Non-adherence to medications has many deleterious effects including poor treatment outcomes, increased economic burden, increased morbidity, hospitalization rate, health care utilization, productivity loss, and mortality.

Aim. To study the factors contributing to medication adherence in Moroccan adult asthma patients.

Materials and methods. General scientific and theoretical research methods were used.

Results and their discussion. WHO defines medication adherence as "the extent to which a person's behavior taking medication, corresponds with agreed recommendations from a health care provider". The major global challenge in asthma treatment and control is non-adherence to prescribed medications. Non-adherence to medications in asthmatic patients can be affected by different factors. These factors are classified by WHO into 5 categories: socioeconomic factors, factors associated with the health care team and system in place, disease-related factors, therapy-related factors, and patient-related factors. Effective asthma management requires that patients should have specific knowledge in order to control their asthma. Accordingly, patients should recognize: the pathophysiology of asthma and the lung changes underlying the disease, the role of different medications used in the treatment of asthma, self-monitoring and the use of inhalers and spacer devices, environmental controls (asthma triggers), and the signs and symptoms of uncontrolled asthma that may indicate an exacerbation. Adherence is also influenced by the patients' knowledge and beliefs about their disease and medications, in particular, associated with how the patients judge their personal need for the treatment (severity of disease), and potential adverse effects of medications.

Conclusions. People with asthma and their families need education to understand more about their asthma. This includes their treatment options, triggers to avoid, and how to manage their symptoms at home. It is important for people with asthma to know how to increase their treatment when their symptoms are worsening to avoid a serious attack. Healthcare providers (including pharmacists) may give an asthma action plan to help people with asthma to take greater control of their treatment.

COMPARATIVE CHARACTERISTICS OF MODERN GROUPS OF DRUGS AFFECTING THE RENIN-ANGIOTENSIN-ALDOSTERONE SYSTEM

Shchokina K. H.
Kharkiv Institute of Medicine and Biomedical Sciences, Kharkiv, Ukraine
Pharmacy department
acya@ukr.net

Introduction. according to the Global Health Estimates published by WHO, cardiovascular diseases are now the top ten leading causes of death. It is known that activation of the renin-angiotensin-aldosterone system (RAAS) plays a key role in the pathogenesis of many cardiovascular diseases. Long-term effects of increased production of renin, angiotensin II and sympathetic hypertension include the development of dyslipidemia, cardiac arrhythmias, hypercoagulation, endothelial dysfunction, insulin resistance, and weight gain. β-blockers, angiotensin-converting enzyme inhibitors (ACE inhibitors), angiotensin II receptor antagonists (ARA II), aldosterone antagonists are used to effectively counteract the negative effects of the RAAS. A direct renin inhibitor, aliskiren, has also been synthesized.

There are currently two ways to eliminate all the negative effects of angiotensin II: by reducing ACE activity and blocking type 1 angiotensin II receptors without disrupting ACE synthesis. The first goal is achieved by using ACE inhibitors, which have become the "gold" standard for treating chronic heart failure (CHF) over almost 30 years of successful clinical use, first-line drugs for treating hypertension, kidney disease, and diabetes. The second task is successfully solved with the help of ARA II. And each time the inevitable question arises: what is better to use in a specific clinical situation: ACE inhibitors or ARA II? In this regard, it was interesting to compare these two groups of drugs.

Aim. The aim of the work was to provide a comparative description of ACE inhibitors and ARA II, and also, based on evidence-based medicine data, to determine the clinical efficacy and safety of drugs in these groups.

Materials and methods. At the first stage of the study, using literature data, we conducted a comparative analysis of ACE inhibitors and ARA II. At the next stage, we analyzed their clinical efficacy and safety. To analyze the clinical efficacy and safety of ACE inhibitors and ARA II, we used well-known evidence-based medicine databases: Cochrane Library, Trip Database, PubMed.

Results. It has been established that each group of drugs has its own advantages and disadvantages. The uniqueness of ACE inhibitors is in providing a double therapeutic effect: they weaken the aggressive effect of angiotensin II (pressor link) and enhance the protective effect of bradykinin (depressor link). However, it is the bradykinin mechanism of action of ACE inhibitors, which plays an important role in the remodeling of the heart and blood vessels, that underlies the occurrence of cough (in 5-7% of patients receiving ACE inhibitors for a long time), which is sometimes the reason for stopping treatment. It is also known that ACE inhibitors cannot completely suppress excessive RAAS activity, since up to 70-80% of angiotensin II is

synthesized in organs and tissues without the participation of ACE using other enzymes (chymase, cathepsin, etc.), and during treatment with ACE inhibitors, the synthesis of angiotensin II can switch from the ACE-dependent pathway to the chymase pathway. This explains the possibility of the phenomenon of "escape" of the antihypertensive effect during treatment with ACE inhibitors, especially against the background of a high-salt diet, and serves as a rationale for the use of drugs that suppress the activity of angiotensin II, regardless of the pathway of its formation. In addition, ACE inhibitors affect the degradation of bradykinin, which, on the one hand, enhances their antihypertensive effect, and on the other hand, leads to the development of side effects (dry cough and angioedema).

The antihypertensive effect of ARA II does not depend on the activity of the RAAS, gender and age of the patient. ARA II does not have the phenomenon of "slippage" of the effect, since their action does not depend on the pathway of formation of angiotensin II. ARA II, having a similar mechanism of action with ACE inhibitors, can also affect the processes of left ventricular remodeling in patients with acute myocardial infarction (MI). However, ARA II do not affect the metabolism of bradykinin, which is associated with many positive effects of ACE inhibitors, in particular, organoprotective. Although it is assumed that it is possible that through stimulation of AT-2 receptors, ARA II can provide additional therapeutic effects, including organoprotective effects. In recent years, it has become known that AT-2 receptors are expressed only in case of damage, for example, in MI and CHF. It has been established that with the blockade of ATI receptors, the effect of angiotensin II on intact AT-2 receptors is enhanced, since its production compensatorily increases with the blockade of ATI receptors. It is possible that with the use of ARA II, reparative processes in tissues are activated through AT-2 receptors and remodeling with pathological proliferation of connective tissue structures is suppressed. Important advantages of ARA II include better tolerability due to the absence of influence on bradykinin formation; stable and longer-lasting inhibition of angiotensin II and aldosterone; additional positive effects associated with stimulation of angiotensin II receptors; the ability to block alternative pathways of angiotensin I transformation into angiotensin II; no effect on hematocrit; uricosuric effect (in some drugs). The main distinguishing feature of ARA II is high safety, a minimal number of side effects comparable to placebo, due to the absence of influence on bradykinin breakdown.

Conclusions. Thus, according to 10 systematic reviews, it was established that both groups of drugs that affect the RAAS are among the most effective and safe groups of drugs for the treatment of arterial hypertension and other cardiovascular diseases. From the point of view of evidence-based medicine, ACE inhibitors and ARA II do not have significant differences in pharmacodynamics, in particular, in the severity of the antihypertensive effect. Today, there is no clear answer to the question of the absolute advantages of any of these groups of drugs. It is certain that ACE inhibitors and ARA II are currently among the most effective and popular drugs.

EVALUATION OF CLINICAL AND PHARMACEUTICAL EFFICACY OF THE TRIAZOLE DERIVATIVES USE IN MODERN GYNECOLOGICAL PRACTICE

Bondariev Ye. V., Tymchenko Yu. V.
National University of Pharmacy, Kharkiv, Ukraine
Department of Clinical Pharmacology
clinpharmacol2023@gmail.com

Introduction. An urgent problem of modern gynecological practice is the effective use of effective drugs in the treatment of chronic inflammatory processes, in particular mycosis infection of the genital area in women of fertile age. In modern conditions, the requirements for the effectiveness and safety of rational therapy are constantly increasing, which improves the directions of effective treatment of patients Determination of etiopathogenetic justification and evaluation of data on efficacy and safety of use in clinical and pharmaceutical studies of a group of antifungal drugs in gynecological patients with manifestations of colpitis and vulvovaginitis of mycosis etiologywith manifestations of candidal pathology and provides appropriate conditions for the prevention of possible complications of the disease. Improving the efficacy and safety of certain groups of antimycosis drugs helps to increase the effectiveness of treatment of vulvovaginal candidiasis in women of fertile age, and also helps to minimize complications in this pathology. Doctors, pharmacists, clinical pharmacists in their practice are constantly improving the algorithms for the use of this group of drugs and preventing dangerous and negative manifestations of treatment.

Aim. Determination of etiopathogenetic justification and evaluation of data on efficacy and safety of use in clinical and pharmaceutical studies of a group of antifungal drugs in gynecological patients with manifestations of colpitis and vulvovaginitis of mycosis etiology.

Materials and methods. Conducting a non-interventional sampling study, clinical and pharmaceutical analysis and clinical processing of questionnaire data of patients with manifestations of colpitis and vulvovaginitis of mycosis etiology of different age groups, fixation of parameters of clinical and laboratory criteria.

Results and discussion. Evaluation of the results of this clinical and pharmaceutical analysis using a group of antifungal drugs, in particular itraconazole, in the treatment of chronic vulvovaginal candidiasis indicates that the main clinical aspect of antifungal therapy is the optimization of the effective and safe use of antimycosis drugs. Based on the data obtained, the main aspects of effective and safe use of the group of antimycosis drugs used to treat candida infection were determined. In general, chronic vulvovaginal candidiasis (CVC) is a "disease of civilization" that affects up to 75-77% of women of fertile age who have had vaginal candidiasis. The causative agent of CVC in 85% of cases is a mixed candida infection resistant to widely used antimycotic agents.

The most modern method of treating chronic mycosis infection is the use of systemic antifungal agents. The advantages of systemic therapy of colpitis and vulvovaginitis include: exposure to the pathogen of any localization and a significant anti-relapse effect. The property of uniform distribution into organs and tissues and the effect on the causative agent of any localization of the pathological process is most pronounced in itraconazole. Itraconazole is an antifungal agent derived from triazole. Its mechanism of action is associated with the ability to inhibit cytochrome P450-dependent enzymes of fungi sensitive to it, which leads to impaired ergosterol synthesis of the fungal cell wall. It has a wider spectrum of antifungal action on candida. The drug has a number of favorable pharmacodynamic and pharmacokinetic properties that increase its clinical efficacy. Itraconazole compares favorably with other representatives of azoles with a wider spectrum of action and much less toxicity. Based on the results of the study, the dynamics of symptoms and laboratory parameters of colpitis and vulvovaginitis in patients with chronic mycosis infection under the influence of itraconazole were assessed. It was determined that the dynamic efficacy rates of itraconazole used in the form of capsules ranged from 85% to 93% in patients with chronic vulvovaginal candidiasis. Based on the analysis of the dynamics of clinical manifestations of colpitis under the influence of itraconazole, positive tolerability was determined in 95% of patients. Therapy of chronic conditions is carried out for 6-7 days with a repeated course of treatment after 4-6 months, taking into account clinical and laboratory criteria.

Conclusions. When evaluating the data of the clinical and pharmaceutical study, the modern algorithm for the effective and safe use of itraconazole, a drug from the group of systemic antifungal drugs, for the treatment of gynecological patients with manifestations of chronic vulvovaginal candidiasis was substantiated. The use of itraconazole effectively ensures the eradication of all types of pathogenic candida, which makes it possible to empirically treat mycosis infection without prior determination of the subspecies of fungi. These results contribute to the improvement of modern criteria for the efficacy and safety of the use of a group of systemic antifungal drugs in the treatment of chronic mycosis infection, which makes it possible to increase the effectiveness of treatment in clinical gynecological practice.

CLINICAL EFFICACY AND SAFETY OF THE ANTIHISTAMINIC DRUG EBASTIN

Berezniakov A., Bondariev Ye.
National University of Pharmacy, Kharkiv, Ukraine
Department of Clinical Pharmacology
an-ber@ukr.net

Introduction. Antihistamines (AHP) are widely used in the treatment of allergic diseases such as allergic rhinitis, atopic dermatitis, urticaria, etc. The widespread use of AHP is due to their ability to reduce the intensity of allergic inflammation. The most common underlying mechanism of these diseases is an immediate allergic reaction (IgE mediated or type I). An allergic reaction, triggered by either immune or non-immune mechanisms, has both an early phase (developing within minutes) and a late phase (occurring 8-12 hours later). The late phase of the allergic response is clinically less pronounced than the early phase, but it is precisely this that leads to the chronicity of the allergic process.

Objective. Understanding the pharmacokinetics and pharmacodynamics, as well as having confidence in the predictability of clinical effects and the safety of H1-histamine receptor blockers, is crucial when choosing the appropriate medication for a specific clinical situation [1].

Materials and methods. First-generation antihistamines (with pronounced sedative and anticholinergic effects), used for the treatment of allergic diseases, are considered classical, whereas second generation antihistamines, which lack these side effects, are regarded as modern.

The pharmacological effects and therapeutic indications of first- and second-generation antihistamines are similar. However, unlike first-generation drugs, second-generation antihistamines contain hydrophilic fragments that reduce their lipophilicity, resulting in poor penetration through the blood-brain barrier, which minimizes central nervous system side effects. However, unlike the first generation, the second-generation drugs contain hydrophilic fragments that help reduce lipophilicity and, as a consequence, poor penetration through the blood-brain barrier, which minimizes central nervous system side effects [2]. The features of second-generation antihistamines include high affinity for H1 receptors, the absence of blockade of other receptor types, lack of tachyphylaxis, no dryness of mucous membranes or impaired mucus clearance, good absorption from the gastrointestinal tract, and no dependency on food intake. They also have a rapid onset of action, as well as sufficient duration of the main effect (up to 24 hours, and more than 48 hours for ebastine), which allows patients to take the medication in a highly compliant regimen – once per day. The chemical structure of ebastine significantly distinguishes it from many systemic antihistamines. What sets ebastine apart is that it does not form stereoisomers and enters the body as a pure substance rather than a mixture of racemates. This advantage ensures the highest affinity for H1 histamine receptors, duration of the clinical effect and a better safety profile by minimizing unwanted pharmacological effects and the absence of toxicity [3].

Results. Ebastine has high bioavailability and after oral administration is rapidly absorbed in the intestines (up to 95%) and extensively metabolized during the first pass through the liver into the pharmacologically active substance – karebastine [4]. Research data on the pharmacokinetics of ebastine demonstrate that its administration does not require dose adjustment depending on gender and age [5]. In both preclinical and clinical studies, ebastine has established itself as a drug with a favorable safety profile. First of all, the guarantee of safety is the absence of toxic effects associated with the medicinal substance, both general and specific (such as carcinogenicity, mutagenicity, and teratogenicity) [6]. Additional evidence of the safety of ebastine includes the absence of mutagenic potential and reproductive toxicity (teratogenicity, feto- and embryotoxicity, decreased fertility) even at doses exceeding 100 mg/kg in preclinical studies. The absence of carcinogenic activity of ebastine has been confirmed in studies using a dosage of 36 mg/kg/day, which is 200 times higher than the therapeutic dose for humans. The safety of long-term use of ebastine was observed in 6 studies involving 1,286 patients with chronic urticaria, of whom 567 received the medicine for a year at doses of 10 or 20 mg. It was noted that the recommended doses were well tolerated by the patients. Out of the reported 1,924 adverse events, 140 were serious, but only 15 could be presumably linked to the use of ebastine [7]. The safety of ebastine (and its pharmacologically active substance, karebastine) regarding its potential impact on cardiac electrophysiology has been thoroughly studied and evaluated in numerous in vitro and in vivo studies, including placebo-controlled, comparative trials and different age groups. It has been established that any changes in the QT/QTc interval associated with ebastine administration are not clinically significant [8].

Conclusion. Numerous studies characterize ebastine as one of the most promising second generation antihistamines with a favorable safety profile, making it a recommended option for managing various allergic reactions.

References:

1. Farzam K., Sabir S., O'Rourke M.C. Antihistamines. [Updated 2023 Jul 10]. In: StatPearls [Internet Resource]: https://www.ncbi.nlm.nih.gov/books/NBK538188/
2. Shabanov D.V., Lutkovskaya Ju.E. Actual Antihistamine Therapy. // Medical Council. 2020;(16):26–35. doi: 10.21518/2079-701X-2020-16-26-35.
3. Sophia Linton., Lubnaa Hossenbaccus., Anne K. Ellis. Evidence-based use of antihistamines for treatment of allergic conditions. // Ann. Allergy Asthma Immunol. 2023;131(4): 412–420. doi: 10.1016/j.anai.2023.07.019
4. Sastre J. Ebastine in the Treatment of Allergic Rhinitis and Urticaria: 30 Years of Clinical Studies and Real-World Experience. J. Investig. Allergol. Clin. Immunol. 2020;30(3):156-168. doi: 10.18176/jiaci.0401.
5. Van Cauwenberge P., de Belder T., Sys L. A review of the second-generation antihistamine ebastine for the treatment of allergic disorders // Expert Opin. Pharmacother. 2004; 5: 1807-1813.
6. Coltman N.J., Roberts R.A., Sidaway J.E.Data science in drug discovery safety: Challenges and opportunities. Exp Biol Med (Maywood). 2023 Nov;248(21):1993–2000. doi:10.1177/15353702231215890.
7. Roberts D. J. A preclinical overview of ebastine. Studies on the pharmacological properties of a novel histamine H1 receptor antagonist // Drugs. 1996; 52 Suppl 1: 8-14.
8. Demidova O. A., Shikh Ye. V., Ismagilov A. D., Sizova Zh. M. Clinical and pharmacological aspects of the safety of the use of drugs that cause QT prolongation // Bezopasnost' i risk farmakoterapii. 2017; 3 (5): 117-125.

ФАРМАКОТЕРАПІЯ ДИСБАЛАНСУ СИСТЕМИ ПРОТЕАЗИ АНТИПРОТЕАЗИ У ОСІБ ЗІ СТАБІЛЬНОЮ СТЕНОКАРДІЄЮ НАПРУГИ В ПОЄДНАННІ З ХРОНІЧНИМ ОБСТРУКТИВНИМ ЗАХВОРЮВАННЯМ ЛЕГЕНЬ

Крайдашенко О. О.

Запорізький державний медико-фармацевтичний університет
м. Запоріжжя, Україна
Кафедра внутрішніх хвороб 3
sashko_kr@ukr.net

Вступ. Серцево-судинна патологія в поєднанні з хронічним обструктивним захворюванням легень (ХОЗЛ) залишаються одними з найбільш поширених захворювань, що потребує сучасних підходів до патогенетичної терапії.

Мета. Оцінити клінічну ефективність терапії екзогенним L-аргіні-ном на фоні базисної терапії у хворих на стабільну стенокардію напруги (СН) ІІ-ІІІ ФК у поєднанні із ХОЗЛ ІІ-ІІІ стадії та вплив її на рівень експресії матрич-них металопротеїназ та активність тканинного інгібітору металопротеїназ.

Матеріали та методи. 38 хворих із СН+ХОЗЛ (середній вік 56,68±1,21 років), що були рандомізовані у 2 групи: 1-а включала 20 пацієнтів, котрі на тлі базисної терапії отримували метаболітотропний препарат L-аргінін (4,2% р-н в/в по 100 мл 1 раз на добу (курс 12 днів)) з подальшим переходом на розчин для перорального застосування по 10 мл 5 разів на добу; група 2 включала 18 хворих, які отримували тільки базисну терапію. Курс лікування 3 міс. ± 3 доби. Вивчалися показники функції зовнішнього дихання, толерантності до фізичних навантажень, якості життя, концентрації матричної металопротеїнази 9 (ММП-9) та її тканинного інгібітору (ТІМП) методом імуноферментного аналізу в крові.

Результати. Проведений аналіз рівня активності ММП-9 та ТІМР до та після проведеної терапії свідчив, що рівень експресії ММП-9 знизився на 41,2% ($p<0,05$) у 1 групі та на 14,8% у 2-й групі к кінцю строку терапії. В той час активність ТІМП через 12 тижнів лікування склала +31,1% ($p<0,05$) та +12,2%, відповідно. Причому кінцеві показники ММП-9 та ТІМП також достовірно відрізнялися між цими двома когортами, підкреслюючи більш статистично значне поліпшення протеаз-антипротеазого статусу при цієї коморбідності саме при застосуванні екзогенного L-аргініну. Все це супроводжувалось більш вираженими позитивними змінами показників функції зовнішнього дихання та толерантності до фізичного навантаження, якості життя у осіб 1-ї групи.

Висновки. Застосування L-аргініну на тлі базисної терапії у хворих на стабільну СН ІІ-ІІІ ФК у поєднанні із ХОЗЛ ІІ-ІІІ стадії сприяло більш вираженій позитивній динаміці як клінічної картини у хворих, так і модуляції порушень протеаз-антипротеазого статусу у вигляді зниження підвищеної експресії ММП-9 та елевації рівня ТІМП на тлі лікування.

ВИВЧЕННЯ ОСОБЛИВОСТЕЙ ПОКАЗНИКІВ ЯКОСТІ ЖИТТЯ ПАЦІЄНТІВ НА ФОНІ ЗАМІСНОЇ ГОРМОНАЛЬНОЇ ТЕРАПІЇ ГІПОТИРЕОЗУ

Отрішко І. А., Вєтрова К. В., Безугла Н. П., Ковальчук А. О.
Національний фармацевтичний університет, м. Харків, Україна
Кафедра фармакології та клінічної фармації
innaotrishko@gmail.com

Вступ. Гіпотиреоз є одним із найпоширеніших ендокринних розладів у всьому світі, частка гіпотиреозу становить 6,4 % у загальній структурі ендокринної патології. Забезпечуючи своєчасну діагностику та належне лікування гіпотиреозу, фахівці медицини та фармації мають змогу полегшити тягар особистих і соціально-економічних проблем, пов'язаних із цим розладом.

Мета. Вивчення особливостей показників якості життя пацієнтів на фоні замісної гормональної терапії гіпотиреозу.

Матеріали та методи. Методологічну основу дослідження складають принципи об'єктивності і системності. У роботі використано комплекс загальнонаукових та спеціальних методів: теоретичний, метод узагальнення, систематизації даних, порівняння, методи вивчення літературних джерел, аналізу та ін.

Результати. Метою лікування гіпотиреозу є нормалізація рівня тиреотропного гормону (ТТГ) і полегшення симптомів. Левотироксин є стандартом лікування гіпотиреозу. Замісна терапія має на меті усунення суб'єктивних та об'єктивних симптомів гіпотиреозу; досягнення нормалізації рівнів сироваткового ТТГ та гормонів щитоподібної залози; уникнення надмірного лікування (ятрогенного тиреотоксикозу), особливо у людей похилого віку. Левотироксин має застосовуватися за 30-60 хв до прийому їжі. Необхідно уникати прийому деяких препаратів (кальцію карбонат, заліза сульфат, алюмінію гідрохлорид) протягом 4 год. Оскільки замісну терапію левотироксином зазвичай призначають на невизначено тривалий термін, важливо передбачати ризики надмірного або недостатнього лікування в різні періоди життя пацієнта. На показники ефективності левотироксину можуть впливати наступні чинники: патофізіологічні; перебіг захворювання, генетичні особливості та супутня патологія або поведінкові (одночасний прийом інших лікарських засобів, пропуск прийому тощо); фармацевтичні – форма випуску, шлях введення, режим дозування, одночасний прийом із іншими ліками; порушення умов зберігання препарату, що погіршує стабільність лікарського засобу через шкідливий вплив вологості, світла і високої температури.

Висновки. Обізнаність щодо можливих проявів дисфункції щитоподібної залози, зокрема розвитку гіпотиреозу, сприятиме своєчасній діагностиці та призначенню адекватного лікування, дозволить уникнути більш важких проявів захворювання та порушень з боку роботи інших органів і систем, покращить якість життя таких пацієнтів.

CLINICAL PHARMACOLOGY FROM THE PERSPECTIVE OF THE WORLD HEALTH ORGANISATION

Seniuk I. V., Kravchenko V. M., Riyad Qamouta
National University of Pharmacy, Kharkiv, Ukraine
Department of Clinical Laboratory Diagnostics, Microbiology and Biological Chemistry
citochrom@gmail.com

Introduction. The World Health Organization (WHO) recently published a major position paper on clinical pharmacology 1. It drew ideas from the International Union of Basic and Clinical Pharmacology (IUPHAR) and the Council for International Organizations of Medical Sciences and complements Aronson's ”manifesto” by providing a vision for the specialty. This includes recognition of and contribution to global health, and leadership in the area of biologics and personalized medicines, in addition to the more traditional roles.

Aim. To explore strategies to ensure that clinical pharmacology is supported by teaching, training and policy organisations, to value clinical pharmacology services in monetary terms.

Materials and methods. The bibliographic databases used are Google Scholar, PubMed, Scopus, Web of Science, Springer and Science Direct.

Results. The proposals consist of three sections: engaging health system and policy makers to ensure the full benefit from clinical pharmacology services is achieved; putting a monetary value on clinical pharmacology; ensuring adequate recruitment of doctors into the specialty. “OMICS” is a neologism for disciplines, such as pharmacogenomics, epigenomics, transcriptomics, proteomics and metabolomics, that involve studies of cellular biological pathways. Choosing wisely is a programme initiated by the American Board of Internal Medicine that aims to promote conversations between clinicians and patients to choose care that is: supported by evidence; not duplicative of other tests or procedures already received; free from harm; truly necessary.

It aims to identify tests or procedures commonly used in specialty fields whose necessity should be questioned and discussed.

Clinical pharmacology is a research and teaching specialty and therefore these academic roles should be funded by Universities. However the discipline makes a large contribution to planning and delivery of clinical care at institutional regional and national levels. This has recently led to its recognition as a medical specialty in the European Union, a recognition that has existed in many Commonwealth countries for decades. Such recognition is a prerequisite for acceptance by other health professionals and governments or institutions, particularly when it comes to the payment for clinical pharmacology services.

At a clinical level the primary focus of clinical pharmacology must remain on quality, safety and patient-centred care but it is also important to emphasize the economic value of clinical pharmacology, recovered through reducing drug expenditures and the costs of unnecessary medications and avoidable adverse effects.

Service on hospital and regional drug committees, contribution to policy documents involving medicines, evaluation of literature to make better decisions, interactions with Industry around providing appropriate clinical support and information for particular drugs, education and training of prescribers, and advice and evaluation of regulatory and pricing documents are also important aspects of a clinical pharmacologist's workload that generate clear benefits to health systems.

Over the last 20 years we have witnessed the loss of training and consultant posts in clinical pharmacology around the world. Young clinicians have trained to be high level clinical pharmacologists but devoid of academic or clinical positions, have taken secure and well-paid jobs in Industry or other physician specialties. Hospital and university employment contracts do not compete well with those in Industry. The direct result of this is that when someone does retire there is often nobody capable and well trained to fill the position. There is a need to develop an all-embracing recruitment policy, and medical students interested in pharmacology, physiology, chemistry or with analytical or mathematical skills in particular should be approached early and mentored into clinical pharmacology posts. Specialist training in clinical pharmacology needs to be reviewed with links facilitated from residency and intern training wherever possible.

Strategies to ensure the vision becomes a reality. Ensure governments, health maintenance organizations (HMOs), consumer health groups and medical schools can see and measure the benefits a clinical pharmacologist adds to the healthcare and academic organization (improved patient care via correct choice and dose of drug, reduced numbers of adverse drug reactions and interactions, cost savings in pharmaceutical utilization and reducing inappropriate drug concentration testing and urine tests, enhanced design of clinical trials, correct interpretation of drug-related laboratory tests, stopping of futile or unhelpful therapies and teaching and guidance of clinical staff in therapeutics).

Develop a plan for a career structure that will attract medical students and young doctors to choose the specialty of clinical pharmacology to ensure the level of clinical service is expanded given an increasingly elderly population with multiple co-morbidities and exposed to polypharmacy. Ensure that development of new pharmacological agents involves the expertise of clinical pharmacologists at different stages of pre-clinical, clinical and post marketing development and use.

Conclusions. Clinical pharmacology is a medical specialty whose practitioners teach, undertake research, frame policy, give information and advice about the actions and proper uses of medicines in humans and implement that knowledge in clinical practice. It involves a combination of several activities: drug discovery and development, training safe prescribers, providing objective and evidence-based therapeutic information to ethics, regulatory and pricing bodies, supporting patient care in an increasingly subspecialized arena where co-morbidities, polypharmacy, altered pharmacokinetics and drug interactions are common and developing and contributing to medicines policies for Governments. Clinical pharmacologists must advocate drug quality and they must also advocate for sustainability of the Discipline. However for this they need appropriate clinical service and training support.

МЕТОДИКИ КОРЕКЦІЇ ПОСТТРАВМАТИЧНОГО СТРЕСОВОГО РОЗЛАДУ

Кахніашвілі А. С., Деримедвідь Л. В.
Національний фармацевтичний університет, м. Харків, Україна
Кафедра фармакології та клінічної фармації
kahniashvilinastya@gmail.com

Вступ. Посттравматичний стресовий розлад (ПТСР) – це гострий або хронічний розлад психічного здоров'я, який розвивається після ситуації, що загрожувала життю та цілісності людини. Цей стан розвивається не одразу після травматичної події, а через певний проміжок часу, який може варіюватися залежно від психічного стану людини. Сьогодні випадки ПТСР на території України частішсе зустрічаються внаслідок воєнних дій, окупації, масованих обстрілів та підвищення рівня злочинності на деокупованих територіях. Отже, ця проблема є актуальною для всіх українців сьогодні та для більшості громадян у майбутньому.

Мета. Розглянути та дослідити методики лікування та корекції ПТСР; визначити фармакологічні групи препаратів, які використовується за даного розладу та виокремити декілька лікарських засобів, які можуть виступати препаратами першої лінії.

Матеріали та методи. Було проаналізовано наукові статті, рекомендації Міністерства охорони здоров’я України, наукову літературу, рекомендації закордонних лікарів.

Результати. У результаті дослідження було встановлено, що ПТСР піддається лікуванню, але чіткі часові межі встановити важко, оскільки психічне здоров’я у кожної людини індивідуальне і терапія може тривати від 3 місяців до року, або більше. Наразі виділяють два основних підходи до лікування – психотерапія та фармакотерапія. До першого відносять багато методик, а саме: раціональна психотерапія, аутотренінг, когнітивна психотерапія, особистоорієнтована терапія, гештальт-терапія, логотерапія, групова терапія та ін. Усі ці методики допомагають подолати причини та усунути ПТСР у пацієнтів, які цим страждають, а також запобігти виникненню цього розладу у людей із групи ризику. Що стосується фармакотерапії, то її ефективність, без залучення психотерапії, становить 60 %. Приблизно третина пацієнтів видужують шляхом медикаментозного лікування. Згідно з сучасними протоколами лікування ПТСР, в терапії широко використовують антидепресанти з групи селективних інгібіторів зворотного захоплення серотоніну (СІЗЗС) – пароксетин та сертралін. СІЗЗС пригнічують зворотне захоплення серотоніну (5НТ) у нейронах; а деякі препарати також мають слабкий ефект відносно зворотного захоплення у нейронах норадреналіну та допаміну. СІЗЗС використовують як препарати першої лінії через їх позитивні ефекти, які усувають багато симптомів ПТСР, а саме: нав’язливі думки та спогади, дратівливість, гіперзбудження, пригнічений настрій та ін.

Варто зазначити, що дана група препаратів викликає синдром відміни при передчасному завершенні лікування, і через це виникає помилкове судження про загострення симптомів ПТСР.

Також до ефективних препаратів з групи СІЗЗС для терапії ПТСР відносять і флуоксетин, як препарат лікування за принципом off-label.

У разі неефективності СІЗЗС, використовують інгібітори зворотного захоплення серотоніну-норадреналіну (ІЗЗСН) як препарати другої лінії. Венлафаксин є ефективним у лікування ПТСР, у той час, як міртазапін, нефазодон та трициклічні антидепресанти також мають ефективність, але не застосовуються у терапії через велику кількість побічних ефектів, проте також належать до препаратів другої лінії. Максимальне дозування антидепресантів на добу: флуоксетин – 80 мг, пароксетин – 50 мг, сертралін – 200 мг, венлафаксин – 300 мг.

При порушеннях сну, спричинені кошмарами, доцільне використання альфа-адреноблокатору празозину у максимальному дозуванні 40 мг на добу. Снодійні препарати використовуються короткочасно для усунення симптомів безсоння. Максимальне дозування снодійних на добу: залеплон – 20 мг, золпідем – 10 мг. Препарати бензодіазепінового ряду використовувати небажано, оскільки вони погіршують симптоми ПТСР, тому їх слід уникати.

Наразі в терапії ПТСР обмежено використовують препарати літію, вальпроат натрію, топірамат, габапентин та ін. препарати другої лінії через вимоги до моніторингу її ефективності.

Антипсихотичні препарати за ПТСР вивчалися в багатьох контрольованих дослідженнях, проте велике багатосайтове дослідження рисперидону не виявило користі для лікування ПТСР. Саме тому вони не рекомендуються в лікуванні.

Також є низка рекомендації для медикаментозної терапії у певних вікових категорій пацієнтів: дітям та підліткам до 18 років не рекомендовано лікування препаратами при ПТСР; якщо у пацієнта наявна депресія, то потрібно лікувати спочатку її; при наркотичній або алкогольній залежності все одно потрібно лікувати ПТСР; якщо у пацієнта складний випадок ПТСР, терапія буде тривати довше; за наявності психозу у пацієнта доцільне використання рисперидону, а інших антипсихотичних засобів слід уникати.

Висновки. Зважаючи на тяжкість перебігу посттравматичного стресового розладу, фахівці в галузі медицини та фармації змогли досягти значних успіхів у лікуванні даного порушення з досягненням повного одужання. Методи корекції можуть бути у вигляді психотерапії та фармакотерапії, що мають значні успіхи та позитивну динаміку у більшості хворих.

Препаратами для безпечного лікування ПТСР можуть бути з наступних груп: антидепресанти (пароксетин, сертралін, венлафаксин), альфа-адреноблокатори (празозин), снодійні (залеплон, золпідем). Окремо застосовуються при ускладненнях бензодіазепіни, антипсихотичний засіб рисперидон, препарати літію, вальпроат натрію, топірамат, габапентин. Деякі препарати можуть використовуватися за принципами off-label.

КЛІНІЧНА ФАРМАКОЛОГІЯ НА ОСНОВІ ІНТЕРФЕРЕНЦІЇ РНК

Кравченко В. М., Сенюк І. В., Галузінська Л. В.
Національний фармацевтичний університет, м. Харків, Україна
Кафедра клінічної лабораторної діагностики,
мікробіології та біологічної хімії
citochrom@gmail.com

Вступ. Протягом багатьох десятиліть розробка лікарських засобів була зосереджена на протеїнах, що викликають захворювання, наприклад, протеїнах патогенних мікроорганізмів та протеїнах з аномальними функціями або рівнями експресії. Однак ні низькомолекулярні препарати, ні протеїнові терапевтичні препарати не впливають безпосередньо на основні гени, які є першопричиною багатьох захворювань людини. На додаток до цього внутрішнього обмеження, малі молекули та протеїнові терапевтичні засоби не можуть отримати доступ до значної кількості мішеней клітин. Знання про те, як генетичні агенти перетворюються на вихідні біологічні зсуви, сприяло появі концепції генної терапії. Фундаментально відрізняючись від стратегії модуляції протеїнової мішені, генна терапія змінює послідовність ДНК, яка є причиною патології.

З моменту першого нормативного схвалення у 1990 році застосування генної терапії в медичних втручаннях дедалі більше набирало обертів, до прикладу винаходом кластеризованого регулярно проміжного короткого паліндромного інструменту повторення в останнє десятиліття. Хоча генна терапія на основі ДНК має потенціал для лікування захворювань за допомогою маніпуляцій з геномом, потенціал нецільового редагування може призвести до незворотних руйнівних наслідків.

РНК-таргетна терапія, яка часто спрямована на зміну експресії мРНК для запобігання синтезу спорідненого проблемного протеїну, навпаки, не тільки безпосередньо модулює біологічну функцію на посттранскрипційному рівні без незворотного перегляду геному людини, але й обходить проблему розробки фармакологічного агента з достатньою специфічністю і спорідненістю до цільового протеїну. На сьогоднішній день Управління з контролю за якістю харчових продуктів і медикаментів (FDA) схвалило п'ять терапевтичних засобів малої інтерферуючої РНК (міРНК). Терапевтичні препарати міРНК, часто довжиною 20-30 нуклеотидів, використовують протеїн людського аргонавта 2 для формування РНК-індукованого комплексу глушіння, який, у свою чергу, забезпечує розщеплення цільової мРНК.

Мета. Проаналізувати та узагальнити клінічні фармакологічні дані, які схвалені FDA терапевтичних засобів на основі міРНК, спрямованих на полегшення майбутньої розробки лікарських засобів.

Матеріали та методи. Було використано інформацію про призначення та клінічні фармакологічні огляди FDA з Drugs@FDA. Проаналізована загальнодоступна література (PubMed), яка стосується клінічної фармакології з використанням міРНК.

Результати. Усі п’ять схвалених міРНК, до яких належать *Patisiran* (2018), *Givosiran* (2019), *Lumasiran* (2020), *Inclisiran* (2021), *Vutrisiran* (2022) – це послідовності дволанцюгові РНК, які складаються з 21-нуклеотидного кодуючого ланцюга та 21-23-нуклеотидного некодуючого ланцюга. У цитоплазмі два ланцюги розкручуються, і некодуючий ланцюг інтегрується в РНК-індукований комплекс сайленсингу. міРНК метаболізуються нуклеазами до олігонуклеотидів меншої довжини, які можуть включати деякі фармакологічно активні види. За винятком лікарського препарата *Patisiran* схвалені міРНК кон’юговані з N-ацетилгалактозаміном для ефективної доставки до печінки, оскільки рецептор азіалоглікопротеїну, який розпізнає специфічний фрагмент, в основному експресується у гепатоцитах. Зазначені міРНК використовують механізм РНК-інтерференції та регулюють каталітичне розщеплення мРНК. Крім того, чотири кон’юговані з N-ацетилгалактозаміном міРНК вводять підшкірно, тоді як *Patisiran* вводять внутрішньовенно.

Кон’юговані з N-ацетилгалактозаміном міРНК не взаємодіють з ензимами та транспортерами, що метаболізують лікарські препарати, оскільки вони ефективно розподіляються в цільовій тканині та не модулюють цитокіни. У дослідженнях *in vitro* ці п’ять міРНК, схвалених FDA, не функціонували як інгібітори, індуктори або субстрати ферментів цитохрому P_{450} або інгібітори та субстрати транспортерів. міРНК можуть індукувати вроджені імунні відповіді та можуть бути обмежено імуногенними.

З огляду на те, що міРНК в основному розщеплюються ендонуклеазами та екзонуклеазами, не є субстратами для ферментів P_{450} і мають короткий період напіврозпаду у плазмі, загалом не було значної дії щодо клінічно значущого впливу печінкової недостатності (незалежно від ступеня) на метаболізм затверджених міРНК. З іншого боку, було зазначено, що на рівні експресії можуть суттєво впливати захворювання печінки, такі як цироз і гепатоцелюлярна карцинома.

У пацієнтів з помірними порушеннями функції печінки вихідні рівні біомаркерів були нижчими, а ефекти фармакодинаміки меншими, ніж ті, що спостерігалися у пацієнтів із нормальною функцією печінки. Для пацієнтів із легким та помірним порушенням функції печінки корекція дози не актуальна. Фармакокінетичний аналіз даних, отриманих у спеціальному дослідженні ниркової недостатності, довів про підвищення максимальної (пікової) концентрації у сироватці крові (C_{max}) у пацієнтів з легким, помірним або тяжким порушенням функції нирок порівняно з пацієнтами з нормальною функцією нирок.

Висновки. міРНК як терапевтичні агенти є зміною парадигми у галузі розробки ліків. Оцінка унікальної ефективності, безпеки та клінічних фармакологічних характеристик цього нового класу терапевтичних засобів активно проводиться. Клінічні фармакологічні характеристики терапевтичних засобів на основі міРНК, схвалених FDA, є унікальними порівняно з характеристиками малих молекул і терапевтичних засобів на основі протеїну.

ТЕРАПЕВТИЧНИЙ ПОТЕНЦІАЛ, БЕЗПЕКА І МІЖЛІКАРСЬКІ ВЗАЄМОДІЇ ROSMARINUS OFFICINALIS L.

Савельєва-Кулик Н. О.

Національний медичний університет імені О.О. Богомольця, м. Київ, Україна

Кафедра клінічної фармакології та клінічної фармації

nsavelyevakulyk@gmail.com

Вступ. Рослинні лікарські засоби використовувались з давніх часів. Останніми десятиліттями увага до досліджень фітозасобів зросла, враховуючи їх відносно обмежений спектр побічних ефектів і ускладнень терапії. Розмарин, *Rosmarinus officinalis L. (Labiatae),* застосовують для полегшення головного, ревматичного і спастичного болю, при дисменореї, епілепсії, нервовому збудженні, істерії, депресивних розладах, зниженні пам'яті та когнітивних розладах (Duke, 2000; Heinrich et al., 2006). Рядом досліджень на тваринних моделях або культивованих клітинах описано протизапальні (Kayashima et al., 2012), антиоксидантні, антиноцицептивні, нейропротекторні (González-Trujano et al., 2007; Bakirel et al., 2008; Hou et al., 2012), а також антидепресивні та прокогнітивні ефекти розмарину та його сполук (Hosseinzadeh et al. 2004; Sasaki et al., 2013). За даними фітохімічних досліджень, з-поміж терпеноїдів, ефірних олій, алкалоїдів та флавоноїдів, які містить розмарин, найпотужнішими ефективними компонентами визнано тритерпени, фенольні дитерпени і фенольні кислоти, включаючи розмаринову, карнозинову, урсолову, бетулінову кислоти, розманол і карнозол (Hosseinzadeh et al., 2003; Borras-Linares et al., 2014). Серед згаданих фенольних сполук розмаринова та карнозинова кислоти виявляють найбільший лікувальний вплив, а саме протизапальний та антиоксидантний (Jayanthy et al., 2014; Birtic et al., 2015; Sedighi et al., 2015). FDA США розмарин класифіковано як «загалом безпечний», або GRAS (CFR182.10; 182.20) (Aguilar et al., 2008), і визначено, що розмаринова кислота має дуже слабку токсичність з LD50 561 мг/кг у гризунів (Tal Friedman, 2015).

Мета. Оцінити терапевтичний потенціал, безпеку застосування і ризик міжлікарських взаємодій Rosmarinus officinalis L. на основі даних клінічних випадків, оглядів, метааналізів, досліджень *in vitro* та *in vivo*.

Матеріали та методи. Аналіз інформації за ключовими словами "Rosmarinus officinalis", "rosemary", "therapeutic effects", "toxicity", "drug interactions" в наукометричній системі PubMed за період 1999-2024 рр.

Результати. Результатами проведеного аналізу встановлено, що розмарин традиційно використовується як болезаспокійливий і протизапальний засіб (al-Sereiti et al., 1999; Heinrich et al., 2006). Останніми роками активно вивчаються його цитостатичні (González-Vallinas et al., 2015; Allegra et al., 2020) і гепатопротекторні властивості (Guimarães et al., 2023), а також повідомляється, що розмарин виявляє високу антиоксидантну та протимікробну активність (Nieto et al., 2018; Uluışık, 2023), має нейропротекторний і кардіопротекторний

вплив (Rahbardar and Hosseinzadeh, 2020; 2024). Визнано, що захисні ефекти розмарину та його компонентів опосередковані пригніченням явищ окисного стресу, зниженням рівня медіаторів запалення, включаючи фактор некрозу пухлини-альфа (TNF-α), інтерлейкін-6 (IL-6), інтерлейкін-17 (IL-17), циклооксигеназу-2 (ЦОГ-2) і ядерний фактор кВ (NF-кВ), а також модуляцією сигнальних шляхів апоптозу та мітоген-активованої протеїнкінази (Alavi et al., 2021).

За даними Європейської агенції з лікарських засобів (EMA, 2010), ефірну олію розмарину рекомендовано в лікуванні пацієнтів з диспепсією та легкими спазматичними розладами, як допоміжний засіб для полегшення незначного м'язового і суглобового болю, при незначних розладах периферичного кровообігу. Незважаючи на те, що загалом розмарин вважається безпечним для системного та місцевого призначення, у чутливих осіб можуть виявлятись ознаки дерматиту та алергічних реакцій. Встановлені можливі токсикологічні ефекти охоплюють прояви гепато- і нефротоксичності: атрофічні і дегенеративні зміни, підвищення рівня азоту сечовини, аспартатамінотрансферази і зменшення вмісту загального білка сироватки крові (Motaghi et al., 2021). Розмарин може викликати репродуктивну токсичність шляхом зниження сперматогенезу, рівня тестостерону, провокувати генотоксичність та аномалії у плодів, посилюючи скорочення цитоплазматичної мембрани, утворення апоптичних тілець і фрагментації міжнуклеосомної ДНК (Nusier et al., 2007; Bezerra et al., 2022).

Встановлено ряд взаємодій рослинних лікарських засобів на основі розмарину з антибіотиками, анксіолітиками та антикоагулянтами з можливістю потенціювання їх активності (Ulbricht et al., 2010; Posadzki et al., 2013). Водночас продемонстровано здатність компонентів ефірної олії розмарину індукувати каталітичну функцію мікросомальних ферментів, зокрема CYP_{2B}, та меншою мірою – UDP-глюкуронозилтрансферази (UGT), а отже і можливість взаємодії з препаратами, біотрансформація яких відбувається за участю цитохромоксидаз CYP_{450} або UGT (Debersac et al., 2001). Індукуючий вплив розмарину на ізоферменти CYP_{2E1}, CYP_{1A2}, CYP_{3A4} (Yaneva et al., 2020) потенційно може підвищувати плазмовий кліренс субстратів вищезгаданих ізоферментів (теофілін, мексилетин, кломіпрамін, клозапін тощо) або навпаки сприяти ризику розвитку токсичних ускладнень у разі утворення вихідними субстратами проміжних токсичних метаболітів, як-от при одночасному застосуванні парацетамолу (Zhou et al., 2003).

Висновки. Ймовірні механізми дії та потенційні міжлікарські взаємодії розмарину необхідно досліджувати, поєднуючи фізіологічні та фармакологічні методи, орієнтовані на відстеження біодоступності. При тривалому та інтенсивному використанні засобів на основі розмарину бажано дотримуватись обережності, зважаючи на можливість нефротоксичного і гепатотоксичного ефектів, а також небажаного впливу на репродуктивну систему і ризик тератогенезу. В цілому, все вказане підкреслює важливість враховувати потенційні міжлікарські взаємодії, призначаючи комплексне лікування, до складу якого входять фітозасоби із вмістом Rosmarinus officinalis L.

ОБІЗНАНІСТЬ ФАРМАЦЕВТІВ ЩОДО КЛІНІЧНОЇ ХАРАКТЕРИСТИКИ ТРИВОЖНИХ РОЗЛАДІВ ТА НАДАННЯ ФАРМАЦЕВТИЧНОЇ ОПІКИ ПРИ ЛІКУВАННІ ПОРУШЕНЬ ДІЯЛЬНОСТІ ЦНС

Штробля А. Л., Гольцева С. С.
Ужгородський національний університет,
м. Ужгород, Україна
anzhela.shtroblia@uzhnu.edu.ua, mf.holtseva.snizhana@student.uzhnu.edu.ua

Вступ. Тривожні розлади є однією з найпоширеніших форм психічних захворювань, що мають значний вплив як на здоров'я окремих осіб, так і на суспільство загалом. За даними епідеміологічних досліджень, приблизно 25% дорослих людей хоча б раз у житті стикаються з тривожними станами, що підкреслює масштаб і серйозність цієї проблеми. Тривога, будучи природною емоційною реакцією на потенційні загрози, виконує адаптивну функцію, але у випадках надмірного або тривалого впливу стресорів вона може перетворитися на патологічний стан. Це значно погіршує якість життя людини, впливаючи на її фізичне, емоційне і когнітивне благополуччя, а також на соціальну активність. Тривожні розлади часто супроводжуються іншими психічними захворюваннями, зокрема депресією, а також можуть бути асоційовані з підвищеним ризиком суїцидальної поведінки та зловживання психоактивними речовинами. Така коморбідність ускладнює лікування, вимагаючи комплексного підходу, що включає не лише медикаментозну терапію, але психологічну підтримку.

Значну роль у лікуванні тривожних розладів відіграє фармацевтична опіка, яку надають фармацевти при зверненні пацієнта в аптеку. Пацієнти часто використовують безрецептурні препарати (ОТС) для зняття симптомів тривоги та стресу, однак неконтрольоване застосування цих засобів може призводити до негативних наслідків, таких як передозування, звикання або розвиток побічних ефектів. У зв'язку з цим фармацевти відіграють ключову роль у консультуванні пацієнтів щодо безпечного використання таких препаратів.
Забезпечення належної фармацевтичної опіки сприяє покращенню клінічних результатів та якості життя пацієнтів, допомагаючи уникнути ризиків, пов'язаних із неправильним використанням ліків.

Мета. Оцінити рівень обізнаності працівників аптек щодо клінічних характеристик тривожних розладів і надання фармацевтичної опіки пацієнтам при лікуванні симптому тривоги.

Матеріали та методи. У дослідженні використовувалося анонімне опитування фармацевтів, яке складалося з 27 запитань. Анкета охоплювала ключові аспекти клінічної характеристики тривожних станів, симптоми, фактори ризику, а також принципи використання безрецептурних препаратів для лікування симптомів тривоги. Усього в дослідженні взяли участь 40 фармацевтів. Для аналізу результатів використовувалися кількісні методи, з розподілом правильних і неправильних відповідей у відсотковому співвідношенні.

Результати та їх обговорення. Результати опитування показали високий рівень обізнаності фармацевтів щодо клінічних характеристик тривожних станів і методів їх лікування. Основні показники:

1. Визначення тривожних станів: 92,5% фармацевтів правильно визначили тривогу як нав'язливе почуття тривоги або загрози.
2. Розуміння симптомів тривоги: 85% респондентів правильно ідентифікували основні симптоми тривожних станів, такі як прискорене серцебиття, безсоння, нервозність.
3. Знання про ОТС-препарати: 92,5% були обізнані щодо правильного використання безрецептурних засобів, зокрема, про можливі побічні ефекти та ризики передозування.

Також було виявлено високий рівень знань про анксіолітики (87,5%), препарати на основі звіробою (92,5%) та гліцин (87,5%).

Дослідження підтвердило, що фармацевти відіграють ключову роль при наданні фармацевтичної опіки пацієнтам із тривожними розладами. Високий рівень їхньої обізнаності щодо клінічних характеристик тривожних станів дозволяє їм не лише надавати кваліфіковані рекомендації щодо ОТС-препаратів, але й запобігати можливим ускладненням, пов'язаним із їх неконтрольованим застосуванням.

Фармацевти здатні своєчасно розпізнати серйозні симптоми, що вимагають негайної медичної допомоги (77,5% респондентів правильно відповіли на питання про загрозливі симптоми). Це свідчить про високий рівень готовності фармацевтів надавати допомогу в рамках фармацевтичної опіки пацієнтам із тривожними розладами, що робить їх важливими учасниками системи охорони здоров’я.

Висновки. Результати дослідження свідчать про високий рівень обізнаності фармацевтів щодо тривожних розладів і надання фармацевтичної опіки. Це забезпечує безпеку та ефективність лікування пацієнтів із порушеннями діяльності ЦНС, підвищуючи якість фармацевтичної допомоги. Важливим аспектом є те, що фармацевти добре розуміють симптоматику тривожних станів, фактори ризику та правильне застосування безрецептурних препаратів, що сприяє покращенню здоров'я населення та зменшенню негативних наслідків, пов'язаних із неконтрольованим застосуванням ліків.

Підвищення рівня обізнаності фармацевтів щодо тривожних розладів та методів лікування є важливим кроком для покращення якості фармацевтичної опіки та покращення якості життя пацієнтів.

ТОПІРАМАТ ДЛЯ ПРОФІЛАКТИКИ МІГРЕНІ

Гриньків Я. О.
Львівський національний медичний університет імені Данила Галицького
м. Львів, Україна
Кафедра організації і економіки фармації, технології ліків та фармакоекономіки ФПДО
yaryna_hrynkiv@ukr.net

Вступ. Препарати для профілактики можуть зіграти важливу роль у зменшенні частоти, важкості та тривалості нападів мігрені, а також потенційно скоротити надмірне використання лікарських засобів при терапії власне фази головного болю.

Мета. Опрацювати джерела інформації щодо ефективності топірамату для профілактики мігрені.

Матеріали та методи. Матеріалами дослідження були офіційно затверджені інструкції до застосування топірамату; публікації у фахових та наукових виданнях. Методи дослідження – вебометричний та порівняльний аналізи, систематизація даних.

Результати: Ефективність топірамату при його застосуванні для профілактики нападів мігрені підтверджена результатами цілої низки клінічних досліджень. Топірамат має кілька механізмів дії, які можуть сприяти профілактиці мігрені: інгібування потенціалзалежних натрієвих каналів, інгібування активованих високою напругою кальцієвих каналів, інгібування опосередкованої глутаматом нейротрансмісії на α-аміно-3-гідрокси- підтипи рецепторів 5-метилізоксазол-4-пропіонової кислоти та каїнату, а також посилення потоку хлориду, опосередкованого рецептором $_{A}$ γ-аміномасляної кислоти.

Для профілактики нападів мігрені у дорослих рекомендована добова доза топірамату становить 100 мг, розподілена на 2 прийоми. Слід розпочинати з прийому 25 мг ввечері протягом тижня. У подальшому дозу збільшувати на 25 мг на добу з інтервалом в тиждень після кожного підвищення дози. Можна застосувати менш значне збільшення дози або триваліші інтервали між збільшеннями, якщо пацієнт погано переносить зазначений режим добору дози. При підборі дози слід керуватися терапевтичною ефективністю.

Вивчення профілю безпеки топіромату засвідчило хорошу переносимість препарату хворими на мігрень. Найчастішими побічними ефектами були слабкість, нудота, відсутність апетиту та парестезії. Ризик розвитку побічних ефектів залежить від дози препарату та є максимальним при використанні дози 200 мг на добу. Найбільша частота побічних ефектів відзначається у перші два місяці терапії, при тривалому лікуванні показник поступово знижується

Висновки. Численні джерела літератури наводять переконливі дані про доцільність застосування топірамату для профілактики мігрені у дорослих.

ВИКОРИСТАННЯ ІНФУЗІЙНИХ РОЗЧИНІВ ПРИ ГІПОКАЛІЄМІЇ У ПАЦІЄНТІВ ЗАГАЛЬНОЇ ХІРУРГІЇ

Карпінчик Л. С., Харченко Ю. В.

Дніпровський державний медичний університет, м. Дніпро, Україна
Кафедра фармакології, загальної та клінічної фармації
209_14@dmu.edu.ua

Вступ. Недостатні абсорбція та споживання, а також надмірне виведення (блювота, пронос) іонів K^+ через шлунково-кишковий тракт (ШКТ) є частою причиною гіпокаліємій у пацієнтів загальної хірургії. Тому, гіпокаліємія є поширеною проблемою в загальній хірургії, що може призводити до серйозних ускладнень, таких як аритмії, м'язова слабкість та порушення функцій органів. Для відновлення рівня внутрішньоклітинного K^+ та у плазмі крові проводять інфузії розчином ГІК (глюкоза-інсулін-калій).

Мета. Проаналізувати доцільність та ефективність використання інфузійного розчину ГІК при гіпокаліємії у хворих загальної хірургії внаслідок розладів шлунково-кишкового тракту.

Матеріали та методи. Проведено аналіз сучасної літератури присвяченої методам відновлення рівня іонів K^+ у плазмі крові та в клітинах за допомогою інфузії. Виконано систематичний пошук у наукових базах даних «National library of medicine», «Google Scholar», «PubMed».

Результати. Основними захворюваннями загальної хірургії є захворювання ШКТ, які спричиняють дефіцит іонів K^+ в організмі, а також енергетичне голодування, за рахунок неможливості в першу чергу надходження та абсорбції клітинами тонкого кишечника іонів та глюкози (служить джерелом енергії для клітин і стимулює вивільнення інсуліну з підшлункової залози), тому варто застосувати внутрішньовенний спосіб відновлення. Наслідком нестачі калію у плазмі крові також є порушення процесу транспорту іонів Na^+ та H^+ в ентероцитах. ГІК є засобом, що сприяє підтриманню належного внутрішньоклітинного і позаклітинного рівня калію, який є основним внутрішньоклітинним іоном, що відіграє важливу роль у клітинному метаболізмі і регуляції різних функцій клітин, зокрема процесі реполяризації. Іони калію активують велику кількість цитоплазматичних ферментів, регулюють внутрішньоклітинний осмотичний тиск. Інсулін, в свою чергу, сприяє переносу калію з позаклітинного простору в клітини, що допомагає знизити рівень калію в плазмі крові. Режим дозування та термін застосування встановлюють індивідуально, контролюючи результати електрокардіографії та рівні калію і глюкози в крові.

Висновки. Поєднання глюкози, інсуліну та калію дозволяє одночасно вирішити кілька проблем, пов'язаних із метаболічними порушеннями, що може бути особливо корисним у постопераційний період. Інфузійний розчин ГІК є ефективним та зручним методом лікування гіпокаліємії у пацієнтів загальної хірургії.

АСОРТИМЕНТ МІОРЕЛАКСАНТІВ У ТЕРАПІЇ ЗАХВОРЮВАНЬ ОПОРНО-РУХОВОГО АПАРАТУ

Хворост О. П., Опрошанська Т. В., Скребцова К. С.
Національний фармацевтичний університет, м. Харків, Україна
Кафедра фармакогнозії та нутриціології,
Кафедра фармацевтичної технології, стандартизації та сертифікації ліків
khvorost09101960@gmail.com

Вступ. На сьогодні проблеми з опорно-руховим апаратом (ОРА) стрімко поширюються серед населення планети. Згідно АТС-класифікації засоби, що впливають на ОРА підрозділяються на: протизапальні та протиревматичні (М01); засоби, що застосовують місцево у разі суглобового і м'язового болю (М02); міорелаксанти (М03); засоби, що застосовуються для лікування подагри (М04); засоби, що застосовуються для лікування захворювань кісток (М05); інші засоби (М09). Допоміжною симптоматичною ланкою терапії захворювань ОРА, зокрема, ноцицептивного болю є міорелаксанти, що зменшують больові симптоми за рахунок вибіркового розслаблення скелетних м'язів, ефективні при м'язових спазмах, болі в спині або шиї, а також за інших проблем, пов'язаних з м'язами та суглобами. Ці препарати можуть усунути гострий больовий синдром, що виник через гіпертонус і спазм різних м'язів.

Мета. Проаналізувати асортимент засобів групи М03 – міорелаксантів, що застосовуються для лікування патологій ОРА.

Матеріали та методи. Аналіз проводили на основі даних Державного реєстру лікарських засобів України, інтернет-торгівельних платформ станом на 2024 рік.

Результати. За результатами здійснених маркетингових досліджень асортименту засобів групи М03 на фармацевтичному ринку України встановлено, що на ринку репрезентовано 34 лікарських засоби цієї групи. Аналіз асортименту досліджуваної групи за лікарськими формами показав, що вони представлені таблетками, капсулами, ліофілізованим порошком для інфузійних розчинів, розчинами для ін'єкцій, порошком для розчинення. Основна їх частина – 53% представлена ін'єкційними формами, таблетки складають 35 %. Більшість лікарських засобів групи М03 на ринку України становлять вітчизняні препарати, їх питома вага складала 53 %. Домінуючим українським підприємством є АТ «Фармак» – 5,9 %. Закордонні засоби-міорелаксанти виробляють 8 країн. Лідерами за загальною кількістю асортиментних позицій препаратів-імпортерів є Польща, Турція та Велика Британія (по 8,8 %). Домінують міорелаксанти центральної дії, АФІ в яких виступають синтетичний н-холінолітик толперізон, а також циклобензаприн. Присутні на ринку міорелаксанти периферійної дії, що містять напівсинтетичне сульфідне похідне колхікозиду тіоколхікозид.

Висновки. Зважаючи на побічні ефекти застосування міорелаксантів, пошук нових вітчизняних безпечних ефективних препаратів є актуальним.

MECHANISMS ASSOCIATED WITH URIC ACID IN NEUROPROTECTIVE STRATEGIES

Tovchiga O. V., Inkielewicz-Stępniak I.
[1]Medical University of Gdańsk, Gdańsk, the Republic of Poland
Department of Pharmaceutical Pathophysiology
olga.tovchiga@gumed.edu.ua

Introduction. Uric acid is known to exert the beneficial and detrimental effects including those directed on the CNS. Among the beneficial ones, there is neuroprotection and possible cognitive enhancement, and significance of these fields for drugs discovery can hardly be overestimated.

Aim. This study aimed at summarization of data in the literature concerning the possible ways of neuroprotection associated with uric acid which could be further exploited in drug discovery or modification of therapeutic strategies. There is a high massive of data concerning the central effects of uric acid, and their contradictory nature causes the comparison of this substance with "double-edge sword" or "two-faced Janus" [1]. That is why it is important to elucidate the mechanisms of its beneficial effects and determine conditions in which they are realized.

Funding sources or registered topic of the research. The work was supported by funding ST-54 from the Medical University of Gdańsk.

Results. There is an impressive example of verification of uric acid effects in stroke (its supplementation was proposed as promising adjuvant intervention for mechanical thrombectomy), namely the approaches similar to those in clinical research were implemented in preclinical studies. Namely, the criteria of efficacy were predefined, the animals of the different age, sex, and with comorbidities were included, with the results being obtained in several independent laboratories. Notably, under such strict and predetermined conditions efficacy of uric acid was confirmed, and it is supposed that its influence is directed on the microcirculatory hypoperfusion and oxidative stress [2]. Indicatively, the results of current systematic analysis [3] have shown that lower uric acid blood concentrations were independently associated with a decreased risk of hemorrhagic transformation in post-acute ischemic stroke patients. The most recent review article once again emphasizes the U-shaped association between serum uric acid levels and ischemic stroke [4].

Of special interest there are data on the mechanisms involved into the protective role of uric acid. Being far from overestimating the prospects of antioxidants for neuroprotection (out of the context of additional mechanisms of multitarget molecules), it should be noted that uric acid is considered as a natural scavenger of peroxynitrite [5], and the significance of this mechanism is established on the models of neurodegeneration [6] and drug-induced neurotoxicity, for instance caused by cysplatin [7]. Just the ability to inactivate peroxynitrite also contributes to the maintenance of blood brain barrier integrity [5], and it has been proven that uric acid is able to inhibit endothelial cell adhesion and the penetration of inflammatory cells in the CNS [8].

Among the fundamental mechanisms of uric acid protective activity in neurodegeneration there are induction of microglia autophagy and enhanced β-amyloid degradation. These processes are realized through an activation of transcription factor EB (TFEB)-related signaling pathways, and this may help elucidating new targets in neuroprotection [9]. In this context, it also may be important that, in addition to the ability of peroxynitrite scavenging, which was discussed above, uric acid leads to the induction of the Nrf2 signaling pathway linked to the glutathione system in the astrocytes [10].

Since hyperuricemia may lead to the unfavourable consequences [1, 11], the concept has been suggested to use the modulators of the transport systems which to increase just the local concentration of uric acid within the brain. The idea to modulate the level of endogenous metabolite seems to be very attractive as a classical example of pharmacological regulation. Namely, the protein ITM2B, which interacts with GLUT9 and regulates GLUT9-mediated urate transport, was characterized as a molecular link between urate homeostasis and neurodegenerative disorders [13].

And, finally, the gut-brain axis, which attracts great attention in many fields, is also important for the realization of the central effects of uric acid. The significant impact of gut microecology on purine metabolism is generally accepted, and many molecules including those of herbal origin may exert their beneficial effects through modulating the composition of gut microbiota, gut mucosal barrier, inflammatory response, the reactions of purine metabolism, and associated transporters [14]. This helps to explain efficacy of molecules with limited systemic bioavailability and also helps to establish new ways of neuroprotection.

Conclusions. The mechanisms associated with uric acid represent show great promise for future developments within neuroprotective strategies.

References

1. Fang P, Li X, Luo JJ, Wang H, Yang XF. A Double-edged Sword: Uric Acid and Neurological Disorders. Brain Disord Ther. 2013;2(2):109. doi: 10.4172/2168-975X.1000109.
2. Leira EC, Planas AM, Chauhan AK, Chamorro A. Uric acid: a translational journey in cerebroprotection that spanned preclinical and human data. Neurology 2023;101:1068–74. https://doi.org/10.1212/WNL.0000000000207825
3. Qian Y, Li N, Li Y, Tao C, Liu Z, Zhang G, Yang F, Zhang H, Gao Y. Association between uric acid and the risk of hemorrhagic transformation in patients with acute ischemic stroke: a systematic review and meta-analysis. Front Neurol. 2024;15:1378912. doi: 10.3389/fneur.2024.1378912.
4. Zhu B, Huang X, Zhang J, Wang X, Tian S, Zhan T, Liu Y, Zhang H, Chen S, Yu C. A New Perspective on the Prediction and Treatment of Stroke: The Role of Uric Acid. Neurosci Bull. 2024 Sep 23. doi: 10.1007/s12264-024-01301-3.
5. Kean RB, Spitsin SV, Mikheeva T, Scott GS, Hooper DC. The peroxynitrite scavenger uric acid prevents inflammatory cell invasion into the central nervous system in experimental allergic encephalomyelitis through maintenance of blood-central nervous system barrier integrity. J Immunol. 2000;165(11):6511-8. doi: 10.4049/jimmunol.165.11.6511.
6. Mattson MP, Goodman Y, Luo H, Fu W, Furukawa K. Activation of NF-kappaB protects hippocampal neurons against oxidative stress-induced apoptosis: Evidence for induction of manganese

superoxide dismutase and suppression of peroxynitrite production and protein tyrosine nitration. J Neurosci Res 1997;49:681–97. https://doi.org/10.1002/(SICI)1097-4547(19970915)49:6<681 :AID-JNR3>3.0.CO;2-3
7. Sun M, Mao X-F, Li Z-M, Zhu Z-H, Gong D-M, Lu L, et al. Endothelial peroxynitrite causes disturbance of neuronal oscillations by targeting caspase-1 in the arcuate nucleus. Redox Biol 2021;47:102147. doi.org/10.1016/j.redox.2021.102147
8. Feng J, Zheng Y, Guo M, Ares I, Martínez M, Lopez-Torres B, et al. Oxidative stress, the blood–brain barrier and neurodegenerative diseases: The critical beneficial role of dietary antioxidants. Acta Pharm Sin B 2023;13:3988–4024. https://doi.org/10.1016/j.apsb.2023.07.010.
9. Xiao Q, Wang J, Tian Q, Tian N, Tian Q, He X, et al. Uric acid mitigates cognitive deficits via TFEB-mediated microglial autophagy in mice models of Alzheimer's disease. Mol Neurobiol 2023. https://doi.org/10.1007/s12035-023-03818-6.
10. Bakshi R, Zhang H, Logan R, Joshi I, Xu Y, Chen X, et al. Neuroprotective effects of urate are mediated by augmenting astrocytic glutathione synthesis and release. Neurobiol Dis 2015;82:574–9. https://doi.org/10.1016/j.nbd.2015.08.022
11. Tovchiga O V, Shtrygol' SYu. Uric acid and central nervous system functioning (a literature review). Biology Bulletin Reviews 2014;4:210–21. https://doi.org/10.1134/S2079086414030086
12. Mandal AK, Mount DB. Interaction Between ITM2B and GLUT9 links urate transport to neurodegenerative disorders. Front Physiol 2019;10:1323. https://doi.org/10.3389/fphys.2019.01323
13. Wang H, Zheng Y, Yang M, Wang L, Xu Y, You S, Mao N, Fan J, Ren S. Gut microecology: effective targets for natural products to modulate uric acid metabolism. Front Pharmacol. 2024 Aug 28;15:1446776. doi: 10.3389/fphar.2024.1446776.

VAGINAL DRUG DELIVERY SYSTEMS: INNOVATIONS THAT CHANGE THE PHARMACOLOGICAL PROPERTIES OF DRUGS

A. Makovetskyi[1], G. Zaychenko[2]
Bogomolets National Medical University, Kyiv, Ukraine
Department of Pharmacology
Andrew@Makovetskyi.com[1], Ganna.Zaychenko@ukr.net[2]

Drug delivery systems (DDSs) have fundamentally transformed the administration of medications, significantly enhancing both their efficacy and safety. These systems include advanced technologies to control drug release, target specific sites within the body, and reduce adverse effects. Among the most noteworthy innovations in this field is the development of vaginal drug delivery systems (VDDS). VDDS have redefined the pharmaceutical landscape by offering innovative solutions that enhance the local and systemic effectiveness of therapeutic agents. By leveraging the unique anatomical and physiological characteristics of the vaginal environment, these systems enable improved bioavailability, prolonged drug retention, and targeted delivery, making them a pivotal advancement in modern therapeutic strategies.

In the realm of obstetrics and gynecology (O&G), VDDS are playing an increasingly important role, offering significant advantages for both local and systemic

treatments. The anatomical and physiological characteristics of the vaginal route make it ideal for delivering drugs that address conditions specific to women's health, such as bacterial infections, hormonal regulation, and contraception. The ability to provide localized therapeutic effects while minimizing systemic side effects makes VDDS a safer and more effective option for treating diseases of the reproductive system in women. Moreover, the rich vascular network of the vaginal mucosa, coupled with the bypassing of hepatic metabolism, enhances the bioavailability of various active pharmaceutical ingredients (APIs), including hormones, peptides, and small molecules. These advantages, coupled with the incorporation of advanced bioadhesive technologies, have led to superior drug retention and distribution, further optimizing therapeutic outcomes for both local and systemic applications. As a result, VDDS continues to generate growing interest as a versatile and effective drug delivery platform.

The knowns forms of vaginal dosage include vaginal rings, gels, creams, pessaries and suppositories. Each form of dosage is adapted for specific therapeutic applications, allowing controlled release of medications while avoiding systemic side effects commonly associated with oral or injection therapy. Such dosage forms as pessaries (vaginal suppositories) have long been used in gynecological practice. These forms can treat both local infections and systemic conditions by leveraging the rich vascularization of the vaginal wall, allowing for effective drug absorption (Jahromi M. M. et al, 2015). While the vaginal rings are particularly remarkable for hormonotherapy, providing continuous release of drugs such as progesterone over long periods of time, providing stable plasma levels with minimal fluctuations, making them suitable for hormone replacement therapy (Cheung J. et al., 2010). Vaginal gels and creams have also been widely studied for conditions such as hyperprolactinemia and HIV prevention. These forms can provide localized therapy with minimal systemic absorption and reduce side effects while maintaining therapeutic efficacy (Vermesh M. et al., 1988).

Recent innovations in vaginal drug delivery systems have considerably changed the treatment of both gynecological and systemic conditions, providing more effective and patient-centered therapeutic solutions. These advances give priority to improving drug bioavailability, extending retention time and simplifying administration, while minimizing adverse effects and improving patient compliance.

Key developments include the use of bioadhesive gels and mucoadhesive polymers, which extend the residence time of drugs within the vaginal mucosa. For example, bioadhesive polymers such as polyacrylate and chitosan have been used to improve the preservation of drugs and reduce leakage, a common limitation of conventional formulations (Acartürk et al.). These materials enable a sustained and localized release of APIs, improving treatment outcomes and enhancing patient compliance, particularly for conditions requiring prolonged drug exposure.

Another significant innovation is the introduction of controlled-release systems like intravaginal rings and hydrogels, which ensure a continuous release of medication over an extended period. These delivery platforms are tailored to ensure pro-

longed therapeutic effects while minimizing the side effects commonly associated with oral or injectable routes. For example, vaginal rings have proven effective in hormone replacement therapy by maintaining steady plasma levels of drugs like progesterone, reducing fluctuations and systemic side effects and minimizing dosing frequency (Cheung J. et al., 2010). Vaginal gels and creams have also been widely studied for conditions like hyperprolactinemia and HIV prevention. These forms can offer localized drug action with high concentrations in vaginal tissues, well above inhibitory levels with minimal systemic absorption (Nishiura K. et al., 2021), reducing side effects while maintaining therapeutic efficacy (Vermesh M. et al., 1988).

Nanotechnology is an essential driver of the progress of DDS, particularly through the development of nanoparticles and liposomes. These nanocarriers allow targeted drug delivery to weakened tissues, minimize exposure to healthy tissues and reduce systemic toxicity – an important breakthrough in anti-tumor therapy (Ahmad W. & Jawed R., 2022). In VDDS, nanoparticle systems have emerged as a transformational solution, particularly for macromolecules such as peptides and proteins, which often face absorption challenges. Nanoparticles improve drug permeability in the vaginal epithelium, allowing the release of more controlled and sustained drugs (Priyanka P. et al., 2022). These technologies are especially valuable in the treatment of a wide range of diseases, ranging from bacterial infections to systemic diseases such as osteoporosis and migraine, by ensuring more effective drug delivery and therapeutic results.

Additionally, thermosensitive gels and self-emulsifying drug delivery systems (SEDDS) have been developed, which change their consistency in response to body temperature or fluid interaction, ensuring efficient drug release (Köllner S. et al., 2017). These technologies are particularly useful for conditions like bacterial vaginosis, HIV prevention, and other infections where controlled drug release is critical. Also, novel approaches like micromachining, biosensors, and responsive drug delivery systems have emerged, which enable real-time monitoring and controlled release of drugs, paving the way for more personalized treatment protocols (Deo S. K. et al., 2003). This trend towards integrating technology and pharmaceuticals continues to push the boundaries of how drugs are administered, with the potential to replace conventional drug forms soon.

Innovations such as bio-compatible polymers and smart materials are further enhancing drug delivery by improving bioavailability and offering more efficient drug transport to specific cells and tissues (Mainardes R. M. & Silva L. P., 2004). Moreover, non-invasive delivery methods, which include mucosal applications, is increasingly utilized to provide painless administration across biological barriers like the skin or mucosa (Vllasaliu D., 2021).

These innovations not only optimize local treatment, but also offer promising methods for delivering systemic drugs, significantly expanding the potential of VDDS in clinical practice. Controlled-release mechanisms, in particular, allow medications to be administered at a controlled rate over an extended period, reducing the frequency of doses and enhancing convenience for patients (Nayak U. Y et al., 2016).

The ongoing exploration of bioadhesive and controlled-release technologies marks a transformative phase in VDDS development, allowing for more effective treatments that improve both patient compliance and therapeutic outcomes (Langer R., 1991). Through the integration of these innovative technologies, drug absorption and therapeutic efficacy are optimized, patient comfort is improved, and many challenges associated with the old formulation are reduced. This progress will transform vaginal drug delivery into a versatile and reliable option for local and systemic treatments, which plays a crucial role in women's health.

In conclusion, innovations in vaginal drug delivery systems, ranging from bio-adhesive polymers to nanoparticle-based platforms, hold significant potential for treating diseases of the female reproductive system.

References:

1. Acartürk, F. (2009). Mucoadhesive vaginal drug delivery systems. *Recent Patents on Drug Delivery & Formulation, 3*(3), 193–205. https://doi.org/10.2174/187221109789105658
2. Ahmad, W., & Jawed, R. (2022). An updated review on novel drug delivery system. *Research Journal of Pharmaceutical Dosage Forms and Technology, 14*(3), 211-216. https://doi.org/10.52711/0975-4377.2022.00034
3. Cheung, J., DiLiberti, C., & Reape, K. Z. (2010). Evaluation of the single dose and steady state pharmacokinetics of a progesterone vaginal ring. *Fertility and Sterility, 94*(4 Suppl), S176. https://doi.org/10.1016/j.fertnstert.2010.07.699
4. Deo, S. K., Moschou, E. A., Peteu, S. F., Bachas, L. G., & Daunert, S. (2003). Responsive drug delivery systems. *Analytical Chemistry, 75*(9), 207A–213A. . PMID: 12751526.
5. Köllner, S., Nardin, I., Markt, R., Griesser, J., Prüfert, F., & Bernkop-Schnürch, A. (2017). Self-emulsifying drug delivery systems: Design of a novel vaginal delivery system for curcumin. *European Journal of Pharmaceutics and Biopharmaceutics, 115*, 268-275. https://doi.org/10.1016/j.ejpb.2017.03.012
6. Langer, R. (1991). Drug delivery systems. *MRS Bulletin, 16*, 47–49. https://doi.org/10.1557/S0883769400056050
7. Mainardes, R. M., & Silva, L. P. (2004). Drug delivery systems: Past, present, and future. *Current Drug Targets, 5*(5), 449–455. https://doi.org/10.2174/1389450043345407
8. Jahromi, M. M., Ghaemi, H., Ajdary Tafti, M., Arabzadeh, A., & Afsharypuor, S. (2015). Vaginal and rectal dosage forms in Iranian traditional pharmacy. *Jundishapur Journal of Natural Pharmaceutical Products, 10*(2), e19935. https://doi.org/10.17795/jjnpp-19935
9. Nayak, U. Y., Mutalik, S., & Udupa, N. (2016). An overview of novel and controlled drug delivery systems. *Indian Drugs, 53*(5), 5-12. https://doi.org/10.53879/id.53.05.10396
10. Nishiura, K., Sharma, S., Sterling, M., Makarova, N., Martin, A., Dinh, C., Mitchell, J., García-Lerma, J. G., Heneine, W., & Dobard, C. (2021). Pharmacokinetics of vaginally applied integrase inhibitors in macaques. *Journal of Antimicrobial Chemotherapy, 76*(11), 2894–2901. https://doi.org/10.1093/jac/dkab293
11. Priyanka, P., Sri Rekha, M., & Devi, A. S. (2022). Review on formulation and evaluation of solid lipid nanoparticles for vaginal application. *International Journal of Pharmacy and Pharmaceutical Sciences, 14*(1), 1–8. https://doi.org/10.22159/ijpps.2022v14i1.42595
12. Vermesh, M., Fossum, G. T., & Kletzky, O. A. (1988). Vaginal bromocriptine: Pharmacology and effect on serum prolactin in normal women. *Obstetrics and Gynecology, 72*(5), 693–698. PMID: 3173919.
13. Vllasaliu, D. (2021). Non-invasive drug delivery systems. *Pharmaceutics, 13*(5), 611. https://doi.org/10.3390/pharmaceutics13050611

РОЛЬ ФАРМАЦЕВТА В ЗАБЕЗПЕЧЕННІ РАЦІОНАЛЬНОГО ЛІКУВАННЯ ПАЦІЄНТІВ З АРТЕРІАЛЬНОЮ ГІПЕРТЕНЗІЄЮ

Туранська Н. П., Вєтрова К. В., Отрішко І. А.
Національний фармацевтичний університет, м. Харків, Україна
Кафедра фармакології та клінічної фармації
vkv_katya@ukr.net

Вступ. На сьогоднішній день серцево-судинні захворювання залишаються найрозповсюденішою групою захворювань, що негативно впливають на якість та тривалість життя людей в усьому світі. Артеріальна гіпертензія (АГ) є одним з найпоширеніших захворювань серцево-судинної системи. За даними ВООЗ у світі 1,28 млрд. людей у віці від 30 до 79 років хворіють на АГ. До основних груп лікарських препаратів, що застосовуються для лікування АГ відносять: інгібітори ангіотензинперетворюючого ферменту, блокатори рецепторів ангіотензину II, β-адреноблокатори, блокатори кальцієвих каналів та діуретики (тіазидні, тіазидоподібні). Зазвичай, лікування АГ потребує застосування комбінацій антигіпертензивних препаратів.

Мета. Визначення ролі фармацевта в забезпеченні раціонального лікування пацієнтів з АГ.

Матеріали та методи. Аналіз та узагальнення даних наукової літератури.

Результати. При зверненні до аптеки відвідувачів з АГ з метою придбання антигіпертензивних засобів фармацевт має запитати рецепт, оскільки такі препарати застосовуються лише за призначенням лікаря. Фармацевт повинен проінформувати пацієнта, що за державною програмою «Доступні ліки» за наявності рецепта лікаря можна отримати препарати для лікування АГ безкоштовно або зі знижкою. Також фармацевт має наголосити, що лікування АГ проводиться постійно, впродовж всього життя, а не курсами до поліпшення стану. В рамках проведення фармацевтичної опіки обов'язковою є консультація відвідувача щодо умов раціонального застосування антигіпертензивних препаратів, можливих проявів їх побічної дії та взаємодії з іншими групами лікарських засобів. Фармацевт повинен проінформувати хворого на АГ, що при виникненні побічних ефектів необхідно припинити використання препарату та звернутися до лікаря. Фармацевт має навчити пацієнта правильно вимірювати артеріальний тиск (АТ) і наголосити на постійному контролі показників АТ (ведення щоденника АТ, використання мобільних додатків тощо). У межах фармацевтичної опіки фармацевт має надати рекомендації щодо немедикаментозного лікування АГ: зменшення ваги тіла, регулярні фізичні вправи, обмеження вживання солі, відмова від паління, зменшення вживання алкоголю тощо.

Висновки. Проведення фармацевтом належної фармацевтичної опіки при відпуску з аптеки антигіпертензивних препаратів сприятиме підвищенню ефективності та безпечності терапії АГ, а також прихильності хворих до лікування.

ФУНКЦІОНАЛЬНЕ НАПОВНЕННЯ РОЛЕЙ ФАРМАЦЕВТІВ ПРИ ПРОФІЛАКТИЦІ ІШЕМІЧНОЇ ХВОРОБИ СЕРЦЯ НА ПЕРВИННОМУ РІВНІ МЕДИЧНОЇ ДОПОМОГИ

Білоусова Н. А.

Національний університет охорони здоров'я України імені П.Л. Шупика, м. Київ, Україна

Кафедра кардіології

arinatala@gmail.com

Вступ. За даними 2021 року Всесвітньої організації охорони здоров'я (ВООЗ) Україна посідала перше місце серед європейських країн за смертністю від серцево-судинних захворювань (ССЗ), що становить 49,82% [1]. Основною причиною смертності від ССЗ в Україні є ішемічна хвороба серця (ІХС) [2]. Доведено, що під впливом соціально-економічних та стресових факторів захворюваність та смертність від ІХС зростає [3]. Отже, гіпотетично захворюваність і смертність від ІХС під час війни в Україні зростає, що потребує відповідної уваги [4].

На Глобальному рівні відбувається переорієнтація фармацевтичних послуг із фармацевтичного забезпечення (ФЗ) на надання ФД з розширенням Ролей і функцій GPP, які забезпечують економічну стійкість не тільки аптечних закладів, а й національних систем охорони здоров'я (ОЗ) [5].

На державному рівні в нову редакцію Закону України «Про лікарські засоби» (ЛЗ) [6], який вступає в дію після закінчення воєнного стану, введено поняття «фармацевтичної допомоги» [7]. На галузевому рівні затверджено протоколи надання фармацевтичної допомоги пацієнтам із ССЗ, які передбачають ФЗ населення ЛЗ та не враховують усього комплексу заходів при наданні ФД, передбачених законодавством.

Таким чином, актуальними є вивчення розробка та впровадження комплексу заходів із надання фармацевтичної допомоги пацієнтам із ІХС на первинному рівні медичної допомоги (МД).

Мета. Метою дослідження є обґрунтування функціонального наповнення Ролей GPP при наданні фармацевтичної допомоги пацієнтам з ІХС на первинному рівні МД.

Матеріали та методи. Матеріалами дослідження стали міжнародні стратегії, нормативно-правові документи вітчизняного та зарубіжного законодавства, клінічні рекомендації ESC/AHA, американські протоколи надання ФД. Використано методи компаративного аналізу, синтезу, моделювання, дедукції, індукції, узагальнення.

Результати. За результатами дослідження встановлено, що на Глобальному рівні відбувається розширення та функціональне наповнення Ролей GPP в процесі надання ФД за рахунок інтеграції фармацевтів в первинну медико-санітарну допомогу в якості членів мультидисциплінарних команд із подальшою міждисциплінарною взаємодією. Визначено, що дані заходи ма-

ють економічний вплив на стійкість національних систем ОЗ. Виокремлено освітню та інформаційну функції фармацевтів у наповненні Ролей GPP. Акцентовано увагу на командній роботі у відповідності із кращими світовими практиками. Запропоновано модель надання ФД пацієнтам із ІХС на первинному рівні МД. Запропоновано розширення протоколів фармацевтів з урахуванням функціонального наповнення Ролей GPP.

Висновки. Розроблено та обґрунтовано модель надання ФД пацієнтам із ІХС на первинному рівні МД на засадах оцінки медичних технологій. Обґрунтовано функціональні складові Ролей GPP з урахуванням зменшення кількості вторинних серцево-судинних подій, інвалідізації населення та зниження навантеження на національну систему ОЗ.

Перелік використаних джерел інформації:

1. Health profile Ukraine. World Life Expectancy. URL: https://www.worldlifeexpectancy.com/country-health-profile/ukraine.
2. Державна служба статистики України. Інститут демографії та соціальних досліджень імені М.В. Птухи Національної Академії Наук України. Населення України. Інформаційно-аналітичні матеріали. Державна служба статистики України. URL: http://db.ukrcensus.gov.ua/PXWEB2007/ukr/publ_new1/2022/publ2022.asp.
3. Kornelia Kotseva, Guy De Backer, Dirk De Bacquer, Lars Rydén, Arno Hoes, Diederick Grobbee, Aldo Maggioni et al. The EUROASPIRE V Investigators, Primary prevention efforts are poorly developed in people at high cardiovascular risk: A report from the European Society of Cardiology EURObservational Research Programme EUROASPIRE V survey in 16 European countries, European Journal of Preventive Cardiology, Volume 28, Issue 4, April 2021, Pages 370–379, https://doi.org/10.1177/2047487320908698
4. Bilousova NA. Primary prevention of a coronary heart disease within the structure of pharmaceutical care provision: functional role fulfillment of the pharmacist. Фарм. час. [Internet]. 2024 Sep. 30 [cited 2024 Oct. 14];(3):42-5. https://doi.org/10.11603/2312-0967.2024.3.14862.
5. International Pharmaceutical Federation (FIP). Economic sustainability and pharmacy: A commentary article. The Hague: International Pharmaceutical Federation; 2023. P. 21. URL: https://www.fip.org/file/5870
6. Про лікарські засоби: Закон України від 28.07.2022 р. № 2469-IX : станом на 17 верес. 2023 р. URL: https://zakon.rada.gov.ua/laws/show/2469-20#Text.
7. Про затвердження протоколів фармацевта : Наказ МОЗ України від 05.01.2022 р. № 7. URL: https://zakon.rada.gov.ua/rada/show/v0007282-22#Text

ОПТИМІЗАЦІЯ СИМПТОМАТИЧНОГО ЛІКУВАННЯ ТРИВОГИ У ПАЦІЄНТІВ З АРТЕРІАЛЬНОЮ ГІПЕРТЕНЗІЄЮ В УМОВАХ АПТЕКИ

Сахарова Т. С.

Одеський національний університет імені І.І. Мечникова, м. Одеса, Україна
Кафедра фармакології та технології ліків
pharmacology.technology20@gmail.com

Вступ. Дані численних епідеміологічних досліджень засвідчують взаємообумовлений зв´язок надмірної тривоги/тривожних розладів (ТР) та артеріальної гіпертензії (АГ). З одного боку, психологічні наслідки стресу, у тому числі тривога, є відомими предикторами АГ, з іншого – ТР погіршують якість життя хворих на АГ та підвищують ризик серцево-судинних ускладнень. Саме тому за наявності певного ТР пацієнти кардіологічного профілю повинні отримувати адекватне протитривожне лікування, рекомендоване відповідними клінічними настановами. Стандартом надання лікарської допомоги, у разі звернення пацієнта зі скаргою на тривогу до аптеки, є «Протокол фармацевта при відпуску лікарських засобів без рецепта «Симптоматичне лікування стресу»», затверджений Наказом МОЗ України № 7 від 05.01.2022 р.

Мета. визначити підходи до оптимального вибору безрецептурних протитривожних лікарських засобів (ПТЛЗ) пацієнтам з АГ на підставі рекомендацій чинного Протоколу фармацевта «Симптоматичне лікування стресу» та принципів раціональної фармакотерапії.

Матеріали та методи. Інформаційний пошук (аналіз літературних джерел та Internet-ресурсів) та індуктивний метод наукового дослідження.

Результати. Проведена оцінка ризику можливої фармакодинамічної та фармакокінетичної взаємодії безрецептурних ПТЛЗ, рекомендованих вищезгаданим Протоколом, з найуживанішими у медичній практиці антигіпертензивними ЛЗ (АГЛЗ). Усім «протокольним» ПТЛЗ (гліцин, мебікар, препарати звіробою, валеріани, пустирника, пасифлори)) притаманний слабкий антигіпертензивний ефект, який не розцінюється як клінічно значущий та нівелюється з часом. Застереження стосуються лише гліцину, доза якого повинна коригуватися на тлі регулярного контролю АТ. Наукові дані щодо фармакокінетичних характеристик розглянутих ПТЛЗ свідчать про відсутність взаємодії з основними АГЛЗ на шляхах проходження організмом. Втім, уваги заслуговують препарати звіробою, БАР яких є потужними індукторами ізоформи цитохрому P_{450} – CYP3A. Ця властивість обмежує (виключає) їх одночасне застосування з ЛЗ різних груп, які є субстратами цього ж ферменту, зокрема, блокаторами кальцієвих каналів (ніфедипін, дилтіазем, верапаміл та інш.).

Висновки. Отримані результати визначають доцільність оптимізації симптоматичного лікування тривоги у хворих на АГ в умовах аптеки для попередження небажаних наслідків поліфармації та досягнення адекватного контролю за АТ.

ДЕЯКІ АСПЕКТИ ЗАСТОСУВАННЯ ШАМПУНІВ ІЗ КЕТОКОНАЗОЛОМ ПРИ ЛУПІ

Ващенко О. О.

Львівський національний медичний університет імені Данила Галицького, м. Львів, Україна

Кафедра технології ліків і біофармації

o_vashchenko@ukr.net

Вступ. Лупа є стійким рецидивуючим станом, що часто діагностують як легку форму себорейного дерматиту. Серед лікарських засобів, які призначають для лікування та профілактики лупи є шампуні із кетоконазолом.

Мета. Проаналізувати основні аспекти застосування шампунів із кетоконазолом для лікування та профілактики лупи.

Матеріали та методи. Інформаційний аналіз наукових фармацевтичних та медичних джерел.

Результати. Етіологія лупи на сьогодні залишається дискусійним питанням, проте серед вчених домінує думка щодо визначальної ролі тріади факторів: генетична схильність, надмірна колонізація грибів *Malassezia spp*., зміна складу шкірного сала. З огляду на етіопатогенез лупи, для лікування цього стану зокрема застосовують протигрибкові шампуні. При виборі шампуню доцільно враховувати наступні рекомендації (Британський національний формуляр): при легких формах лупи застосовувати шампуні із цинку піритіоном, селеном, дьогтем, саліциловою кислотою; при більш складних і стійких формах – шампуні із кетоконазолом (ШК).

Кетоконазол – синтетичне похідне імідазолдіоксолану, що виявляє протигрибкову активність до дерматофітів і дріжджів; особливо виражений ефект щодо *Malassezia spp*. На фармацевтичному ринку України ШК зареєстровані як лікарські засоби. Ці шампуні можна застосовувати із 12-річного віку: для лікування лупи – 2 рази на тиждень протягом 2-4 тижнів; для профілактики – 1 раз на тиждень/2 тижні. Окрім частоти застосування, на ефективність препарату впливає час експозиції, тому рекомендовано шампунь після нанесення витримати на голові 3-5 хв, а потім змивати. Серед побічних ефектів ШК можливі місцеві алергічні реакції; волосся може стати жирним або сухим; в окремих випадках, в основному у пацієнтів з хімічно ушкодженим або сивим волоссям, спостерігається незначна зміна відтінку кольору волосся. Кокранівська база містить публікації, що вказують на ефективність ШК при себорейному дерматиті, яка є порівнюваною із ефективністю стероїдів при короткострокових дослідженнях, при цьому частота побічних реакцій при застосуванні кетоконазолу є нижчою; у порівнянні із 1% цинку піритіоновим шампунем, ШК демонструє вищу ефективність та нижчий рівень рецидивів лупи.

Висновки. Враховуючи дані щодо ефективності та безпечності, шампуні із кетоконазолом доцільно застосовувати для лікування складних форм лупи.

ТРАВМИ, ЩО НАНЕСЕНІ ДОМАШНІМИ ТВАРИНАМИ, ТА ЇХ ЛІКУВАННЯ

Катурова В. В.
КНП ХОР «Обласна клінічна лікарня», м. Харків, Україна
Стоматологічне відділення № 3 (дитяче)

Вступ. Щодня у домашньому побуті виникають травми від домашніх тварин. Найчастіше людей кусають домашні собаки, рідше – кішки. Кількість укушених ран щелепно-лицьової ділянки за останні роки значно зросла. На їх частку припадає 10 % від загальної кількості хворих з ізольованими ранами.

Мета. Метою даної роботи став аналіз особливостей перебігу травм від тварин та їх лікування.

Матеріали та методи. Досліджували дані наукової літератури з практичної медицини. На основі даних оцінено особливості перебігу ран, отриманих від тварин та наведені методи їх лікування. Дослідження виконано з використанням бібліосемантичного аналізу (інформаційно-аналітичного узагальнення даних із наукових літературних джерел).

Результати. Укус тварини – це рана з надзвичайно високою, майже стовідсотковою ймовірністю інфікування. За даними статистики приблизно, 5–20 % укусів собак і 30–60 % укусів котів стають інфікованими. Серед інфекційних агентів, які можуть потрапити в організм при укусі тварини, є збудники смертельно небезпечних хвороб – сказу та правця. Також, укуси частіше множинні, іноді супроводжуються пошкодженням нервових стовбурів, великих судин, дефектами м'яких тканин, поєднанням різних видів ран і різними термінами ранового процесу (на 3-5-й день набряк тканин може збільшуватися через виникнення запального процесу).

Відразу після укусу лікарі рекомендують рану промити 10 % розчином господарського мила або йодовідоном. Зазвичай це рани з нерівними, рваними і розчавленими краями, тому їх нежиттєздатні шари необхідно вирізати. Укушені рани слід дренувати протягом 5-7 днів, тому що всі вони на 100 % інфіковані. Хірург разом з рабіологом визначає показання до вакцинації проти сказу відповідно до наказу МОЗ України «Про профілактику захворювань на сказ людини», оскільки вірус сказу поширюється периневрально зі швидкістю 3-5 мм на годину і при неправильно обраній тактиці лікування дитина може померти. При укусах легкого і середнього ступеня тяжкості, нанесених не щепленими тваринами, призначають: 2-разове введення вакцини типу Фермі (з мозку овець) або культуральної інактивованої вакцини проти сказу (CAV) в один день: дітям до 10 років – 1,5 мл з інтервалом 30 хвилин, дітям 10 років і старше – 3 мл. Також первинною профілактикою інфікування є застосування антибактеріальних препаратів амоксицилін/ клавуланова кислота (500/125 мг 3 рази на добу для дорослих) або комбінація тетрацикліну (доксициклін 150 мг один раз на добу або 100 мг двічі на добу) та метронідазолу (400–500 мг 3 рази на добу).

Висновки. Отже, аналіз показав, що в побуті люди отримують травми від тварин, серед яких укушені рани найпоширеніші. Від правильної та своєчасної наданої медичної допомоги може залежати не тільки здоров’я, але й життя постраждалого. Тому, фахівці рекомендують відразу звертатися до лікарень.

ВИВЧЕННЯ ДУМКИ ПАЦІЄНТІВ І ФАРМАЦЕВТІВ ЩОДО ЗАСТОСУВАННЯ РЕЦЕПТУРНИХ ЛІКАРСЬКИХ ЗАСОБІВ

Ривак Т. Б., Сувала Н. С.

Львівський національний медичний університет імені Данила Галицького, м. Львів, Україна

Кафедра менеджменту в охороні здоров'я, фармакотерапії і клінічної фармації

tanusha1905@gmail.com

Вступ. Професійні обов'язки фармацевта в умовах аптеки передбачають реалізацію (відпуск) споживачам лікарських засобів (ЛЗ) як безрецептурних, так і рецептурних. Згідно з визначенням ВООЗ, головним інструментом фармацевта при відпуску ЛЗ є фармацевтична опіка. Як складові фармацевтичної опіки, консультування пацієнтів і вирішення лікопов'язаних проблем *(drug-related problems, DRPs)*, є ключовими напрямками професійної діяльності фармацевта. Результати міжнародних досліджень доводять, що зосередження фармацевтів на проблемах і потребах пацієнтів під час відпуску рецептурних ЛЗ та їх фахові консультації, допомагають підвищити ефективність і безпеку лікування.

Мета. Вивчити думку пацієнтів і фармацевтів щодо застосування рецептурних ЛЗ на основі анкетного опитування.

Матеріали та методи. Дослідженням охоплено 79 респондентів, із них 50 пацієнтів і 29 фармацевтів, які заповнили відповідні анкети в *Google Forms*. Методи: системного підходу, клініко-фармацевтичний, анкетного опитування.

Результати. Встановлено, що 66,0% опитаних приймають рецептурні ЛЗ, 45,0% отримують їх за допомогою електронного рецепта, 37,5% – паперового рецепта довільного формату та 32,5% – паперового рецепта, виписаного на бланку Ф1. Разом із тим, 60,0% респондентів зазначили, що купують рецептурні ЛЗ без рецепта. Найчастіше пацієнти застосовують наступні групи рецептурних ЛЗ: нестероїдні протизапальні засоби – 47,5%, антибактерійні – 37,5%, антигіпертензивні – 35,0%, антитромботичні – 20,0%, антидепресанти – 17,5%. Переважна більшість (85,0%) пацієнтів оплачують повну вартість за рецептурні ліки, проте 25% – отримують їх безоплатно та 17,5% – із доплатою. Лише 15,0% опитаних отримують ЛЗ за державною програмою «Доступні ліки», а 30,0% – не знають про неї. Найчастіші запитання, які виникали у пацієнтів щодо призначених рецептурних ЛЗ, стосувалися: способу застосування – 67,5%; побічних реакцій – 57,5%; тривалості лікування – 47,5%; взаємодії з іншими ЛЗ – 40,0%; протипоказань – 32,5%; дій у випадку пропуску дози – 32,5%; поєднання ліків з їжею та напоями – 30,0%. Відповіді на ці запитання респонденти отримують самостійно з інструкції/інтернету – 80,0% та від фармацевта – 62,5%. У той же час, пацієнти хотіли б отримувати додаткову інформацію щодо раціонального застосування рецептурних ЛЗ у вигляді консультації в аптеці, яка, на їх думку, мала б тривати 3-5 хвилин. Труднощі, з якими зіткнулися респонденти при отриманні ліків за рецептом: 1) нерозбірливий почерк лікаря (важко розібрати спосіб застосування, назву та/або дозу ЛЗ) – 35,0%, 2) технічні пробле-

ми в аптеці (відсутність інтернету, збій в програмі) – 15,0%; 3) не приходило смс-повідомлення з кодом підтвердження для електронного рецепта – 10,0%; 4) помилки в дозуванні та 5) неправильна кількість таблеток в електронному рецепті – по 5,0%. Більше ⅓ (35,0%) опитаних були змушені повторно звертатися до лікаря через помилки в рецепті, з'ясовані в аптеці.

Частка фармацевтів, яким відомий термін лікопов'язані проблеми (DRPs) складає 62,1%, із них 37,9% спостерігали їх у своїй практиці. Зокрема, у випадку потенційних взаємодій між ЛЗ, виписаними в рецепті, 55,2% фармацевтів порадять повторно звернутись до лікаря для попередження негативних наслідків, 41,4% – попередять про це пацієнта. Найчастіше у фармацевтів виникають труднощі з наданням фармацевтичної опіки при відпуску: антидепресантів – 44,8%; антипсихотичних – 41,4%; наркотичних анальгетиків – 34,5%; протиепілептичних – 34,5%; гормональних – 24,1%; анксіолітиків і антигіпертензивних ЛЗ – по 17,2%. У своїй професійній діяльності, для отримання інформації про ЛЗ, фармацевти зазвичай використовують: 1) додаток «Tabletki.ua» – 93,1%; 2) державний реєстр ЛЗ і 3) компендіум – по 44,8%. Стосовно зручності та швидкості відпуску ЛЗ, то для 48,3% фармацевтів не має суттєвого значення форма виписаного рецепта, проте 20,7% надає перевагу електронним рецептам і така ж частка – паперовим рецептам довільного формату. На практиці фармацевтам доводилось комунікувати з лікарем для уточнення певних питань при відпуску ЛЗ за рецептом: 1) у разі нерозбірливого почерку – 75,9%; 2) на прохання пацієнта для уточнення інформації щодо ЛЗ та/або схеми лікування (зокрема, через погану пам'ять) – 24,1%; 3) через некоректно зазначену дозу ЛЗ – 17,2%.

Висновки. Таким чином, більшість (66,0%) опитаних пацієнтів приймають рецептурні ЛЗ та 62,5% потребують додаткової інформації від фармацевта стосовно їх раціонального застосування у вигляді 3-5-хвилинної консультації в аптеці.

За результатами дослідження встановлено необхідність: поглиблення знань фармацевта з надання фармацевтичної опіки при відпуску окремих груп рецептурних ліків (антидепресанти, антипсихотичні, протиепілептичні тощо); навичок роботи з сучасними доказовими ресурсами, включно з інструментами для полегшення прийняття клінічних рішень, і міжпрофесійної фахової комунікації, що дозволить швидко та ефективно вирішувати ідентифіковані DRPs.

Разом із тим, проблему нерозбірливого почерку лікарів у рецептах, яку найчастіше зазначають як пацієнти, так і фармацевти, на нашу думку, можна вирішити перевіривши рецепт за допомогою штучного інтелекту з системою оптичного розпізнавання символів і рукописного тексту, зокрема он-лайн ресурси *2OCR, Handwriting OCR* та інші.

TREATMENT OF HYPERESTROGENISM IN MEN OF FERTILE AGE WITH TESTICULAR TESTOSTERONE DEFICIT

Bayraktar V. M.
Mechnikov National University, Odesa, Ukraine
virolviro6@gmail.com

Introduction. Problem of hyperestrogenism in men of reproductive age frequently could be indicated by different reasons. Food, alcohol, smoking, inflammatory diseases, condition of liver, pancreas, kidneys, stress, depression. To specify hyperestrogenism first of all necessary to test hormones in the blood. Since of hyperestrogenism diagnosed necessary to get qualified treatment. Because without treatment increases weight of body, growing mammary glands, due to fat growing abdomen, decreases libido. All of this shows for hyperestrogenic condition and functional possibilities of liver. High estrogene levels in men suppress secretion of follicle stimulating hormone synthesis, which leads to impaired sperm formation. Men with high levels of estradiol may have difficulty getting or maintaining erection. Excess of estrogene in men leads to increase weight of body and increase deposition of fat. High level of estrogens in man provokes hair loss and appearance of rashes on the face. Pathological manifestations testicular insufficiency accompanied by increasing estrogene secretion it makes much more difficult hormone replacement therapy.

Aim. To confirm estrogene decrease production in testicles and conversion in liver after treatment with antiestrogene klostilbegit and alpha gonadotropine chorionic.

Material and methods. Suffering with hyperestrogenism 20 men in reproductive age 18-35 years old were selected for research. Preliminary hormonal testing and treatment with antiestrogene remedy Klostilbegit (Clomifene citrate) 50 mg in tablets. Klostilbegit made in pharmaceutical PLC, Egis, Budapest, Hungary. At the same time we used alpha gonadotropine chorionicum 1500 IU for injections Ovigil HP, producing in international company Shrree Venkatesh, India. For protection and increase functional possibility of liver we used in the course of treatment glutargine 750.0 mgs. in tablets, produced in pharmaceutical company Zdorovye LLC, Kharkiv, Ukraine.

For testing of blood hormones we used test kits producing in company, Xema, Kiyv, Ukraine. Test kits for immune enzyme analisys: testosterone, estradiol (E_2), progesterone, luteinizing hormone, follicle stimulating hormone, prolactin.

For detection of liver function we used test kits of enzymes such as: lactate dehydrogenase, alanine aminotransferase, aspartat aminotransferase, glutamate aminotransferase, alkaline phosphatase. Test kits produced by company Costa Brava, Spain.

Analyzer for enzyme immunofluorescent analisys used Tosoh AIA-900 Easy reader+. Tosoh Corporation, Tokio, Japan.

Blood tests of hormones and enzymes in tested men made in the clinical laboratory at university clinic.

Results and discussion. Klostilbegit or clomiphene citrate it is non steroidal antiestrogene and necessary to use 50 mg.in tablet twice a day after meal and glutargine 750 mg. at the same time 1 tablet twice a day with interval 12 hours. Alfa gonadotropine chorionic injection in dose 1500 IU once a day. For course of treatment 14 days. After that necessary to take blood test to specify if hyperestrogenism in man decreased during treatment. Also necessary to take biochemical test of liver enzymes to assesst functional possibilities of liver. Clomifene citrate usually well recover producing of testosterone and twice increase testosterone concentration in sufferers during two weeks of treatment. This is possible to confirm by blood test of hormones. Clomifene citrate has positive action for quality of sperm, especially when indicated olygospermia, decreased spermatozoa count in sperm. Since hormonal activity increased and it was confirmed in blood test it will be necessary to stop treatment as a first course.

Glutargine it is L-arginine glutamate, combination of arginine and glutamic acid used as hepatoprotector for treatment and improve functional opportunity of liver.

Table 1

Hormonal status in blood M±m, n=20

No	Hormone test	Normal range	Before treatment	After two weeks treatment
1.	Luteinizing hormone (LH), mIU/ml	1.24-8.62	1.3±0.24	12.2±1.26
2.	Follicle stimulating hormone (FSH), mIU/ml	1.27-19.26	3.1±0.62	16.7±1.82
3.	Prolactin (PRL) ng/ml	3.3-13.5	47.4±2.86	8.4±0.94
4.	Testosterone (T), ng/dl	262-870	164±5.43	428±3.12
5.	Estradiol (E2), pg/ml	up to 47	86±3.16	35±3.46
6.	Progesterone, ng/ml	0.15-1.97	2.29±0.47	0.23±0.07

Proteinic and steroid hormone activity in the blood of sufferers with hyperestrogenism presented in table 1. Before of treatment we tested hormonal activity for each sufferer. It showed concentration of luteinizing and follicle stimulating hormone and testosterone decreased. Instead of that increased level of prolactine, estradiol and progesterone. It is working very demonstratively negative feedback when plrolactine increases, luteinizing and follicle stimulsting hormone automatically decreases. Testosterone decrieses instead of it esradiol and progesterone increases. Only therefore we proposed to blocking secretion of estrogens and proposed antiestrogenic remedy clomifene citrate together with gonadotropine chorionic injections to improve actions in target organ of testicular tissue stimulating testosterone production without estrogene secretions. In this case we received optimal effect of hormone balancing in the body.

Antiestrogene non steroidal origin such as clomifene citrate with combination of alpha gonadotropine chorionic with direct action to target organ, testicle doing effectively treatment. Estrogen depending receptors inhibits during treatment with

clomifene citrate brings a number of endocrine system changes such as: decrease of secretion of estradiol, progesterone and prolactine. Instead of it increases secretion of luteinizing and follicle stimulating hormone and testosterone.

Table 2

Functional status of liver M±m, n=20

No	Enzyme test	Normal range	Before treatment	After two weeks treatment
1.	Lactate dehydrogenase (LDH)	115-450 IU/L	386±4.91	258±5.83
2.	Alanine aminotransferase (AlAT)	0-45 IU/L	37±2.48	24±1.27
3.	Aspartat aminotransferase (AsAT)	0-40 IU/L	32±2.07	20±1.43
4.	Gamma Glutamyl transferase (GGT)	11-61 IU/L	54±3.26	23±2.14
5.	Alcaline phosphatase (AF)	up to 115 IU/L	84±2.74	68±3.64

Before to use hormonal therapy we checked liver function and proposed in the course of treatment glutargine to protect liver and improve functional possibility of liver. Therefore due to glutargine improved estrogene conversion in liver. As showed test result of blood, activity of liver enzymes improved functional possibilities. It is visible in table 2, before and after two weeks treatment. Functional opportunity of liver is very important. Because estrogens reduce accumulation of triglycerides in the liver, stimulate lipolysis and improve the oxidation of fatty acids in mitochondria. In the fatty tissue produces peptidic hormone leptin. It stimulates production of releasing hormones which in turn activate synthesis of estrogens. Estrogens in the body eliminates via urine and bile. Therefore very important use hepatoprotective remedies and improve functional opportunities. Under the influence of estrogens activity of liver proteins (MDR3), which ensures the transport of phosphatidylcholine into the bile canaliculi decreses. All these factors determine the increase in bile lithogenicity. Including in the course of treatment glutargine 750.0 mg with hepatoptoprotective effect improve functional opportunity of liver. This is confirmed by blood test results of enzyme activity. Increasing concentration of estrogens can lead to decrease outflow of bile and accumulation of toxic bile acids (BA) which subsequently causes intrahepatic cholestasis.

Conclusions. Treatment with clomifen citrate in dose 50 mg twice a day in combination with alpha gonadotopine chorionic 1500 IU injections (Ovigil) once a day and glutargin 750 mg optimizes estrogens level in the body. Treatment during two weeks showed increasing testosterone, increased concentration of luteinizing hormone and follicle stimulating hormone. At the same time decreased concentration of estradiol, progesterone and prolactin. Hormonal condition in men stabilized, normalized enzyme activity in liver. Conversion estrogens in liver improved and stabilized in tested men.

BEE-HERBAL PRODUCTS AND PROBIOTICS COMBINATION FOR THERAPEUTIC ENHANCEMENT IN SKIN DISEASES

Michael Tarapata, Oleksandr Kukhtenko
National University of Pharmacy, Kharkiv, Ukraine
The Department of Pharmacy and Industrial Technology
automcorporation@gmail.com

Introduction. Acne (also known as Acne Vulgaris) is a common cutaneous inflammatory disorder of the pilosebaceous unit, which runs a chronic course that affects people all over the world. Acne exhibited a prevalence rate of 9.4% within the global population, ranking it eighth among skin diseases worldwide. About 87% of teenagers have this common skin problem, and the disease can persist into adulthood. Acne vulgaris is commonly observed in adolescents and young adults, but it can affect individuals of various ages.

Clinical presentation can vary from very mild facial comedonal lesions to more severe forms characterized by disfiguring inflammatory manifestations, which can lead to hyperpigmentation, scarring, adverse psychological effects, and other clinical sequelae. Hyperpigmentation is frequently seen, particularly in darker skin tones, and can persist for many months to years in some individuals. Currently, there's no 100% effective cure for acne, acne can't be stopped from occurring or cured, but it can be treated quite successfully.

As a result, there is a growing interest in alternative and complementary therapies, including herbal medicine, bees' products, and probiotics. This paper explores the complexity and potential benefits of these natural remedies in managing acne.

Aim. To substantiate the feasibility of a comprehensive approach to acne treatment using plant products, bee products, probiotics and potential advantages.

Materials and methods. Analytical, bibliosemantic.

Results and their discussion. Dermatology is a division of medicine, which constantly changes at a vast rate. Alongside technology, more complex medications, methods and treatments are continuously developed for the treatment of all common skin diseases. Acne is diagnosed by a simple visual inspection. Also a practitioner may take a swab or scraping of a lesion or pustule for microbiological examination or culture to rule out other sources of infection. The ways of assessing for acne severity grading including two main methods, namely, lesion counting and comparison of patient with a photographic standard for medical records. Next one the computational assessment method, the emphasis is on computational imaging techniques of measurements using algorithms. The Global Acne Grading System (GAGS) was first developed in 1997. The GAGS divides the face (forehead, each cheek, nose, and chin), chest, and back into six areas, and the severity in each zone is then assessed on a scale of 0 to 4 (0, no lesions; 1, comedones; 2, papules; 3, pustules; and 4, nodules).

Classification of Acne Severity by severity:

Mild - < 20 comedones, or < 15 inflammatory lesions, or < 30 total lesions.

Moderate – 20 to 100 comedones, or 15 to 50 inflammatory lesions, or 30 to 125 total lesions.

Severe - > 5 cysts, or total comedone count > 100, or total inflammatory lesion count > 50, or > 125 total lesions.

The pathogenesis of acne occurring by dysbiosis, and it's refers to an imbalance of microbial species and a reduction in microbial diversity within certain bodily microbiomes - i.e., the accumulation of microorganisms existing in a specific part of the human body. The human gut microbiota plays a significant role in maintaining gut-skin homeostasis, and whole body. Disruption of this equilibrium is characterized by the overgrowth of some bacterial species, pathogens, fungal and an overall loss of community diversity. This dysbiosis can lead to chronic diseases impaired wound healing, increased inflammation, and greater risk for diverse infection in the body and on the skin. Dysbiosis take to metabolic deregulation that affect immunologically the the skin barrier. Acne develops when dead skin builds up it and block the pores - hair follicles and trap the sebum beneath the skin's surface. Then bacterias can trigger inflammation and infection resulting in more severe acne or infections. There are many additional treatments acne, actual additional alternative of treatment included herbal combinations with propolis and fermented products. Propolis - a resinous substance collected by bees from plants, it exhibits strong antimicrobial, antioxidant, and anti-inflammatory properties. Inhibits the growth of acne-causing bacteria*.

The complexity of herbal and bee product action in acne management can be attributed to their diverse phytochemical profile. Anti-inflammatory components from herbal extracts can inhibit the activity of pro-inflammatory cytokines, while antibacterial agents can reduce microbial population of infections. Moreover, many herbal extracts demonstrate antioxidant properties that combat oxidative stress, a contributor to inflammatory processes in acne.

In ancient times, fermentation was used to boosts the food's shelf life and to increase the new quality of food. At present, the process has gained popularity due to its immense health benefits- healthy probiotics, nutritional value. Naturally fermented foods are getting a lot of attention from health experts as dietary supplements promoting health of gut microbiome - live microorganisms crucial to healthy digestion. Researchers are beginning to link these tiny creatures to all sorts of health conditions from metabolism deregulation, skin diseases to obesity to neurodegenerative diseases. Pre- and probiotics that is a line of research having many promising prospects on the gut-skin axis and other diseases.

Conclusions. Despite the promising benefits of herbal, bee products and fermented products, there is a need for rigorous scientific evaluation to establish their safety and efficacy. While many natural substances are generally well-tolerated, allergic reactions and skin sensitivities may occur. Moreover, varying concentrations of active ingredients across different products can lead to inconsistent results. Clinical trials and standardized formulations are essential to ascertain dosage and long-term safety.

ПЕРЕВАГИ КОМБІНОВАНОЇ ТЕРАПІЇ ЦУКРОВОГО ДІАБЕТУ КОМБІНАЦІЄЮ МЕТФОРМІНУ З ІНГІБІТОРАМИ НАТРІЙ-ГЛЮКОЗНОГО КОТРАНСПОРТЕРА 2-ГО ТИПУ

Деримедвідь Л. В.[1], Риженко І. М.[2], Богуцька О. Є.[3], Кузенков Р. В.[2]

[1]Національний фармацевтичний університет, м. Харків, Україна

[2]Інститут підвищення кваліфікації спеціалістів фармації НФаУ

[3]Херсонський державний університет, м. Івано-Франківськ, Україна

clinpharmacol_ipksph@nuph.edu.ua

Вступ. У структурі ендокринних захворювань цукровий діабет (ЦД) входить до трійки найбільш поширенних ендокринних патологій і посідає або перрше або друге місце після захворювань щитоподібної залози. Майже 42 млн людей у світовій спільноті хворіють на ЦД. В Україні кількість хворих на ЦД дорівнює 1,4 млн, проте ще є значна кількість людей з ЦД, які мають недіагностовану хворобу. Пацієнти з ЦД I типу (інсулінзалежний) складають лише 5-10% від загальної кількості хворих ЦД. ЦД II типу (інсуліннезалежний) складає 85-90% усіх випадків ЦД і зустрічається переважно серед осіб старшої вікової групи. Подальше стрімке зростання ЦД II типу в усьому світі дозволяє схарактеризувати його як пандемію сторіччя. У теперішній час спостерігається систематичне оновлення лікарських засобів, які використовуються в схемах фармакотерапії ЦД II типу, причому досить часто акцент робиться на фіксовані комбінації гіпоглікемічних препаратів. Це дозволяє лікам із різними механізмами дії впливати на різноманітні патогенетичні ланки захворювання, доповнюючі фармакологічні ефекти, зменшувати дози, покращувати контроль глікемії, знижувати частоту побічних реакцій та підвищувати комплаєнс хворих.

Мета дослідження. Проаналізувати дію деяких фіксованих комбінацій бігуанідів з інгібіторами натрій-глюкозного котранспортера 2-го типу, зареєстрованих в Україні.

Методи дослідження. Аналітичний, бібліосемантичний.

Результати дослідження. Серед значної кількості різноманітних комбінацій цукрознижувальних засобів насамперед привертають до себе увагу фіксовані комбінації метформіну з інгібіторами натрій-глюкозного котранспортера 2-го типу (ІНГКТ -2): дапагліфлозином (Ксигдуо пролонг) або з емпагліфлозином (Сінджарді).

Гіпоглікемічна дія метформіну обумовлена такими основними механізмами, як пригнічення глюконеогенезу в печінці, підвищення чутливості рецепторів скелетних м'язів до інсуліну, покращення захоплення глюкози периферичними тканинами та її утилізація, а також сповільнення всмоктування глюкози в кишківнику. Незалежно від його впливу на рівень глюкози, препарат покращує ліпідний профіль, зменшує масу тіла та артеріальний тиск у хворих з ЦД та артеріальною гіпертензією. Метформін не впливає на бета- клітини підшлункової залози, і тому не стимулює секрецію інсуліну і не призводить до розвитку гіпоглікемічного стану. Метформін знижує смертність та ризик ро-

звитку ускладнень серцево-судинних захворювань у хворих з ЦД II типу та ожирінням.

Зовсім інші механізми гіпоглікемічної дії в ІНГКТ -2, які пригнічують реабсорбцію відфільтрованої глюкози в проксимальних канальцях нефрона та знижують нирковий поріг для глюкози шляхом інгібування SGLT2, який експресується на високому рівні в нирках (експресія в інших тканинах відсутня). Це призводить до збільшення екскреції глюкози з сечею, осмотичному діуретичному ефекту, а також втраті калорій, що пов'язано зі зниженням рівня жирів і зменшенням маси тіла. Подібно метформіну, препарати даної групи сприяють зниженню ризику гіпоглікемії. Крім того, знижується реабсорбція натрію, що супроводжується помірним діурезом і зменшенням перед- та постнавантаження на серце, що сприяє зниженню ригідності артерій та артеріального тиску хворих з ЦД та артеріальною гіпертензією. Використання ІНГКТ -2 сприяє зниженню серцево-судинної смертності.

Взагалі глюкозурія, натрійурез та осмотичний діурез, які виникають внаслідок патофізіологічних механізмів дії ІНГКТ -2 сприяють покращенню серцево-судинних прогнозів у хворих з даною патологією та надлишковою масою тіла. Проте за даними досліджень останніх років, є певні відмінності щодо ступеня ремоделювання серця під впливом різних препаратів з групи ІНГКТ -2. Так, терапія емпагліфлозином у хворих з ЦД II типу та серцевою недостатністю, асоціюється зі збільшенням фракції викиду лівого шлуночка у порівняні з дапагліфлозином. За даними літератури, застосування емпагліфлозину на 20%-30% зменшує відносний ризик смерті від серцево-судинних захворювань або госпіталізації у дорослих пацієнтів із серцевою недостатністю зі збереженою фракцією викиду. Також емпагліфлозин є засобом першої лінії у пацієнтів із ЦД II типу та хронічною хворобою нирок (ХНН).

Висновки. Таким чином фіксовані комбінації бігуанідів і ІНГКТ -2, які мають різні механізми дії, дозволяють значно краще посилити глікемічний контроль хворих з ЦД II типу.

Препарати із фіксованим комбінаціям бігуанідів і ІНГКТ -2, доцільно призначати для зниження ризику госпіталізацій через серцеву недостатність у дорослих з ЦД II типу та встановленими серцево-судинними захворюваннями, або множинними факторами ризику з боку серцево-судинної системи та/або ХНН.

ВИБІР НЕСТЕРОЇДНИХ ПРОТИЗАПАЛЬНИХ ЗАСОБІВ МІЖ ВІТЧИЗНЯНИМИ ТА ЗАКОРДОННИМИ ПРЕПАРАТАМИ

Севастьянова Т. В., Яшна П. О., Антонов В. С.
Херсонський державний університет, Херсон, Україна
Медичний факультет, кафедра хімії та фармації
p.yashna.222@gmail.com

Анотація. Нестероїдні протизапальні препарати (НПЗП) є важливими фармацевтичними засобами, широко використовуваними для зняття болю та запалення. Проблема вибору між вітчизняними та закордонними НПЗП в умовах сучасного фармацевтичного ринку України є актуальною, оскільки на вибір споживачів впливають різні економічні та соціальні фактори. Метою дослідження було проаналізувати вплив ефективності, ціни, доступності та інших факторів на вибір НПЗП серед українських споживачів. Дослідження проводилося методом анкетування, в якому взяли участь 101 респондент. Результати показали, що основними критеріями вибору є ефективність та ціна, а найбільш популярними препаратами залишаються ібупрофен, парацетамол та німесил. Ці результати допомагають розробляти стратегії маркетингу та управління в фармацевтичній галузі.

Ключові слова: *Нестероїдні протизапальні препарати, споживчий вибір, ефективність, ціна, доступність, фармацевтичний ринок, маркетинг, Україна.*

Abstract. Nonsteroidal anti-inflammatory drugs (NSAIDs) are important pharmaceutical agents widely used to relieve pain and inflammation. The problem of choosing between domestic and foreign NSAIDs in the conditions of the modern pharmaceutical market of Ukraine is relevant, since various economic and social factors influence the choice of consumers. The purpose of the study was to analyze the influence of effectiveness, price, availability and other factors on the choice of NSAIDs among Ukrainian consumers. The research was conducted by the questionnaire method, in which 101 respondents took part. The results showed that the main selection criteria are effectiveness and price, and the most popular drugs remain ibuprofen, paracetamol and nimesil. These results help to develop marketing and management strategies in the pharmaceutical industry.

Keywords: *Nonsteroidal anti-inflammatory drugs, consumer choice, effectiveness, price, availability, pharmaceutical market, marketing, Ukraine.*

Вступ: Нестероїдні протизапальні препарати (НПЗП) належать до числа найбільш поширених лікарських засобів, особливо враховуючи безрецептурний режим їхнього продажу в аптечних закладах. Нестероїдні протизапальні засоби досить часто застосовуються в медичній практиці, оскільки мають виражені протибольові та протизапальні ефекти [1]. Загалом відмічається тенденція до збільшення вживання НПЗП у 2-3 рази кожні 10 років. За даними ВООЗ, кожний третій мешканець планети регулярно використовує НПЗП [2].

Зараз пропонується широкий асортимент таких препаратів, які представлені як вітчизняними, так і закордонними виробниками. В Україні зареєстровані декскетопрофен, диклофенак, ібупрофен, кеторолак, кетопрофен, піроксикам, німесулід, етодолак, мелоксикам, кислота мефенамінова, целекоксиб, парекоксиб, рофекоксиб, діацереїн тощо[3,4,7]. Міжнародний досвід регулювання фармацевтичного сектору спонукає до орієнтації на існуючих та потенційних закордонних партнерів. Так, на державному рівні визначена стратегія руху України до Європейського Союзу, яка передбачає удосконалення вітчизняного нормативного забезпечення щодо виробництва і контролю якості ліків до стандартів прийнятих в ЄС, що у свою чергу, виробникам України дасть змогу виходити із своєю фармацевтичною продукцією на ринок ЄС та США [5,6].

Мета: Дослідити стратегії вибору нестероїдних протизапальних лікарських препаратів. Проаналізувати уподобання споживачів та вплив управлінсько-організаційних, маркетингових та соціально-економічних факторів на вибір між вітчизняними та закордонними продуктами в фармацевтичній галузі.

Матеріали та методи дослідження: Дослідження проводилось за допомогою власноруч розробленого опитувальника, який включав питання про фактори, які впливають на вибір нестероїдних протизапальних препаратів. У анкетуванні взяли участь 101 респондент.

Результати та їх обговорення: За результатами опитування встановлено: частка жінок становить 85,1%, чоловіків – 14,9% (рис. 1).




Рис. 1 Розподіл учасників опитування за статтю.

Більшість респондентів (70,3%) належать до вікового діапазону 16-30 років (рис. 2).




Рис. 2 Розподіл учасників опитування за віком.

Відсоток студентів/працівників медичної сфери складає 69,3% (рис. 3).




Рис. 3 Розподіл учасників опитування за сферою діяльності.

Суб'єктивний рівень обізнаності в сфері НПЗП за п'ятибальною шкалою: 1- 5%; 2- 12,9%; 3- 45,5%; 4- 27,7%; 5- 8,9% (рис. 4).

Віддають перевагу вітчизняним НПЗП — 22,8%, закордонним — 27,7%, однак для майже половини опитуваних осіб країна-виробник не має значення (49,5%) (рис. 5).




Рис. 4 Суб'єктивний рівень обізнаності опитуваних у сфері НПЗП за п'ятибальною шкалою.




Рис. 5 Надання переваги вибору НПЗП за країною-виробником.

Основними факторами, що впливають на вибір конкретного НПЗП (респондентам дозволялось обрати декілька відповідей одразу) є: ефективність — 88,1%, ціна — 57,4%, доступність у місцевості — 46,5%, форма випуску — 43,6%, рекомендації лікаря — 41,6%, країна виробник — 24,8% (рис. 6).




Рис. 6 Фактори, що впливають на вибір конкретного НПЗП.

Найбільша довіра до препаратів (респондентам дозволялось обрати декілька відповідей) Німеччини (72,3%), на другому місці Україна (62,4%), третє — Швейцарія (39,6%); далі в рейтингу займають місця США (28,7%), Франція (27,7%), Бельгія (22,8%), Нідерланди (16,8%), Італія (14,9%), Японія (10,9%), Іспанія (8,9%), Індія (7,9%), Ірландія (6,9%), Китай (2%) відповідно (рис. 7).




Рис. 7 Розподіл опитуваних за рівнем довіри до НПЗП країн виробників препаратів.

Перше місце серед найпопулярніших НПЗП (респонденти могли обрати декілька відповідей одразу) займає ібупрофен (70,3%), друге — парацетамол (56,4%), третє — німесил (51,5%); далі в рейтингу займають місця кетопрофен

(10,9%), дексалгін (7,9%), кеторолак (6,9%), декскетопрофен (5%), лорноксикам та налгезін (4%), ортофен та індометацин (3%), піроксикам разом з напроксеном та ацеклофенаком (2%), артоксан (1%), інші(6,9%) (рис. 8).




Рис. 8 Популярність НПЗП при виборі опитуваними.

Ситуація в країні, серед опитуваних, значно не впливає (44,6%), скоріше впливає чим ні (29,7%), значно впливає (25,7%) на вибір НПЗП (рис. 9).




Рис. 9 Вплив економічної ситуації в країні на вибір НПЗП.

Реклама та маркетинг у 15,8 % випадків впливає на вибір НПЗП (рис. 10).




Рис. 10 Вплив реклами та маркетингу на вибір НПЗП**.**

Висновки: Згідно з результатами нашого дослідження, основними факторами, які впливають на вибір нестероїдних протизапальних лікарських засобів, є їх ефективність, ціна та доступність у місцевості. Ці відомості можуть бути важливими для розробки стратегій управління та маркетингу. На даний момент основною проблемою, що перешкоджає цьому, є економічна ситуація в країні та менша доступність закордонних засобів.

Також, варто звернути увагу на активну участь молодої аудиторії в опитуванні, що може свідчити про необхідність розробки ефективних маркетингових стратегій, спрямованих на цей сегмент споживачів.

Висока довіра до препаратів виробництва Німеччини та України підкреслює значення якості та довіри до виробників у виборі медичних препаратів. Що стосується самого вибору препаратів, ібупрофен, парацетамол та німесил виявилися найпопулярнішими серед респондентів.

Отже, залучення даних з управлінсько-організаційних, маркетингових та соціально-економічних досліджень є ключовим аспектом у формуванні стратегій вибору нестероїдних протизапальних лікарських засобів у фармацевтичній галузі.

Перелік використаних джерел інформації:

1. Малик Н. В., Клименко Д. Ю., Гамаюн А. М. (2023, January). Застосування нестероїдних протизапальних засобів в практиці сімейного лікаря. In The 1 st International scientific and practical conference “Innovations and prospects in modern science”(January 15-17, 2023) SSPG Publish, Stockholm, Sweden. 2023. - С. 75 - 77.
2. Шупер В. О., Дорогокупець А. В. «Аналіз особливостей застосування нестероїдних протизапальних препаратів». The 11th International scientific and practical conference “Eurasian scientific discussions” (November 21-23, 2022) Barca Academy Publishing, Barcelona, Spain. 2022. - С. 79 - 82.
3. Настанова МОЗ 00883. (08.09.2017). Безпечне застосування нестероїдних протизапальних препаратів (НПЗП).

4. Наказ Міністерства охорони здоров'я України від 22 квітня 2021 року №792 «Про затвердження тринадцятого випуску Державного формуляра лікарських засобів та забезпечення його доступності», С. 387.
5. Овчаренко Т., Єрмолаєва В. Механізм впровадження міжнародних стандартів GMP у фармацевтичний сектор економіки України. Вісник Київського національного університету імені Тараса Шевченка, 2009, С. 90-93.
6. British National Formulary 80 (BNF), September 2020 — March 2021, за ред. Pharmaceutical Press, Great Britain, 2020.
7. Державний реєстр лікарських засобів України. Доступно за посиланням: http://www.drlz.com.ua/

РОЛЬ РАНОЛАЗИНУ ЯК АНТИАРИТМІЧНОГО ПРЕПАРАТУ В СУЧАСНІЙ МЕДИЧНІЙ ПРАКТИЦІ

Русанова А. І.
КНП ХОР Обласна клінічна лікарня, м. Харків, Україна
anastasiaa.rusanova@gmail.com

Вступ. Аритмії – це порушення серцевого ритму, які можуть мати різну природу і ступінь тяжкості від незначних перебоїв до небезпечних для життя станів. Ефективне лікування аритмій є одним із головних викликів у сучасній кардіології, оскільки традиційні антиаритмічні препарати часто мають побічні ефекти та обмежену ефективність. У цьому контексті інтерес до нових методів лікування постійно зростає.

Ранолазин — це лікарський засіб спочатку розроблений для лікування хронічної стенокардії, але дослідження показали, що він має значний антиаритмічний потенціал. Цей лікарський засіб відрізняється своїм унікальним механізмом дії, що робить його корисним у клінічній практиці, особливо у випадках коли стандартні антиаритмічні препарати мають обмежену ефективність або викликають небажані побічні ефекти. Особливий інтерес викликає його можливе застосування за схемою «таблетка у кишені», коли препарат приймається пацієнтом при перших ознаках аритмії для швидкого купірування епізоду.

Мета. Дослідити ефективність ранолазину як антиаритмічного препарату, зокрема його застосування в ролі «таблетка у кишені» для купування пароксизмальних епізодів фібриляції передсердь. Вивчити його вплив на відновлення синусового ритму, частоту серцевих скорочень, побічні ефекти, а також оцінити безпеку і переносимість препарату у порівнянні з іншими антиаритмічними засобами.

Матеріали та методи. Дослідження базувалося на аналізі літератури та клінічних випробувань де ранолазин використовувався для лікування аритмій. Основним об'єктом дослідження є пацієнти з пароксизмальною формою фібри-

ляцією передсердь. Оцінювалися дози препарату прийняті пацієнтами при появі симптомів аритмії. Аналізувалися результати ЕКГ для моніторингу відновлення нормального ритму, а також оцінювалися побічні ефекти препарату (особливо зміни інтервалу QT). Методологія включала ретроспективний огляд клінічних досліджень та спостереження за пацієнтами, які використовували ранолазин у різних дозуваннях – оцінювалися швидкість відновлення синусового ритму, побічні ефекти, а також додаткове лікування для стабілізації стану.

Результати. Основний механізм дії ранолазину пов'язаний з інгібуванням пізньої фази натрієвого потоку. Пізній натрієвий потік виникає при різних патологічних станах, таких як ішемія міокарда та/або серцева недостатність, коли спостерігається значне збільшення кількості пізніх натрієвих каналів на поверхні кардіоміоцитів. Як наслідок, посилений пізній потік натрію викликає підвищення внутрішньоклітинної концентрації натрію. У відповідь на це активується обмін натрій-кальцій у зворотному режимі та викликає підвищення концентрації кальцію. У кардіоміоцитах це може погіршити механічну релаксацію та викликати електричну нестабільність, подовжуючи потенціал дії, що потенційно може викликати шлуночкові аритмії, включаючи «torsades de pointes». Ранолазин селективно інгібує пізній потік натрію у кардіоміоцитах, тим самим обмежуючи надлишок натрію та накопичення кальцію. Це призводить до зменшення діастолічного напруження стінки та покращення коронарного кровотоку, тим самим сприяючи зниженню виникнення ішемії, аритмій та покращенню загальної функції серця.

Завдяки своєму механізму дії ранолазин проявляє антиаритмічну дію, яка стосується кількох типів аритмій. Особливу увагу приділяють його ефективності при лікуванні шлуночкових аритмій, які можуть стати причиною раптової серцевої смерті, та фібриляції передсердь – одного з найпоширеніших типів порушення ритму серця. У дослідженнях було показано, що препарат допомагає відновити нормальний синусовий ритм у пацієнтів з персистуючою або пароксизмальною фібриляцією передсердь, а також знижує частоту повторних епізодів цієї аритмії.

Ранолазин має потенціал для використання як «таблетка у кишені» при лікуванні пароксизмальних аритмій, зокрема фібриляції передсердь. Дослідження показали, що прийом ранолазину в дозах 1000-2000мг виявився ефективним і безпечним для купування епізодів фібриляції передсердь у більшості пацієнтів. У 70-80% випадків нормальний синусовий ритм був відновлений протягом кількох годин після прийому препарату. Однак, через появу у деяких пацієнтів побічних ефектів, таких як запаморочення, нудота і незначне подовження інтервалу QT, що потребувало моніторингу, цей підхід ще потребує подальших досліджень і клінічної перевірки, перш ніж він буде широко застосовуватись у медичній практиці.

Ранолазин має кілька переваг перед традиційними антиаритмічними засобами:

1. Відсутність значного впливу на частоту серцевих скорочень і артеріальний тиск. Багато антиаритмічних препаратів можуть викликати зниження артеріального тиску та частоти серцевих скорочень, що обмежує їх використання в певних групах пацієнтів. Ранолазин виявився нейтральним щодо цих параметрів, що дозволяє використовувати його у ширшого кола хворих, зокрема у пацієнтів з артеріальною гіпотензією або брадикардією.

2. Мінімальний ризик проаритмічних ефектів. Більшість антиаритмічних препаратів (особливо класу I та III) мають проаритмічні властивості, тобто можуть викликати нові або загострювати наявні аритмії. Ранолазин демонструє низький ризик проаритмічних ефектів, що робить його безпечнішим варіантом для лікування пацієнтів з високим ризиком аритмій.

3. Комбінаційна терапія. Ранолазин може застосовуватися як монотерапія, так і в комбінації з іншими антиаритмічними препаратами, що дозволяє підвищити ефективність лікування складних випадків аритмій. Проте, така комбінація потребує ретельного контролю, особливо через можливе подовження інтервалу QT.

Ранолазин зазвичай добре переноситься пацієнтами. Однак, як і будь-який лікарський засіб, він має певні побічні ефекти, які можуть виникати у деяких пацієнтів. Найчастіше відзначаються такі симптоми, як запаморочення, головний біль, нудота і слабкість. У рідкісних випадках можливе подовження інтервалу QT на ЕКГ, що може спричинити серйозні аритмії. Тому пацієнти, що мають фактори ризику подовження інтервалу QT, потребують ретельного контролю при прийомі препарату.

Висновки. Ранолазин у майбутньому може бути перспективним, ефективним і безпечним антиаритмічним препаратом, який завдяки своєму специфічному механізму дії має значний потенціал у лікуванні різних типів аритмій, зокрема шлуночкових аритмій та фібриляції передсердь. Його ключові переваги включають мінімальний вплив на частоту серцевих скорочень і артеріальний тиск, а також низький ризик проаритмічних ефектів. На даний момент точна доза ранолазину для використання як «таблетки у кишені» при аритміях ще не затверджена жодними міжнародними рекомендаціями. Однак є дані з досліджень, які вказують на можливі ефективні дози, що використовувалися при лікуванні пароксизмальної фібриляції передсердь. У деяких випадках ранолазин використовувався в комбінації з антиаритмічними препаратами для підвищення ефективності лікування. Водночас необхідно враховувати можливі побічні ефекти та проводити регулярний моніторинг ЕКГ у пацієнтів, які мають фактори ризику.

УДК 616.348-008.1-085.246

ПОШУК ДОКАЗОВОЇ БАЗИ КЛІНІЧНОЇ ЕФЕКТИВНОСТІ ПРОБІОТИКІВ ПРИ ЛІКУВАННІ ГОСТРОЇ ДІАРЕЇ

Ткачова О. В.[1], Белік Г. В.[2]

Національний фармацевтичний університет, м. Харків, Україна

[1]Кафедра клінічної фармакології ІПКСФ

[2]Кафедра фармакології та клінічної фармації

tkachevaov@gmail.com

Анотація. У статті представлені результати аналізу доказової бази клінічної ефективності різних пробіотиків: B. lactis, L. rhamnosus, S. boulardii, L. acidophilus та B. bifidum, що застосовують для лікування гострої діареї у дітей та дорослих. Для проведення аналізу використовували бази даних доказової медицини: Cochrane Library та PubMed. Отримані результати 11 систематичних оглядів та метааналізів виявились неоднозначними. Більш ефективному скороченню тривалості гострої діареї у дітей та дорослих сприяло поєднане застосування декількох штамів пробіотиків, тобто синбіотиків. Застосування лише одного штаму пробіотика Lactobacillus acidophilus виявилось не ефективним при лікуванні гострої діареї.

Abstract. The article presents the results of the analysis of the evidence base of the clinical effectiveness of various probiotics: B. lactis, L. rhamnosus, S. boulardii, L. acidophilus and B. bifidum, used for the treatment of acute diarrhea in children and adults. Databases of evidence-based medicine: Cochrane Library and PubMed were used for the analysis. The results of 11 systematic reviews and meta-analyses were mixed. A more effective reduction in the duration of acute diarrhea in children and adults was facilitated by the combined use of several strains of probiotics, that is, synbiotics. The use of only one strain of the probiotic Lactobacillus acidophilus was not effective in the treatment of acute diarrhea.

Вступ. Згідно з визначенням Всесвітньої організації охорони здоров'я (ВООЗ), основним критерієм діарейного синдрому є збільшення добової кількості калових мас до 200 г (поліфекалії) на тлі підвищення відсоткового вмісту води в екскрементах хворого до 85-95% (у нормі 60-75%) [1].

Дані статистичних досліджень ВООЗ свідчать про те, що діарея є найчастішою причиною звернення пацієнтів за медичною допомогою. У представників дорослого населення країн Європи та США зустрічається в середньому по одному епізоду гострої діареї на рік. У світі щорічно реєструються 1,5-1,7 млрд випадків діареї різної етіології. Ця патологія пов'язана з високими економічними витратами і посідає друге місце у світі серед причин швидкої смерті, яка настає протягом 2-3 днів від початку захворювання [2].

Діарею класифікують за тривалістю перебігу:

• гостра (тривалістю до 14 днів);
• хронічна (тривалістю понад 2-3 тижні).

Провідним етіологічним фактором гострої діареї є гострі кишкові інфекції. Хронічна діарея носить неінфекційний характер, яка найчастіше зумовлена функціональними розладами кишечника, а також хронічними захворюваннями тонкої кишки, підшлункової залози, ендокринопатіями, запальними захворюваннями товстої кишки [3].

У педіатричній практиці при гострій вірусній та бактеріальній діареї в комплексному лікуванні доведено ефективність пробіотичних та синбіотичних препаратів на основі Lactobacillus rhamnosus, Lactobacillus acidophilus, Saccharomyces boulardii, Bacillus clausii, Bacillus lactis, у дорослих – Enterococcus faecium, Bacillus lactis, а також Lactobacillus rhamnosus, Lactobacillus acidophilus, Saccharomyces boulardii. Ефект пробіотиків пояснюється тим, що мікроорганізми, які входять до їх складу, заміщають функції власної нормальної кишкової мікрофлори [3, 4].

Відомо, що лікування антибіотиками порушує мікрофлору шлунково-кишкового тракту, що викликає низку клінічних симптомів, насамперед діарею. Це особливо важливо у дітей та дорослих, яким часто призначають антибіотики. Пробіотики широко використовуються в різних клінічних ситуаціях. В усьому світі пробіотики широко рекомендують для використання, зокрема після призначення антибіотиків [4].

Відомо, що фарм- і медпредставники, які просувають серед лікарів та фармацевтів лікарські засоби своєї фармацевтичної компанії, пропонують ознайомитися з інформацію про отримані результати клінічного дослідження та вважають її достовірною. Однак, в окремих рандомізованих клінічних дослідженнях (РКД) та оригінальних наукових статтях часто не міститься доведеної вторинної інформації та відповідей на клінічне питання щодо ефективності певного лікарського засобу для лікування певного захворювання, яке заявляється медичними представниками.

Знайти авторитетну відповідь на подібне питання можна лише за даними систематичних оглядів та метааналізів, які містять результати РКД з якісною методологією з вивчення конкретного препарату для лікування конкретного захворювання [5]. Щоб лікар/фармацевт не шукав таку інформацію особисто, вона внесена до вітчизняних та міжнародних стандартів лікування (накази керівництва) певного захворювання, на основі яких розроблено уніфіковані клінічні протоколи медичної допомоги хворим.

В опублікованих доступних наукових вітчизняних дослідженнях дуже мало представлено інформації щодо доказової бази клінічної ефективності пробіотиків при лікуванні діареї [4, 6].

У міжнародних виданнях опубліковано чимало клінічних досліджень для оцінки здатності пробіотиків запобігати викликаній антибіотиками діареї в загальній популяції, однак, мало відомо про те, які штами та дози пробіотиків можуть принести найбільшу користь дітям та дорослим.

Мета дослідження. Зібрати, узагальнити та проаналізувати доказову базу клінічної ефективності різних штамів пробіотиків, в тому числі штаму Lactobacillus acidophilus при лікуванні та для профілактики гострої діареї у дітей та дорослих.

Матеріали та методи дослідження. Для пошуку доказової бази та аналізу клінічної ефективності пробіотиків при лікуванні гострої діареї застосовували відомі бази даних доказової медицини: Cochrane Library та PubMed, що містять систематизовану первинну та/або вторинну інформацію з певного клінічного питання.

Результати та їх обговорення. В проведеній роботі було проаналізовано 11 систематичних оглядів (СО) та метааналізів. Загальна кількість учасників, включених до проведеного аналізу, становила понад 10 тисяч осіб.

Отримані результати аналізу доказової бази пробіотиків для профілактики та лікування гострої діареї у дітей та дорослих наведені у таблиці.

Ефективність різних штамів пробіотиків, в тому числі й L. acidophilus разом з іншими штамами пробіотиків вивчена у численних рандомізованих клінічних випробуваннях (РКВ), які були опубліковані протягом 1996-2021 років. Отримані результати показали, що розширення використання та різноманітність пробіотиків для вибору відповідного типу пробіотичних препаратів стає складним завданням як для медичних працівників, так і для населення.

На сьогодні в багатьох СО та метааналізах підтверджені позитивні ефекти пробіотиків в цілому та зокрема штамів B. lactis, L. rhamnosus, S. boulardii та L. acidophilus. B. bifidum щодо нормалізації функції кишечнику, лікування та профілактики діареї, в тому числі ААД. Але, при виборі відповідного пробіотика часто ігноруються два важливих фактора – специфічність пробіотичного штаму та специфічність захворювання для оцінки його ефективності. Огляди та метааналізи часто поєднують різні типи пробіотиків, що призводить до помилкових висновків про ефективність препаратів. Успіх широкого клінічного застосування пробіотиків, особливо у педіатрії, полягає не тільки в їх антидисбіозній ефективності, але й у визнанні цього методу як одного з найбільш нешкідливих у сучасній медицині.

Висновки. Проведений пошук доказової бази пробіотиків дозволив виявити 11 систематичних оглядів щодо ефективності та безпеки його застосування під час лікування діареї у дорослих та дітей. Загальна кількість пацієнтів, що були включені до систематичних оглядів, проведених протягом 1996-2021 років, становила понад 10 тис. осіб. Численні дослідження показали, що застосування пробіотиків штамів B. lactis, L. rhamnosus, S. boulardii, L. acidophilus та B. bifidum при гострій діареї (як інфекційної, так і небактеріальної) у немовлят, дітей та дорослих виявилося недостатньо ефективним при застосуванні лише одного моноштаму пробіотика та більш ефективним у скороченні тривалості – при поєднанні декількох штамів або використанні синбіотиків.

Таблиця

Результати пошуку доказової бази ефективності пробіотиків для профілактики та лікування гострої діареї у дітей та дорослих

№ п/п	Назва дослідження	Вид дослід-ження, рік	Отримані результати клінічних досліджень
1	2	3	4
1.	Probiotics in the treat-ment and preven-tion of acute infectious diarrhea in infants and children: a systematic review of published randomized, double-blind, placebo-controlled trials.	Система-тичний огляд з мета-аналі-зом, 2001 [7]	*Мета:* визначити вплив пробіотиків у лікуванні та профілактиці гострої ін-фек-ційної діареї у немовлят та дітей. У дослід-женнях лікування викори-стовувалися такі штами пробіотиків: штам Lactobacillus (L.), L. reuteri, L. acidophilus та L. bulgaricus, Saccharomyces boulardii, Streptococcus ther-mophilus lactis. 17 Кокрановських СО віповідали критеріям включення. *Результати:* Діарея в дослідженнях лікування була бактеріальною, викли-каною ротавірусом людини, або мала не встановлене походження. Пробіо-тики значно скорочували тривалість гострої інфекційної діареї у немовлят і дітей, особливо діареї, спричиненої ротавірусом. Не вдалося зробити жод-них висновків про використання пробіотиків для профілактики діареї через клінічну та статистичну неоднорідність досліджень.
2.	Probiotics in prevention of antibiotic associated diarrhoea: meta-analysis.	Мета-аналіз, 2002 [8]	*Мета:* оцінити ефективність пробіотиків у профілактиці та лікуванні діареї, пов’язаної із застосуванням антибіотиків. *Результати:* включені 9 РКВ (n=1214), у яких пробіотична терапія для профілактики діареї порівнювалася з плацебо (обидва застосовувалися в комбінації з антибіотиками). Мета-аналіз показав, що пробіотики можна використовувати для запобігання діареї, пов’язаної з антибіотиками, і що S. boulardii і лактобацили можуть бути використані в цій ситуації. Остаточна ефективність пробіотиків при лікуванні діареї, пов’язаної з прийомом анти-біотиків, залишається недоведеною.

1	2	3	4
3.	Efficacy of probiotic use in acute diarrhea in children: a meta-analysis.	Мета-аналіз, 2002, [9]	*Мета:* визначити ефективність вико-ристання пробіотиків (у т.ч. і) у скорочен-ні тривалості підвищеного діурезу у дітей з гострим діарейним захворюванням. *Результати:* мета аналіз підтвердив ефективність пробіотика в скороченні тривалості симптомів гострої небак-теріальної діареї у дітей віком до 5 років. Пробіотики, особливо лактобацили, скорочували тривалість гострого діарейного епізоду у немовлят або дітей приблизно на один день.
4.	Probiotics for the Prevention of Antibiotic-Associated Diarrhea in Children.	Систематичний огляд з мета-аналізом 2016, [10]	*Мета:* оцінити ефективність про-біотиків у профілактиці антибіотико-асоційованої діареї (ААД) у дітей. Пошук було проведено у Medline, Embase та Кокранівській бібліотеці. У дослідження включено лише РКВ. *Висновки авторів:* включено 6 плацебо-контрольованих РКД (766 дітей). Лікування пробіотиками порівняно з плацебо знижувало ризик ААД з 28,5 до 11,9% (відносний ризик, ВР 0,44, 95% ДІ 0,25-0,77, модель випадкового ефекту). Таким чином, пробіотики знижують ризик ААД у дітей. На кожні 7 пацієнтів, у яких розвинеться діарея при лікуванні антибіотиками, на одного менше розвиватиметься ААД, якщо вони отримуватимуть пробіотики.
5.	A Meta Analysis of Probiotic Efficacy for Gastrointestinal Diseases	Мета-аналіз, 2012 [11]	*Мета:* оцінити ефекти пробіотиків для лікування та профілактики різних шлунково-кишкових захворювань (ШКЗ) з різною етіологією та механізмами дії. *Висновки:* пробіотики корисні при лікуванні та профілактиці ШКЗ. Ефективність не спостерігалася щодо діареї мандрівників, некротичного ентероколіту або пробіотичних видів L. acidophilus, L. plantarum та B. infantis. При виборі пробіотиків для лікування або профілактики ШКЗ, найбільш важливими факторами, які слід враховувати, є тип захворювання та вид (штам) пробіотиків

1	2	3	4
6.	Preventing pediatric antibiotic-associated diarrhea and Clostridium difficile infections with probiotics: A meta-analysis.	Мета-аналіз, 2013, [12]	*Мета:* оцінити ефективність та безпе-ку пробіотиків для профілактики: діареї, пов'язаної з прийомом антибіотиків, та інфекцій, викликаних Clostridiumdifficile (ІСД). *Результати:* пробіотики (всі штами) значно знижували частоту дитячої ААД (RR = 0,42, 95% CI: 0,33-0,53) і значно знижу-вали ІСД (ВР = 0,35, 95% ДІ: 0, 13-0,92), але ефективність значно варіюється в залежності від штаму пробіотика. Значно знижували ААД у дітей: штами Saccharomyces boulardii (ВР=0,43, 95% ДІ: 0,32-0,60) та Lactobacillus rhamnosus (ВР = 0,36, 95% ДІ: 0,19-0,69). В інших пробіотиків була дуже низька ефективність.
7.	What do Cochrane systematicreviews say about probiotics as preventive intervenetions?	Систематичний огляд з мета-аналізом 2017, [13]	*Мета:* дати відповіді на питання чи слід призначати пробіотики з профілактичною метою? Які типи пробіотиків слід використовувати в якій дозі та як довго? Узагальнено всі кокранівські СО про профілактичні ефекти пробіотиків у клінічній практиці. *Результати:* в жодному із СО не представлено доказів високої якості щодо профілактики захворювань за допомогою пробіотиків. Необхідні додаткові випробування, щоб краще дізнатися про пробіотики і підтвердити, коли їх використання є корисним та рентабельним.
8.	Strain-Specificity and Disease-Specificity of Probiotic Efficacy: A Systematic Review and Meta-Analysis	Систематичний огляд з мета-аналізом 2018, [14]	*Мета:* визначити, чи мають штами пробіотиків ефективність щодо діареї. Включено 228 РКД і знайдено докази як специфічності штаму, так і специфічності захворювання щодо ефективності різних штамів пробіотиків. Значні докази ефективності були виявлені для 7 (70%) пробіотичних штамів серед чотирьох профілактичних показань та 11 (65%) пробіотичних штамів серед п'яти показань до лікування. *Висновки:* штам-специфічна ефективність для запобігання діареї, пов'язаної з прийомом антибіотиків у дорослих, була чітко продемонстрована для Lactobacillus [наприклад, сумішшю Lactobacillus acidophilus CL1285, Lactobacillus caseiLBC80R та LactoBacillus + Lacto Bacillus. 14001 (Bio -K+)). Отже, ефективність пробіотиків залежить як від штаму, так і від хвороби.

1	2	3	4
9.	A meta-analysis of the effects of probiotics and synbiotics in children with acute diarrhea.	Мета-аналіз, 2019, [15]	*Метою* було визначення ефективності пробіотиків та синбіотиків при гострій діареї (ГД) у дітей, а також вивчалися склади пробіотиків, типи втручань. У метааналіз включено 34 РКВ з 4911 пацієнтами. *Висновки:* синбіотики були більш ефективними, ніж пробіотики, у скороченні тривалості діареї та госпіталізації, а Saccharomyces boulardii та Bifidobacterium bifidum були більш ефективними, ніж Lactobacillus, у скороченні тривалості діареї.
10.	Are probiotics and prebiotics safe for use during pregnancy and lactation? A systematic review and meta-analysis	Систематичний огляд з мета-аналізом 2021, [16]	*Мета:* оцінка даних про побічні ефекти пробіотиків, пребіотиків та синбіотиків під час вагітності та годування груддю. *Результати:* В 11 зі 100 РКД, проведених у 25 країнах, повідомлялося про незначні побічні ефекти. Побічні ефекти, пов'язані з використанням пробіотиків та пребіотиків, не становили серйозної небезпеки для здоров'я матері чи дитини.
11.	Probiotics for the Prevention of Antibiotic-Associated Diarrhea in Outpatients-A Systematic Review and Meta-Analysis	Систематичний огляд з мета-аналізом 2017, [17]	*Мета:* полягала в оцінці користі та шкоди пробіотиків, які використовуються для профілактики ААД в амбулаторних умовах. Було проведено пошук у базі даних PubMed і було отримано загалом 17 РКД із 3631 учасником, які були включені до огляду. Мета-аналіз було проведено для первинного результату: захворюваності на ААД. *Результати:* об'єднані результати виявили, що AAD був присутній у 8,0% групи пробіотиків порівняно з 17,7% у контрольній групі (ВР 0,49, 95% ДІ 0,36 до 0,66), а результати щодо видів були подібними щодо пробіотичні штами L. rhamnosus GG і S. boulardii . Проте загальна якість включених досліджень була помірною. Мета-аналіз десяти досліджень, у яких повідомлялося про побічні явища, не продемонстрував статистично значущих відмінностей у частоті побічних дій між групою втручання та контрольною групою. *Висновки:* використання пробіотиків є корисним та виглядає безпечним для профілактики ААД серед амбулаторних пацієнтів.

Перелік використаних джерел інформації:

1. Кушнір І.Е. Патофізіологічні аспекти синдрому діареї і підходи до патогенетичної терапії. Раціональна фармакотерапія. 2018. №2 (47). С. 36-42.
2. Усачова О.В., Печугіна В.В. Вплив цинку на розвиток інфекційної діареї у дітей. Педіатрія. 2020. № 4 (55) р.
3. Голубовська О.А., Галушко О.А. Гострі кишкові інфекції: сучасні методи інтенсивної терапії. Гострі та невідкладні стани у практиці лікаря. 2016. 2 (59). С. 15-18.
4. Скрипник І.М., Приходько Н.П. Антибіотикоасоційована діарея: роль пробіотиків у лікуванні та профілактиці. Укр. мед. часопис. 2021. Т. III/IV. №2 (142). С. 1-5.
5. Яковлєва Л.В., Кириченко О.М. Систематичний огляд. Фармацевтична енциклопедія. URL: https://www.pharmencyclopedia.com.ua/article/544
6. Радутна О. Властивості пробіотиків та їх вплив на організм людини. Тенденції та перспективи розвитку науки і освіти в умовах глобалізації: Збірник матеріалів Міжнародної наукової інтернет-конференції (30 вересня 2024 р.). С. 545-548.
7. Szajewska H., Mrukowicz J.Z. Probiotics in the treatment and prevention of acute infectious diarrhea in infants and children: a systematic review of published randomized, double-blind, placebo-controlled trials. J. Pediatr Gastroenterol Nutr. 2001; 33 Suppl 2:S.17-25.
8. D'Souza A.L., Rajkumar C., Cooke J.., Bulpitt CJ. Probiotics in prevention of antibiotic associated diarrhoea: meta-analysis. BMJ. 2002;324(7350):1361.
9. Huang J.S., Bousvaros A., Lee J..W, Diaz A., Davidson E.J. Efficacy of probiotic use in acute diarrhea in children: a meta-analysis. Dig Dis. Sci. 2002;47(11):2625-34.
10. Szajewska H., Canani R.B., Guarino A., Hojsak I., Indrio F., Kolacek S., Orel R., Shamir R., Vandenplas Y., van Goudoever J.B., Weizman Z. ESPGHAN Working Group for ProbioticsPrebiotics. Probiotics for the Prevention of Antibiotic-Associated Diarrhea in Children. J. Pediatr. Gastroenterol Nutr. 2016;62(3):495-506.
11. Ritchie M.L., Romanuk TN. A meta-analysis of probiotic efficacy for gastrointestinal diseases. PLoS One. 2012;7(4):e34938.
12. McFarland L.V., Goh S. Preventing pediatric antibiotic-associated diarrhea and Clostridium difficile infections with probiotics: A meta-analysis. World J. Meta-Anal. 2013. Vol. 1(3). P. 102-120.
13. Braga V.L., Rocha L.S., Bernardo D.D., Cruz C.O., Riera R. What do Cochrane systematic reviews say about probiotics as preventive interventions? Sao Paulo Med J. 2017;135(6):578-586.
14. Lynne V. McFarland, Charlesnika T. Evans and Ellie J. C. GoldsteinStrain-Specificity and Disease-Specificity of Probiotic Efficacy: A Systematic Review and Meta-Analysis. 2018. Front Med (Lausanne). Vol. 5: 124.
15. Yang B., Lu P., Li M.X., Cai X.L., Xiong W.Y., Hou H.J., Ha X.Q. A meta-analysis of the effects of probiotics and synbiotics in children with acute diarrhea. Medicine (Baltimore). 2019;98(37):e16618.
16. Sheyholislami H. and Connor K. L. Are probiotics and prebiotics safe for use during pregnancy and lactation? A systematic review and meta-analysis. Nutrients. 2021.Vol. 13(7), 2382.
17. Blaabjerg S., Artzi D.M., Aabenhus R. Probiotics for the Prevention of Antibiotic-Associated Diarrhea in Outpatients-A Systematic Review and Meta-Analysis. Antibiotics (Basel). 2017;6(4):21.

THE SCIENTIFIC TALKS OF MANIFESTATION SOME TOPICAL ISSUE ASPECTS OF FEATURES OF PHARMACOLOGY OF THE ANTIARRHYTHMIC DRUGS IN GENERAL

Nodar Sulashvili[1], Luiza Gabunia[2], Margarita Beglaryan[3], Nana Gorgaslidze[4], Ketevan Ghambashidze[5], Marika Sulashvili[6]

[1]Tbilisi State Medical University-Scientific Research-Skills Center; Georgian National University SEU- Faculty of Medicine; Sulkhan-Saba Orbeliani University - Faculty of Medicine; Alte University International - School of Medicine; Shota Meskhia Zugdidi State University - Pharmacy Program; David Aghmashenebeli University of Georgia - School of Medicine; University of Georgia- School of Health Sciences, Biochemistry and Pharmacology Direction; East European University - Faculty of Medicine; Tbilisi Humanitarian Teaching University - Faculty of Dentistry and Pharmacy; Tbilisi, Georgia; Yerevan State Medical University after Mkhitar Heratsi - Department of Pharmaceutical Management, Yerevan, Armenia.
[2]Tbilisi State Medical University - Department of Medical Pharmacology, Tbilisi State Medical University- Scientific Research-Skills Center, Tbilisi, Georgia.
[3]Yerevan State Medical University After Mkhitar Heratsi, Yerevan, Armenia. Department of Pharmaceutical Management.
[4]Tbilisi State Medical University, Department of Social and Clinical Pharmacy, Tbilisi, Georgia.
[5]Tbilisi State Medical University, Tbilisi, Georgia. Pathophysiology Department; Scientific Research Skills Center.
[6]Tbilisi State Medical University, Tbilisi, Georgia. Department of Molecular and Medical Genetics.
n.sulashvili@ug.edu.ge

Abstract. The aim of the research was to study and analyze characteristics and topical issue aspects of some features of pharmacology of the antiarrhythmic drugs in general. Antiarrhythmic drugs play a crucial role in the management of various cardiac arrhythmias, which can significantly impact patient health and quality of life. This article provides an in-depth overview of the clinical pharmacology of antiarrhythmic drugs, including their mechanisms of action, therapeutic uses, pharmacokinetics, and potential side effects. Understanding these aspects is essential for optimizing treatment and managing arrhythmias effectively. Arrhythmias, characterized by abnormal heart rhythms, are a significant concern in cardiology. Antiarrhythmic drugs are used to restore normal heart rhythm and prevent arrhythmias. This article reviews the clinical pharmacology of antiarrhythmic drugs, focusing on their classification, mechanisms of action, clinical uses, and safety considerations. Antiarrhythmic drugs are vital for the management of various arrhythmias. Understanding their mechanisms of action, clinical applications, pharmacokinetics, and potential side effects is essential for optimizing treatment and ensuring patient safety. Continued research and clinical experience will help refine the use of these drugs and improve outcomes for patients with arrhythmias. An ion channel is a part of the membrane of a nerve cell, which carries ions, that is, positively and negatively charged. The particle provides selective permeability and participates in the regulation of the body's vital processes. It is responsible for regulating the flow of ions in the plasma and membrane of all cells, controlling the intracellular concentration of various ions. The alternating closing and

opening of sodium, potassium and calcium ion channels ensures the propagation of the nerve impulse. Blockage of ion channels causes serious changes in the body. In today's medicine, a number of diseases are explained precisely by the malfunction of ion channels.

Keywords: *Clinical pharmacology, antiarrhythmic drugs, therapeutic uses, proarrhythmic.*

Introduction. Antiarrhythmic drugs are essential in the treatment of various cardiac arrhythmias, conditions characterized by abnormal heart rhythms that can lead to significant morbidity and mortality. This review explores the clinical pharmacology of these drugs, focusing on their mechanisms of action, therapeutic applications, pharmacokinetics, and potential adverse effects. By understanding these aspects, clinicians can better manage arrhythmias and optimize patient outcomes [1-3].

In the human cell membrane, there are sodium, potassium, calcium, etc. Ion channels, where Na+, K+, Ca2+ ions move respectively, which participate in the regulation of the body's vital processes. Ion channels ensure the excitability of the nervous system, the transmission of nerve impulses from the nerve to the muscle and the secretion of hormones, conditions our thinking, the work of the heart muscle and respiratory diaphragm, and more. That is why the blocking of the mentioned channels causes serious changes in the body. For example, the development of chronic fatigue syndrome is associated with dysfunction of the entire group of ion channels, including sodium and potassium channels [4-6].

Epilepsy is among the diseases caused by dysfunction of ion channels, which is caused by dysfunction of high-conductance potassium channels. The widespread cardiovascular disease long QT syndrome is associated with mutations in genes encoding potassium channels in the heart muscle, which increase the activity of potassium channels and change its normal flow in the heart muscle. Dysfunction of calcium channels leads to a state of ataxia, when coordination of movement is impossible. Therefore, the normal functioning of ion channels is very important for human health [7-8].

The complex intracellular signaling pathway in the body is preceded by the binding of a signaling molecule or ligand to its receptor. There are many different receptors and ligands, but they all have one thing in common, they are precisely paired. A receptor recognizes only one specific ligand, and a ligand binds to only one "target" receptor. Binding of a ligand to a receptor is followed by a change in the shape or activity of the receptor, resulting in signal transduction or a direct change in the cell. Ion channels are one of the "targets" for therapeutic agents [9-10].

Sodium channels There are widely present potential-dependent, potential-independent (ligand-dependent, passive, etc.) sodium channels in body cells. Sodium channels are localized in the cytoplasmic membranes of the axon hillock of neurons, dendrites and axons, in the membrane of the perisynaptic region of the neuromuscular synapse, in the sarcolemma of striated muscle fibers and in the myocardium. The distribution density of sodium channels in these structures is different. In medical prac-

tice, the so-called substances are widely used for the blockade of fast-current sodium channels. Anesthetics (Novocaine, Dicaine, Lidocaine, Sovcaine, Procaine, etc.). Anesthesia is achieved by blocking sodium channels, eliminating the ability to generate nerve impulses in afferent nerve fibers, which ultimately blocks the transmission of signals from sensory pain receptors to the central nervous system. potassium channels. There are voltage-gated and voltage-gated potassium channels. Passive, ligand-dependent and other types of channels are distinguished among the latter. Potassium channels are found in the membranes of the same cells and tissues that contain sodium channels. Na+ and K+ ions are the most important cations, the distribution and movement of which determines the appearance and change of the electrical potential. More than three dozen potassium channels have been described. calcium channels. The family of calcium channels is widely represented in the cells of nervous and muscle tissues. The main places of their localization are membranes of presynaptic terminals of sarcoplasmic and endoplasmic reticulum of muscles, sarcolemma of cardiomyocytes and cell membranes of other tissues [11-13].

Antiarrhythmic drugs are vital for managing various cardiac arrhythmias that affect the heart's rhythm and can lead to severe complications. This comprehensive review explores the clinical pharmacology of antiarrhythmic drugs, detailing their mechanisms of action, classification, therapeutic uses, pharmacokinetics, and potential adverse effects. By understanding these aspects, healthcare professionals can optimize treatment strategies and improve patient outcomes [14-16].

Antiarrhythmic drugs play a crucial role in managing various types of cardiac arrhythmias. However, their therapeutic benefits are often accompanied by a risk of toxicity, which can lead to adverse effects and complications. Understanding the toxicological profiles of these medications is essential for optimizing patient care and minimizing risks [17-19].

Aim. The aim of the research was to study and analyze characteristics and topical issue aspects of some features of pharmacology of the antiarrhythmic drugs in general.

Methods. The material of the article was the data from scientific publications, which were processed, analyzed, overviewed and reviewed by generalization and systematization. research studies are based on a review/overview assessment of the development of critical visibility and overlook of the modern scientific literature. use the following databases (for extensive literature searches to identify the characteristics and topical issue aspects of some features of pharmacology of the antiarrhythmic drugs in general.): PubMed, web of science, clinical key, Tomson Reuters, Google scholar, Cochrane library, and Elsevier foundations. national and international policies and guidelines were also reviewed.

Results and discussion. Antiarrhythmic drugs are a cornerstone in the treatment of cardiac arrhythmias. Understanding their classification, mechanisms of action, pharmacokinetics, and adverse effects is essential for optimizing patient care and improving outcomes. Continued research and advancements in drug development and monitoring technologies will further enhance the management of arrhythmias and

contribute to better patient outcomes. Cardiac arrhythmias are disorders of the heart's rhythm that can cause significant morbidity and mortality. Antiarrhythmic drugs are central to the management of these conditions, helping to restore and maintain normal heart rhythms [20-21]. Antiarrhythmic drugs are essential in managing cardiac arrhythmias but carry a risk of toxicity that can lead to significant complications. Understanding the toxicological profiles of these drugs and implementing appropriate management strategies are crucial for optimizing patient care and minimizing risks. Regular monitoring and dose adjustments based on individual patient needs and responses can help mitigate toxicity and ensure effective treatment [22-23].

Atrial fibrillation (AF), the most common cardiac arrhythmia worldwide, is driven by complex mechanisms that differ between subgroups of patients. This complexity is apparent from the different forms in which AF presents itself (post-operative, paroxysmal and persistent), each with heterogeneous patterns and variable progression. Our current understanding of the mechanisms responsible for initiation, maintenance and progression of the different forms of AF has increased significantly in recent years. Nevertheless, antiarrhythmic drugs for the management of AF have not been developed based on the underlying arrhythmia mechanisms and none of the currently used drugs were specifically developed to target AF. With the increased knowledge on the mechanisms underlying different forms of AF, new opportunities for developing more effective and safer AF therapies are emerging. In this review, we provide an overview of potential novel antiarrhythmic approaches based on the underlying mechanisms of AF, focusing both on the development of novel antiarrhythmic agents and on the possibility of repurposing already marketed drugs. In addition, we discuss the opportunity of targeting some of the key players involved in the underlying AF mechanisms, such as ryanodine receptor type-2 (RyR2) channels and atrial-selective K^+-currents (I_{K2P} and I_{SK}) for antiarrhythmic therapy. In addition, we highlight the opportunities for targeting components of inflammatory signaling (e.g., the NLRP3-inflammasome) and upstream mechanisms targeting fibroblast function to prevent structural remodeling and progression of AF. Were critically appraise emerging antiarrhythmic drug principles and future directions for antiarrhythmic drug development, as well as their potential for improving AF management [24-26].

The initiation of an AF episode requires both a trigger and a substrate. Focal spontaneous firing, also known as ectopic activity, represents the major AF-triggering mechanism. Mechanisms leading to ectopic activity include enhanced automaticity, early afterdepolarizations (EADs) and delayed afterdepolarizations (DADs). In case of enhanced automaticity, the depolarizing current during diastole is large enough to depolarize the resting membrane potential across the threshold for action potential (AP) generation, causing ectopic firing. Transient spontaneous depolarizations can also occur following complete AP repolarization, known as DADs. DADs may cause atrial ectopy if the depolarization is large enough to reach the threshold for Na+-channel activation and AP generation. The depolarizing current underlying DADs is created by increased Na^+ influx through the Na^+/Ca^{2+}-exchanger (NCX), which is activated in response to spontaneous Ca^{2+}-release events (SCaEs) from the intracellular

stores of the sarcoplasmic reticulum (SR) through the ryanodine receptor type-2 (RyR2) channels. Finally, if the AP duration (APD) is excessively prolonged, l-type Ca^{2+} channels can be reactivated, promoting a depolarizing current during the plateau phase or early in the repolarizing phase of the AP, creating EADs [27-29].

The evolution of the arrhythmogenic substrate due to atrial remodeling is promoted by cardiac and non-cardiac diseases, risk factors such as obesity and aging, and AF itself. Atrial electrical remodeling includes Ca^{2+}-handling abnormalities and changes in ion current properties that promote re-entry by abbreviating APD and effective refractory period (ERP). Atrial structural remodeling results from atrial fibrosis, atrial dilation and connexin changes that induce re-entry-promoting heterogeneous conduction slowing [30-31].

Depending on the type of AF, the key mechanisms for initiation and maintenance of the arrhythmia differ and the type and relative contribution of triggers as well as substrates change. The proper consideration of these different mechanisms is important for the discovery of specific targets and to enable tailored therapy for each type of AF [32-33].

Atrial cardiomyocytes from patients with pAF show DAD-mediated cellular triggered activity without clear evidence for electrical remodeling such as shortening of APD or hyperpolarization of the resting membrane potential. The increased incidence of DADs in pAF patients is likely partly caused by an increased Ca^{2+} uptake into the SR by the SR Ca^{2+}-ATPase type-2a (SERCA2a) leading to enhanced SR Ca^{2+} load, along with RyR2 dysregulation [40]. The open probability of RyR2 is higher in pAF, but in contrast to patients with cAF this is associated with an increase in RyR2 expression but not channel hyperphosphorylation [34-35].

There is also evidence for the potential contribution of atrial fibrosis in pAF patients. Compared to patients in sinus rhythm, pAF patients displayed atrial fibrosis in the area of crista terminalis and the inferior pulmonary veins (PV). Similar trends were observed in the Bachman's Bundle and left atrium, albeit no significant differences were found in these regions [36-37]. Ectopic activity arising from the PVs plays a particularly important role in pAF patients, with several potential proarrhythmic electrophysiological characteristics compared to the rest of the atria identified in animal models. However, the molecular mechanisms promoting ectopic activity in the PVs of humans remain largely unknown due to the very limited access to tissue samples. Animal models may provide valuable insight into the molecular mechanisms of pAF, but relevant animal models with spontaneous development of paroxysmal AF are scarce and the PVs do not appear to play a major role for the induction and maintenance of AF in the majority of animal models. Horses develop spontaneous AF, which can easily be detected by implantable loop recorders. Spontaneous PV firing has been recorded in horses, suggesting similar pAF mechanisms in horses and humans. However, the cellular and molecular mechanisms of ectopy have not been assessed in horses so far. Nevertheless, horses could prove to be a valuable animal for studying the true nature of pAF [38-39].

Overall, the current evidence suggests that the leading arrhythmogenic mechanism in patients with pAF is DAD-mediated ectopic activity caused by Ca^{2+}-handling abnormalities without clear evidence of atrial electrical remodeling. Although atrial fibrosis could contribute, the potential role of structural remodeling in pAF patients requires further investigation and validation. The cellular and molecular determinants of re-entry and AF maintenance are best characterized in patients with persistent AF and in animal models with artificially maintained tachycardia to simulate the arrhythmia and to study the specific effects of the high atrial rate [40-41].

Electrical remodeling in cAF or due to atrial tachycardia causes rapid ERP/APD shortening, primarily by increasing K+ currents and reducing l-type Ca^{2+} current (ICa, L). Although the agonist-dependent acetylcholine-activated inward-rectifier K^+ current (IK, ACh) is decreased in cAF, it develops constitutive activity and, together with an increased basal inward-rectifier K+ current (IK1) and slow delayed-rectifier K^+ current (IKs), shortens APD and hyperpolarizes resting membrane potential, promoting re-entry. The increase in IK1 is partly caused by increased protein expression of Kir2.1 and enhanced single-channel open probability promoted by phosphatase-dependent channel dephosphorylation. The constitutive activity of IK, ACh in patients with cAF is due to increased phosphorylation by protein kinase C epsilon. The functional expression of transient outward K^+-current (Ito) is lower in cAF patients, whereas IKr is unchanged [42-43].

Some studies also reported a reduced ultra-rapid delayed-rectifier K^+ current (IKur) in cAF patients, while other studies did not find current alterations. Similarly, some studies revealed increased small-conductance Ca^{2+}-activated K^+ (SK) currents (ISK) in cAF, while other work showed a decreased ISK. Since the expression of SK channels and IKur is larger in atrial compared to ventricular tissue, and under physiological conditions SK channels play a minor role in the ventricles, these channels are considered atrial predominant anti-AF targets. However, ventricular SK-channel expression is increased in heart failure and SK-channel inhibition can promote ventricular arrhythmogenesis under specific conditions. The atrial-specific TASK-1 channel, a member of the two-pore domain K+ channel family, is also upregulated in cAF, contributing to APD shortening, but is downregulated in AF patients with concomitant left-ventricular dysfunction. Thus, the degree of TASK-1 remodeling in AF patients is difficult to predict. Together, the alterations in these K^+ currents likely contribute to the proarrhythmic shortening of the atrial ERP/APD in cAF patients [44-45].

The high atrial rates during AF promote a potentially toxic cellular $Ca2^+$ overload. Consequently, ICa, L is reduced due to a calcineurin-dependent mechanism, contributing to re-entry-promoting APD shortening. Increased nuclear Ca^{2+} releases via inositol 1,4,5-triphosphate with subsequent Ca^{2+}-calmodulin protein kinase II (CaMKII) activation accelerates the nuclear export of histone deacetylase type-4 also contributing to the downregulation of ICa, L. ICa, L downregulation in turn reduces the Ca^{2+}-transient amplitude and atrial contractility. Ca^{2+}-handling abnormalities in cAF are primarily due to increased CaMKII activity, leading to RyR2 hyperphosphorylation at Ser2814 and leaky RyR2 channels causing potentially proarrhythmic

SCaEs. Protein kinase A (PKA)-mediated RyR2 hyperpolarization at Ser2808 has also been found in cAF patients, which promotes dissociation of the stabilizing calstabin2 (FKBP12.6) subunit. SR Ca^{2+} load is unchanged in cAF patients despite the larger SR Ca^{2+} leak. This can either be due to enhanced phosphorylation of phospholamban or a downregulation of the SERCA2a-inhibitory sarcolipin. Increased PKA-mediated activation of the PP1-inhibitory protein inhibitor-1, which inhibits PP1 in the SR microdomain, could explain local phospholamban hyperphosphorylation [46-48].

Structural remodeling, particularly atrial fibrosis, makes important contributions to cAF and is largely irreversible. Besides being caused by the underlying heart disease, atrial fibrosis could result from the high atrial rate during AF. Together with connexin remodeling, collagen deposition due to atrial fibrosis creates re-entry-promoting conduction heterogeneities in cAF. Of note, myofibroblasts may also couple directly to cardiomyocytes through gap junctions, which may promote DAD-mediated triggered activity in cardiomyocytes [49-50].

Conceptually, the development of POAF requires a vulnerable preoperative substrate interacting with surgery-induced triggers such as activation of the autonomic nervous system and inflammation. The pre-existing substrate for POAF includes increased pro-inflammatory signaling, with the increased expression of activated components of the NLRP3 (NACHT, LRR and PYD domains-containing protein inflammasome being the most prominent. The NLRP3 inflammasome is a multiprotein complex responsible for the cleavage and release of the pro-inflammatory cytokines IL-1β and IL-18. Increased expression of NLRP3 inflammasome components was also identified in atrial cardiomyocytes from pAF and cAF patients, pointing to a common inflammatory signature in AF. In agreement, knock-in mice with cardiomyocyte-specific NLRP3 activation revealed an ERP abbreviation, likely due to an increase in IKur, as well as atrial fibrosis and hypertrophy, both contributing to the increased AF susceptibility. In addition, a higher frequency of SCaEs, likely due to CaMKII-dependent RyR2-hyperphosphorylation, along with ICa, L alternans were also noted in POAF patients Interestingly, DADs were absent under resting conditions and only became evident upon acute application of IL-1β, mimicking one of the major post-operative inflammatory triggers. Acute IL-1β application further promoted the CaMKII-dependent RyR2 hyperphosphorylation, thereby exacerbating the pre-existing Ca^{2+}-handling abnormalities. However, the exact consequences of enhanced NLRP3-inflammasome activity in other types of AF, as well as in patients with obesity and diabetes, who are at an increased risk of developing AF, remain incompletely understood [51-53].

Similar to patients with pAF, patients with POAF do not display changes in APD and have no major changes in K+ currents. Thus, the leading mechanism in POAF appears to be ectopic (triggered) activity promoted by the acute inflammatory response without evidence of re-entry promoting APD and ERP changes, although structural changes such as connexin remodeling and fibrosis might also contribute, particularly in the left atrium [54-55].

The current ESC guidelines on management of AF focus predominantly on appropriate anticoagulation, followed by symptom control with either rate or rhythm control and management of risk factors and comorbidities. Initially, large clinical trials comparing rate and rhythm control such as Atrial Fibrillation Follow-up Investigation of Rhythm Management (AFFIRM) and Rate Control vs. Electrical Cardioversion for Persistent Atrial Fibrillation (RACE) did not find significant differences in outcome between rate and rhythm control. However, early modern rhythm control of AF might reduce the incidence of adverse cardiovascular outcomes, as shown in the Early Treatment of Atrial Fibrillation for Stroke Prevention Trial. This has again highlighted the need for improved pharmacological options for rhythm control. The challenge with traditional AADs is not only their modest efficacy, but rather their safety at the ventricular level, with ischemia and structural heart disease being a contraindication for most currently available AADs due to the high risk of drug-induced proarrhythmic. Currently used drugs are not based on knowledge of the underlying arrhythmia mechanisms and none of these drugs have been developed to specifically target AF. With the growing understanding of the fundamental mechanisms of AF it will be likely possible to develop new and improved antiarrhythmic treatment modalities based on precise knowledge of the underlying mechanisms, including upstream therapy preventing electrical and structural remodeling [56-57].

Cellular triggered (ectopic) activity is present in all forms of AF, although the silencing of atrial Ca2+-handling may also occur in cAF under specific conditions. In vivo, atrial ectopy may play a more important role in pAF and POAF than for patients with cAF. As described in Section 2, RyR2 channels play a major role in the Ca2+-handling abnormalities typically seen in AF patients. One possibility for reducing ectopic activity is, therefore, targeting abnormal RyR2 channels either by using direct RyR2-channel blockers or RyR2-function modulators. Besides being Na+-channel blockers, the Class Ic AADs such as flecainide and propafenone also inhibit RyR2 channels, reducing the open probability of RyR2 in vitro via an open-state blocking mechanism (i.e., shortened duration, but increased frequency of RyR2 channel openings, thereby increasing Ca^{2+} spark mass, while reducing the occurrence of spontaneous cell-wide Ca^{2+} waves), potentially preventing the formation of proarrhythmic Ca^{2+}-dependent DADs. The antiarrhythmic actions of flecainide on RyR2 channels have also been demonstrated in a cardiac calsequestrin knockout catecholaminergic polymorphic ventricular tachycardia (CPVT) mouse model. Moreover, a flecainide analogue that lacks RyR2-blocking properties was unable to prevent CPVT, clearly indicating that the primary mode of action of flecainide in CPVT is inhibition of RyR2. In a small clinical trial including CPVT patients, flecainide, in combination with a β-blocker, was very effective in preventing exercise-induced ventricular ectopy. In AF patients, novel routes of flecainide administration might increase cardioversion efficacy and compliance. In preclinical studies in pigs inhalative delivery of flecainide gives rise to a similar pharmacokinetic profile to flecainide given intravenously and causes a rapid cardioversion of AF. The clinical potential of in-

halative flecainide was recently tested in a clinical trial (Inhalation of Flecainide to Convert Recent Onset SympTomatic Atrial Fibrillation to sinus rhythm [58-60].

Ranolazine has been shown to reduce RyR2 open probability and to desensitize Ca^{2+}-dependent RyR2 activation, thereby preventing Ca^{2+} overload and EADs. Ranolazine is used for treatment of angina pectoris and ischemic heart diseases and possess clear antiarrhythmic properties. Unfortunately, ranolazine also carries an increased risk of proarrhythmia at the ventricular level by inhibiting IKr and prolonging APD and QT interval [61-62].

Carvedilol is a third-generation non-selective β-blocker with α1-receptor blocking properties which is used as antihypertensive drug, for the treatment of heart failure with reduced ejection fraction, and in patients with ventricular dysfunction following myocardial infarction. Carvedilol and its analogues (VK-II-86, VK-II-36, CS-I-34, CS-I-59) also exhibit RyR2-blocking properties, reducing the incidence of both DADs and EADs. The effect of carvedilol may be partly due to its β-blocking and antioxidant actions, which may reduce RyR2 phosphorylation or oxidation. Interestingly, the carvedilol analogues have no or minimal β-blocking properties, but still inhibit DAD-promoting SCaEs due to a direct reduction in RyR2 open probability. Clinically used carvedilol consists of a mixture of S-and R-carvedilol enantiomers. In contrast to the S-enantiomer, R-carvedilol does not possess β-blocking properties, but specifically blocks RyR2 channels, preventing arrhythmias ex vivo and in vivo in mice. Although the preclinical studies have shown positive signals, there are no clinical data addressing the antiarrhythmic effects of carvedilol analogues. Besides direct RyR2-channel blockers, several RyR2 modulators are available or are under development. Dantrolene for instance is a RyR1 inhibitor and is clinically used to treat potentially life-threatening malignant hyperthermia. Dantrolene has also been shown to stabilize cardiac RyR2 channels by improving the inter-domain interaction between the channel N-terminal and central domains and by enhancing the binding between calmodulin and RyR2, all of which exert antiarrhythmic effects. In a sheep model of left atrial myocardial infarction dantrolene reduced the occurrence of spontaneous AF episodes. The effects of dantrolene have also been studied in isolated cardiomyocytes from patients with cAF, showing that dantrolene was able to reduce SR Ca^{2+} leak and suppress cellular DADs. Interestingly, when a CaMKII inhibitor (autocamtide-2-related inhibitory peptide, AIP) was applied in addition to dantrolene, no additive effects on the SR Ca^{2+} leak were observed. This either indicates that the effect on RyR2 is so strong that no additional benefit can be obtained by targeting CaMKII or that dantrolene has secondary direct or indirect effects on CaMKII activity. Dantrolene also reduced AF incidence in CaMKIIδC overexpressing mice. In isolated cardiomyocytes from these mice dantrolene reduced SR Ca^{2+} leak and the incidence of both DADs and EADs. These results call for additional studies investigating whether repurposing dantrolene could be an option for effectively treating AF. Other RyR2 modulators are drugs that are capable of increasing the binding between RyR2 and its regulatory subunit calstabin2 (FKBP12.6), which acts like an endogenous RyR2-channel inhibitor by stabilizing the closed state of the channel. S107 inhibited the dis-

sociation of calstabin2 from RyR2, reduced SR Ca^{2+} leak in mouse atrial cardiomyocytes [63-64].

Drugs targeting CaMKII are of potential interest because they target multiple effector proteins including RyR2. There are drugs in preclinical development with promising efficacy for atrial arrhythmias such rimacalib (SMP-114), RA608, hesperadine which are available for oral administration. RA608 is an ATP-competitive CaMKII inhibitor selectivity targeting the predominant cardiac forms, namely CaMKIIδ and CaMKIIγ. The ATP-competitiveness of RA608 is a great advantage, as it enables the drug to bind both the active and inactive forms of CaMKII. Application of RA608 to human atrial cardiomyocytes reduces CaMKII-dependent SR Ca2+ leak and prevents induction of AF in mice. Rimacalib is also an ATP-competitive CaMKII inhibitor, which is currently evaluated in a clinical trial for the treatment of rheumatoid arthritis. Rimacalib reduced Ca2+ SR leak in human atrial and ventricular cardiomyocytes. Hesperadin is an ATP-competitive selective CaMKIIδ-inhibitor that was recently shown to improve ischemia/reperfusion induced cardiac damage [15, 18, 26].

Class III AADs such as amiodarone, dofetilide and sotalol are K^+ channel blockers that prolong ERP, thereby destabilizing re-entry. However, many Class III AADs (e.g., vernakalant, amiodarone, and its derivates dronedarone, celicarone, budiodarone and SAR11464A) are multi-channel blockers with effects from all AAD classes. As mentioned above, the safety profiles of these AADs at the ventricular level is a major limitation hindering the development of novel AADs. The challenge with targeting the typical ERP/APD changes in cAF patients is to design drugs that prolong atrial ERP without clinically relevant pro-arrhythmic effects in the ventricles, especially in patients with ischemia and structural heart diseases, which are very common in AF patients [7, 24, 46].

The antiarrhythmic properties of SK channels have been investigated in several animal models utilizing a number of different compounds. However, the expression of the three SK-channel subtypes differs greatly between species. Moreover, SK channels undergo complex remodeling in response to a wide range of pathophysiological stimuli and the reduction in SK current during cAF observed in some studies may limit the efficacy of SK-channel blockers in certain subpopulations of cAF patients. Thus, careful evaluation of SK-channel inhibitors in well-defined patient populations will be critical. The currently leading SK-channel blocker candidate, AP30663, has proven to be safe for use in humans in phase-1 studies and has entered phase-2 clinical trials [37, 39, 58].

One of the recently added atrial selective targets for potential rhythm control therapy is TASK-1. Two already well-known anti-arrhythmic drugs, namely, vernakalant and amiodarone, both inhibit TASK-1 channels. Selective TASK-1 inhibitors have also been developed and have been used in preclinical investigations. Recent in vivo and ex vivo investigations in pigs, cellular recordings from isolated human atrial cardiomyocytes, and computational modeling suggests that in the atria, the respiratory stimulant doxapram acts as a selective TASK-1 blocker with strong antiarrhythmic properties. Doxapram is currently being investigated in the DOxapram

Conversion to Sinus rhythm (DOCTOS) trial for cardioversion of both paroxysmal and persistent non-valvular AF.

The development of novel AADs faces many challenges and there are many obstacles for the translation from basic discoveries to clinical applications. Besides the high regulatory hurdles and the industry concerns, the strategies used in contemporary AAD development also need careful consideration. First, the complexity of the clinical condition in which AF develops requires a tailored approach already by first clinical exposure of a novel AAD in order to have a real chance to show a positive signal. Safety concerns also play a major role because almost all available and novel AADs possess a proarrhythmic potential. Therefore, atrial selectivity combined with minimal non-cardiac toxicity is often considered the most promising approach for the development of novel AADs for AF. However, many of the putative anti-AF targets presented above are not exclusively present in the atria. For example, although inhibition of CaMKII could be a promising anti-AF approach, CaMKII is expressed in many tissues and its inhibition could give rise to serious extra-cardiac side effects. Thus, successful drug design and appropriate formulation and application need to be carefully considered in early stage of drug development to ensure high atrial selectivity and minimal extra-cardiac effects, particularly in the brain. Atrial-specific gene therapy would be the ultimate option to ensure appropriate selectivity and besides concerns and issues that need consideration in the clinical setting an atrial-specific adeno-associated serotype 9 (AAV9) vectors has already been successfully developed and used in mice to generate efficient and atrial-specific gene expression following a single dose of systemic delivery. The development of RNA-based therapeutics is also quickly emerging [41, 49, 62].

The most promising atrial-selective drugs are those targeting TASK-1 and SK channels. TASK-1 channel inhibitors have shown promising results in preclinical studies and the first clinical trial using this approach is ongoing. Although SK channels are highly expressed in the central nervous system, the small molecules currently under development do not seem to pass the blood–brain barrier and do not exert serious neurological adverse effects in animal models or in the Phase I clinical trial. However, SK channels are not exclusively expressed in the atria of humans, suggesting only moderate atrial selectivity of SK channel inhibition. Perhaps because of this a QT prolongation was noted in the Phase I study, making further clinical investigations necessary to demonstrate the efficacy and validate the safety of SK-channel blockers. Antiarrhythmic drugs will remain a cornerstone of rhythm control therapy for millions of AF patients. Understanding of the molecular mechanisms underlying different forms of AF, identifying novel opportunities for developing mechanism-based antiarrhythmic approaches including AF-specific ion-channel blockers, targeting of Ca^{2+}-handling abnormalities (CaMKII, RyR2), and modulation of upstream pathways (notably inflammatory signaling). Numerous challenges, particularly related to achieving atrial selectivity and identifying the optimal (combination of) targets for different forms of AF, still need to be overcome and well-designed clinical trials are needed to confirm safety and efficacy of these novel approaches. Nevertheless,

the rapid developments in preclinical studies on AF therapies suggest that there is still hope for novel pharmacological approaches for the most common clinically relevant arrhythmia [18, 27, 47, 64].

Conclusion. Antiarrhythmic drugs are vital for the management of various arrhythmias. Understanding their mechanisms of action, clinical applications, pharmacokinetics, and potential side effects is essential for optimizing treatment and ensuring patient safety. Continued research and clinical experience will help refine the use of these drugs and improve outcomes for patients with arrhythmias. The classification, mechanisms of action, and potential side effects must be well understood to optimize treatment and improve patient outcomes. Ongoing research and clinical experience continue to refine the use of these medications, enhancing their efficacy and safety in arrhythmia management.

References:

1. Lip G.Y., Fauchier L., Freedman S.B. et al. Atrial fibrillation. Nat. Rev. Dis. Primers. 2016; 2:16016.
2. Kirchhof P., Camm A.J., Goette A., et al. Early Rhythm-Control Therapy in Patients with Atrial Fibrillation. N. Engl. J. Med. 2020; 383:1305–1316.
3. Marrouche N.F., Brachmann J., Andresen D., et al. Catheter Ablation for Atrial Fibrillation with Heart Failure. N. Engl. J. Med. 2018; 378:417–427.
4. Heijman J., Guichard J.B., Dobrev D. et al. Translational Challenges in Atrial Fibrillation. Circ. Res. 2018; 122:752–773.
5. Heijman J., Hohnloser S.H., Camm A.J. Antiarrhythmic drugs for atrial fibrillation: Lessons from the past and opportunities for the future. Europace. 2021;23: ii14–ii22.
6. Heijman J., Algalarrondo V., Voigt N. et al. The value of basic research insights into atrial fibrillation mechanisms as a guide to therapeutic innovation: A critical analysis. Cardiovasc. Res. 2016; 109:467–479.
7. Dan G.A., Dobrev D. Antiarrhythmic drugs for atrial fibrillation: Imminent impulses are emerging. Int. J. Cardiol. Heart Vasc. 2018; 21:11–15.
8. Hermans A.N.L., Pluymaekers N., Lankveld T.A.R. et al. Clinical utility of rhythm control by electrical cardioversion to assess the association between self-reported symptoms and rhythm status in patients with persistent atrial fibrillation. Int. J. Cardiol. Heart Vasc. 2021; 36:100870.
9. Reiffel J.A. When Silence Isn't Golden: The Case of "Silent" Atrial Fibrillation. J. Innov. Card. Rhythm Manag. 2017; 8:2886–2893.
10. Melduni R.M., Schaff H.V., Bailey K.R. et al. Implications of new-onset atrial fibrillation after cardiac surgery on long-term prognosis: A community-based study. Am. Heart J. 2015; 170:659–668.
11. Dobrev D., Aguilar M., Heijman J. et al. Postoperative atrial fibrillation: Mechanisms, manifestations and management. Nat. Rev. Cardiol. 2019; 16:417–436.
12. Higa S., Lo L.W., Chen S.A. Catheter Ablation of Paroxysmal Atrial Fibrillation Originating from Non-pulmonary Vein Areas. Arrhythmia Electrophysiol. Rev. 2018; 7:273–281.
13. Heijman J., Muna A.P., Veleva T. et al. Atrial Myocyte NLRP3/CaMKII Nexus Forms a Substrate for Postoperative Atrial Fibrillation. Circ. Res. 2020; 127:1036–1055.
14. Dobrev D., Voigt N., Wehrens X.H. The ryanodine receptor channel as a molecular motif in atrial fibrillation: Pathophysiological and therapeutic implications. Cardiovasc. Res. 2011; 89:734–743.
15. Heijman J., Linz D., Schotten U. Dynamics of Atrial Fibrillation Mechanisms and Comorbidities. Annu. Rev. Physiol. 2021; 83:83–106.

16. Shih T., Ledezma K., McCauley M. et al. Impact of traditional risk factors for the outcomes of atrial fibrillation across race and ethnicity and sex groups. Int. J. Cardiol. Heart Vasc. 2020; 28:100538.
17. Linz D., Dobrev D. Sleep apnea and atrial fibrillation: Update 2020. Int. J. Cardiol. Heart Vasc. 2020; 31:100681.
18. Nattel S., Dobrev D. Electrophysiological and molecular mechanisms of paroxysmal atrial fibrillation. Nat. Rev. Cardiol. 2016; 13:575–590.
19. Mitrofanova L.B., Orshanskaya V., Ho S.Y., Platonov P.G. Histological evidence of inflammatory reaction associated with fibrosis in the atrial and ventricular walls in a case-control study of patients with history of atrial fibrillation. Europace. 2016;18: iv156–iv162.
20. Saljic A., Jespersen T., Buhl R. Anti-arrhythmic investigations in large animal models of atrial fibrillation. Br. J. Pharmacol. 2021; 179:838–858.
21. Buhl R., Hesselkilde E.M., Carstensen H. et al. Detection of atrial fibrillation with implantable loop recorders in horses. Equine Vet. J. 2021; 53:397–403.
22. Buhl R., Nissen S.D., Winther M.L.K., et al. Implantable loop recorders can detect paroxysmal atrial fibrillation in Standardbred racehorses with intermittent poor performance. Equine Vet. J. 2021; 53:955–963.
23. Linz D., Hesselkilde E., Kutieleh R. et al. Pulmonary vein firing initiating atrial fibrillation in the horse: Oversized dimensions but similar mechanisms. J. Cardiovasc. Electrophysiol. 2020; 31:1211–1212.
24. Darkow E., Nguyen T.T., Stolina M. et al. Small Conductance Ca^{2+}-Activated K^+ (SK) Channel mRNA Expression in Human Atrial and Ventricular Tissue: Comparison Between Donor, Atrial Fibrillation and Heart Failure Tissue. Front. Physiol. 2021; 12:650964.
25. Diness J.G., Bentzen B.H., Sørensen U.S. et al. Role of Calcium-activated Potassium Channels in Atrial Fibrillation Pathophysiology and Therapy. J. Cardiovasc. Pharmacol. 2015; 66:441–448.
26. Schmidt C., Wiedmann F., Voigt N. et al. Upregulation of K(2P)3.1 K^+ Current Causes Action Potential Shortening in Patients with Chronic Atrial Fibrillation. Circulation. 2015; 132:82–92.
27. Schmidt C., Wiedmann F., Zhou X.B. et al. Inverse remodelling of K2P3.1 K^+ channel expression and action potential duration in left ventricular dysfunction and atrial fibrillation: Implications for patient-specific antiarrhythmic drug therapy. Eur. Heart J. 2017; 38:1764–1774.
28. Qi X.Y., Vahdati Hassani F., Hoffmann D., Xiao J., et al. Inositol Trisphosphate Receptors and Nuclear Calcium in Atrial Fibrillation. Circ. Res. 2021; 128:619–635.
29. Molina C.E., Abu-Taha I.H., Wang Q. et al. Profibrotic, Electrical, and Calcium-Handling Remodeling of the Atria in Heart Failure Patients with and Without Atrial Fibrillation. Front. Physiol. 2018; 9:1383.
30. Nattel S. Molecular and Cellular Mechanisms of Atrial Fibrosis in Atrial Fibrillation. JACC. Clin. Electrophysiol. 2017; 3:425–435.
31. Li N., Brundel B. Inflammasomes and Proteostasis Novel Molecular Mechanisms Associated With Atrial Fibrillation. Circ. Res. 2020; 127:73–90.
32. Wang X., Chen X., Dobrev D., Li N. The crosstalk between cardiomyocyte calcium and inflammasome signaling pathways in atrial fibrillation. Pflug. Arch. Eur. J. Physiol. 2021; 473:389–405.
33. Yao C., Veleva T., Scott L., Jr., Cao S., Li L. et al. Enhanced Cardiomyocyte NLRP3 Inflammasome Signaling Promotes Atrial Fibrillation. Circulation. 2018; 138:2227–2242.
34. Scott L., Jr., Fender A.C., Saljic A. et al. NLRP3 inflammasome is a key driver of obesity-induced atrial arrhythmias. Cardiovasc. Res. 2021; 117:1746–1759.
35. Nattel S., Sager P.T., Hüser J., Heijman J., Dobrev D. Why translation from basic discoveries to clinical applications is so difficult for atrial fibrillation and possible approaches to improving it. Cardiovasc. Res. 2021; 117:1616–1631.

36. Carstensen H., Kjær L., Haugaard M.M. et al. Antiarrhythmic Effects of Combining Dofetilide and Ranolazine in a Model of Acutely Induced Atrial Fibrillation in Horses. J. Cardiovasc. Pharmacol. 2018; 71:26–35.
37. Carstensen H., Hesselkilde E.Z., Haugaard M.M. et al. Effects of dofetilide and ranolazine on atrial fibrillatory rate in a horse model of acutely induced atrial fibrillation. J. Cardiovasc. Electrophysiol. 2019; 30:596–606.
38. Zhang J., Zhou Q., Smith C.D., et al. non-β-blocking R-carvedilol enantiomer suppresses Ca^{2+} waves and stress-induced ventricular tachyarrhythmia without lowering heart rate or blood pressure. Biochem. J. 2015; 470:233–242.
39. Avula U.M.R., Hernandez J.J., Yamazaki M. et al. Atrial Infarction-Induced Spontaneous Focal Discharges and Atrial Fibrillation in Sheep: Role of Dantrolene-Sensitive Aberrant Ryanodine Receptor Calcium Release. Circul. Arrhythmia Electrophysiol. 2018;11: e005659.
40. Pabel S., Mustroph J., Stehle T., et al. Dantrolene reduces CaMKIIδC-mediated atrial arrhythmias. Europace. 2020; 22:1111–1118.
41. Hartmann N., Pabel S., Herting J., et al. Antiarrhythmic effects of dantrolene in human diseased cardiomyocytes. Heart Rhythm. 2017; 14:412–419.
42. Zhang J., Liang R., Wang K., et al. Novel CaMKII-δ Inhibitor Hesperadin Exerts Dual Functions to Ameliorate Cardiac Ischemia/Reperfusion Injury and Inhibit Tumor Growth. Circulation. 2022
43. Mustroph J., Drzymalski M., Baier M. et al. The oral Ca/calmodulin-dependent kinase II inhibitor RA608 improves contractile function and prevents arrhythmias in heart failure. ESC Heart Fail. 2020; 7:2871–2883.
44. Diness J.G., Kirchhoff J.E., Speerschneider T. et al. The K(Ca)2 Channel Inhibitor AP30663 Selectively Increases Atrial Refractoriness, Converts Vernakalant-Resistant Atrial Fibrillation and Prevents Its Reinduction in Conscious Pigs. Front. Pharmacol. 2020; 11:159.
45. Bentzen B.H., Bomholtz S.H., Simó-Vicens R. et al. Mechanisms of Action of the KCa2-Negative Modulator AP30663, a Novel Compound in Development for Treatment of Atrial Fibrillation in Man. Front. Pharmacol. 2020; 11:610.
46. Wiedmann F., Beyersdorf C., Zhou X.B. et al. Treatment of atrial fibrillation with doxapram: TASK-1 potassium channel inhibition as a novel pharmacological strategy. Cardiovasc. Res. 2021
47. Ferreira R.R., Abreu R.D.S., Vilar-Pereira G. et al. TGF-β inhibitor therapy decreases fibrosis and stimulates cardiac improvement in a pre-clinical study of chronic Chagas' heart disease. PLoS Negl. Trop. Dis. 2019;13: e0007602.
48. Li L.Y., Lou Q., Liu G.Z., et al.. Sacubitril/valsartan attenuates atrial electrical and structural remodelling in a rabbit model of atrial fibrillation. Eur. J. Pharmacol. 2020; 881:173120.
49. Künzel S.R., Hoffmann M., Weber S., et al.. Diminished PLK2 Induces Cardiac Fibrosis and Promotes Atrial Fibrillation. Circ. Res. 2021; 129:804–820.
50. Wu Q., Liu H., Liao J. et al. Colchicine prevents atrial fibrillation promotion by inhibiting IL-1β-induced IL-6 release and atrial fibrosis in the rat sterile pericarditis model. Biomed. Pharmacother. Biomed. Pharmacother. 2020; 129:110384.
51. De Jesus N.M., Wang L., Lai J. et al. Antiarrhythmic effects of interleukin 1 inhibition after myocardial infarction. Heart Rhythm. 2017; 14:727–736.
52. Aschar-Sobbi R., Izaddoustdar F., Korogyi A.S. et al. Increased atrial arrhythmia susceptibility induced by intense endurance exercise in mice requires TNFα Nat. Commun. 2015; 6: 6018.
53. Kryshtal D.O., Blackwell D.J., Egly C.L. et al. RYR2 Channel Inhibition Is the Principal Mechanism of Flecainide Action in CPVT. Circ. Res. 2021; 128:321–331.
54. Kannankeril P.J., Moore J.P., Cerrone M. et al. Efficacy of Flecainide in the Treatment of Catecholaminergic Polymorphic Ventricular Tachycardia: A Randomized Clinical Trial. JAMA Cardiol. 2017; 2:759–766.

55. Stocco F.G., Evaristo E., Silva A.C. et al. Comparative Pharmacokinetic and Electrocardiographic Effects of Intratracheal and Intravenous Administration of Flecainide in Anesthetized Pigs. J. Cardiovasc. Pharmacol. 2018; 72:129–135.
56. Crijns H., Elvan A., Al-Windy N. et al. Open-Label, Multicenter Study of Flecainide Acetate Oral Inhalation Solution for Acute Conversion of Recent-Onset, Symptomatic Atrial Fibrillation to Sinus Rhythm. Circul. Arrhythmia Electrophysiol. 2022;15: e010204.
57. Gal P., Klaassen E.S., Bergmann K.R. et al. First Clinical Study with AP30663—A K(Ca) 2 Channel Inhibitor in Development for Conversion of Atrial Fibrillation. Clin. Transl. Sci. 2020; 13:1336–1344.
58. Krisai P., Blum S., Schnabel R.B. et al. Canakinumab After Electrical Cardioversion in Patients with Persistent Atrial Fibrillation: A Pilot Randomized Trial. Circul. Arrhythmia Electrophysiol. 2020;13: e008197.
59. Zarpelon C.S., Netto M.C., Jorge J.C. et al. Colchicine to Reduce Atrial Fibrillation in the Postoperative Period of Myocardial Revascularization. Arq. Bras. De Cardiol. 2016; 107:4–9.
60. Tabbalat R.A., Hamad N.M., Alhaddad I.A. et al. Effect of ColchiciNe on the InciDence of Atrial Fibrillation in Open Heart Surgery Patients: END-AF Trial. Am. Heart J. 2016; 178:102–107.
61. Bessissow A., Agzarian J., Shargall Y. et al. Colchicine for Prevention of Perioperative Atrial Fibrillation in patients undergoing lung resection surgery: A pilot randomized controlled study. Eur. J. Cardiothorac. Surg. 2018; 53:945–951.
62. Yancy C.W., Jessup M., Bozkurt B. et al. 2017 ACC/AHA/HFSA Focused Update of the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. J. Card. Fail. 2017; 23:628–651.
63. Glahn K.P.E., Bendixen D., Girard T. et al. Availability of dantrolene for the management of malignant hyperthermia crises: European Malignant Hyperthermia Group guidelines. Br. J. Anaesth. 2020; 125:133–140.
64. Saljic A, Heijman J, Dobrev D. Emerging Antiarrhythmic Drugs for Atrial Fibrillation. Int J Mol Sci. 2022 Apr 7;23(8):4096.

Розділ 2

БЕЗПЕКА ФАРМАКОТЕРАПІЇ ТА СИСТЕМА ФАРМАКОНАГЛЯДУ

НЕБАЖАНІ ПОБІЧНІ РЕАКЦІЇ ЛІКАРСЬКИХ ЗАСОБІВ: ФАРМАКОНАГЛЯД В ЗАПОРІЗЬКІЙ ОБЛАСТІ. ПІДСУМКИ І ПІВРІЧЧЯ 2024 РОКУ.

Крайдашенко О. В., Косогор Ю. А., Каптур Л. М.

Запорізьке регіональне відділення ДЗ «Державний експертний центр МОЗ України», м. Запоріжжя, Україна

krayd@ukr.net

Вступ. Лікарська безпека є одним з пріоритетних напрямків сучасної медицини та фармації. Медичне застосування препаратів може супроводжуватися виникненням побічних реакцій, які обумовлені фармакологічними властивостями лікарських засобів (ЛЗ) або специфічною реакцією організму на введення медикаменту.

Мета. Аналіз випадків небажаних реакцій (НПР) ЛЗ в Запорізькому регіоні при їх медичному застосуванні за І півріччя 2024 р.

Матеріали та методи. Вивчення спонтанних повідомлень про випадки НПР ЛЗ, які надійшли до регіонального відділення ДЗ «Державний експертний центр МОЗ України» за І півріччя 2024р.

Результати. За даними МОЗ України, станом на кінець 2023 р, за час війни зруйновано 144 медичних закладів та ще понад 1000 – пошкоджено. Найбільше втрат зафіксовано на Донеччині, Харківщині, Херсонщині, Миколаївщині та Київщині. На сьогодні точна кількість ЗОЗ невідома та зменшена майже вдвічі. Також слід зазначити, що в останні місяці постійно підпадає під обстріли Харківська, Херсонська, Дніпропетровська, Запорізька, Миколаївська, Чернігівська, Сумська, Донецька та Луганська області.

Внаслідок військової агресії рф проти України чотири з п’яти районів Запорізької області знаходяться на тимчасово окупованій території. При чому 53% закладів охорони здоров’я (ЗОЗ) області розташовані на цій території і зв’язок з ними відсутній. Крім того, на підконтрольній Україні території області п’ять закладів охорони здоров’я знаходяться в зоні активних бойових дій. Сектор охорони здоров’я області є одним із найбільш постраждалих внаслідок війни: протягом 2023 року зруйновано 51 об’єкт охорони здоров’я, з яких 7 – повністю, 44 – частково.

До Запорізького регіонального відділення ДЗ «Державний експертний центр МОЗ України» за І півріччя 2024 р. надійшло 692 (той самий період 2023р. – 493) повідомлень про НПР ЛЗ.

Аналіз результатів моніторингу НПР ЛЗ у ЗОЗ Запорізької області показує, що за 6 місяців 2024 року найбільша кількість (37%) спостерігалася при призначенні антибактеріальних препаратів (АБ), що можна пояснити не лише їх фармакодинамічними властивостями, але й частотою їх застосування, в тому числі без обґрунтованих на те показань. До теперішнього часу в українських аптеках можна вільно придбати найновіші та найпотужніші АБ. Це є важливим підґрунтям для застосування їх із метою самолікування і, відповідно, до розви-

тку тяжких НПР та до подальшого стрімкого прогресування резистентності мікроорганізмів до АБ. Значна частота НПР цієї групи ЛЗ насамперед пов'язана з високою активністю лікарів Запорізьких регіональних фтизіопульмонологічних ЗОЗ у сфері фармаконагляду.

Серед НПР АБ на 1-му місці за частотою були алергічні прояви (45,1%): ангіоневротичний набряк та окремі його прояви у вигляді набряку губ, повік, вух, шкіри обличчя, слизових, відчуття садіння у горлі; а також шкірні прояви алергії: папульозна висипка, плямисто-еритематозна, кропив'янка. На 2-му місці за частотою серед НПР АБ (27,3%) були порушення з боку шлунково-кишкового тракту, що здебільшого спостерігалися при їх застосуванні всередину: нудота, блювання, діарея, анорексія, біль у животі тощо. НПР із боку ЦНС та органів чуття зареєстровані у 6,1% випадків НПР АБ (ця група ПР у переважній більшості спостерігалася при тривалому використанні АБ для лікування туберкульозу).

Серед інших груп ліків НПР розподілилися таким чином: 16,6% – засоби для лікування серцево-судинних хвороб, 10,8% – анальгетики та нестероїдні протизапальні препарати. 4,3% НПР викликали засоби для лікування хвороб дихальної системи, 1-3% - для лікування травної системи, цукрового діабету, вітамінні, нейротропні ЛЗ, місцеві анестетики, інші препарати.

Серед НПР інших груп ЛЗ значну частку становили алергічні реакції – 23,7% (із них у 63,2% випадків – шкірні прояви у вигляді висипки різного типу, свербіж). Зниження артеріального тиску (АТ) зареєстровано у 3,7% випадків.

Нейро-психічні розлади - 21,4%: шум, дзвін у вухах та зниження слуху, запаморочення, порушення сну, настрою, депресія, відчуття страху, тривоги головний біль, парестезії. Порушення з боку дихальної системи (дискомфорт у грудній клітці, задишка, кашель) виявлені у 10,1% випадків.

Значна частина НПР ЛЗ стосувалася органів шлунково-кишкового тракту (29,5%): нудота, блювання, діарея, біль в епігастрії тощо. Біль у суглобах та м'язах – 3,8%, набряки нижніх кінцівок – 2,5%.

Серед найбільш частих причин виникнення НПР слід зазначити такі: ігнорування лікарями даних лікарського та алергологічного анамнезу; ігнорування лікарями положень, викладених в інструкції для медичного застосування препаратів; недостатня ознайомленість лікарів про численні синоніми торгових найменувань генеричних препаратів, що містять однакову діючу речовину; недостатні знання лікарів про клініко-фармакологічну характеристику конкретного препарату.

Висновки. Кінцевою метою фармаконагляду є зниження захворюваності і смертності, які обумовлені ЛЗ. Досягти цього можна тільки при здійсненні комплексу заходів, найважливіші з яких: удосконалення роботи національної служби фармаконагляду, впорядкування відпуску з аптек і раціоналізація застосування АБ препаратів, зміна стереотипів в роботі лікарів з метою скорочення числа нераціональних призначень ЛП, зміна ставлення хворих до необґрунтованого прийому ліків, особливо при самолікуванні.

ЕПІДЕМІОЛОГІЯ ОТРУЄНЬ АНТИДЕПРЕСАНТАМИ

Баюрка С. В.[1], Карпушина С. А.[2]
[1]Національний фармацевтичний університет, м. Харків, Україна
Кафедра фармацевтичної хімії
serhii.baiurka@gmail.com
[2]Уманський національний університет садівництва, м. Умань, Україна
Кафедра біології
svitkrp@gmail.com

Вступ. Отруєння препаратами антидепресивної дії займають провідні позиції у світі серед усіх груп лікарських препаратів, що пов'язано зі значним поширенням захворювань на депресію, а також високим суїцидальним ризиком цієї категорії хворих. Так, за даними ВООЗ, захворювання на депресію протягом найближчих десяти років, наряду з серцево-судинними та онкологічними захворюваннями, посядуть перше серед усіх вікових категорій людей. Для суспільства сучасної України характерним є надзвичайно високий рівень психоемоційних навантажень, що призводить до суттєвого погіршення рівня психічного здоров'я населення. Для профілактики отруєнь лікарськими препаратами важливе значення має розробка методів їх аналітичної діагностики. Токсичність окремих груп антидепресантів, зокрема, кардіотоксичність, трициклічних антидепресантів, є причиною гострих та смертельних отруєнь препаратами зазначеної групи.

Мета. Узагальнити основні причини інтоксикацій, та фактори, що сприяють прояву токсичних ефектів антидепресантів різних груп.

Матеріали та методи. Дані наукових публікацій стосовно епідеміології антидепресантів різних груп.

Результати. Трициклічні антидепресанти характеризуються високою терапевтичною ефективністю, але мають серйозні побічні ефекти, насамперед, кардіотоксична дія, токсичне ураження нервової системи, судоми та кому. Іншими факторами, які сприяють поширенню отруєнь антидепресантами, є перехід захворювання на депресію в хронічну форму. Трициклічні антидепресанти характеризуються високими значеннями об'єму розподілу, що свідчить про їх інтенсивний розподіл по органах і тканинах та невисокі значення концентрації у крові. Отже, висока чутливість методики визначення є необхідною умовою для використання в аналітичній діагностиці отруєнь антидепресантами зазначеної групи.

Передозування селективними інгібіторами зворотнього захвату серотоніну, насамперед, характеризуються з нейро- та кардіотоксичністю. Аналіз 469 випадків передозувань зазначеної групи антидепресантів показав, що найбільш розповсюдженим токсичним ефектом був «серотоніновий синдром», який спостерігали в 14% випадків. Передозування циталопрамом частіше супроводжується судомами та нейротоксичністю. Дослідження несмертельних випадків передозувань есциталопрамом (середня доза 140 мг, інтервал передозувань 10-560 мг)

показало, що зазначений антидепресант проявляє серотонінову токсичність. Для флувоксаміну відмічено антихолінергічну дію. Описана в літературі інтоксикація флувоксаміном супроводжувалась загрозливим для життя симптомом – епілептичним статусом, при цьому концентрація зазначеного антидепресанта у сиворотці складала 1970 мкг/мл (при терапевтичній 160-220 мкг/мл). Моноотруєння флувоксаміном рідко призводять до гострої інтоксикації та летального отруєння, 77% випадків гострих та смертельних отруєнь, з якими асоціюється флувоксамін, спричинені сумісним надходженням разом з флувоксаміном іншого антидепресанта, бензодіазепінів, нейролептиків або етанолу. В літературі наведено випадок гострого суїцидального отруєння тразодоном зі смертельним висхідом, яке супроводжувалось аритмією та поліорганною недостатністю, концентрація зазначеного антидепресанта в крові складала 25,4 мкг/мл. Описано випадок смертельного отруєння після надходження 30 мг венлафаксину, що супроводжувались симптомами, які включали судоми, тахікардію. Аналітична діагностика отруєння показала наявність в крові 21,82 мг/л венлафаксину та 3,33 мг/л *О*-дезметилвенлафаксину (терапевтичні концентрації становлять, відповідно, 0,1-0,5 та 0,2-0,4 мг/л). В літературі наведено випадок гострого отруєння флуоксетином зі смертельним висхідом. При проведенні клінічної діагностики було відмічено симптоми судомної активності, артеріальний тиск не визначався. Токсикологічне дослідження показало, що концентрація флуоксетину в крові при цьому становила 33 мг/л, норфлуоксетину – 12 мг/л. Описано смертельне отруєння дитини, яка проходила лікування метилфенідатом, клонідином та флуоксетином. Причиною отруєння стало порушення метаболізму лікарських засобів внаслідок генетичного дефекту Р-450 CYP2D системи, що призвело до значного підвищення їх токсичності. Клінічна картина отруєння включала симптоми генералізованих судом, що переросли на епілептичний статус та призвели до зупинки серця. В іншому випадку смертельного отруєння «серотоніновий синдром» розвинувся в результаті сумісного надходження до організму моклобеміду, кломіпраміну та флуоксетину.

Комбінована фармакотерапія депресій з використанням антидепресантів різних груп, застосування антидепресантів в загальній медичній практиці для лікування супутніх психічних та депресивних розладів, а також вживання лікарських препаратів антидепресивної дії не за призначенням робить актуальним розробку методів хіміко-токсикологічного аналізу біологічних об’єктів на наявність препаратів вказаної групи.

Висновки. Трициклічні антидепресанти є найбільш небезпечними серед інших груп препаратів антидепресивної дії, вони мають у декілька разів вищу летальність при передозуваннях, ніж отруєння селективними інгібіторами зворотнього захвату серотоніну.

ФАРМАЦЕВТИЧНА ОПІКА В РАМКАХ РЕАБІЛІТАЦІЇ ПАЦІЄНТІВ ІЗ ЗЛОЯКІСНИМИ НОВОУТВОРЕННЯМИ ПІСЛЯ КУРСУ ХІМІОТЕРАПІЇ

Бут О. В., Хайтович М. В.
Національний медичний університет імені О.О.Богомольця, м. Київ, Україна
kfkfnmu@gmail.com

Вступ. Онкологічні захворювання є однією з найбільших причин смертності у світі. Згідно статистики ВООЗ, станом на 2022 рік, смертність становила близько 11,2 млн. випадків, що на 10% більше ніж у 2020 році. В Україні станом на 2022 рік на обліку перебувало близько 1,2 млн пацієнтів з підтвердженим діагнозом раку, з них 20,1% - рак органів дихання, 20,2% - рак молочної залози, 18,7% - рак прямої кишки, захворювання шкіри - 13,5%. При цьому перша стадія захворювання виявлена у 26,2 % хворих, друга – у 27,1%, третя – у 15,2% та четверта – у - 17,9%.

Хіміотерапія, хоч і є ефективним методом лікування, часто супроводжується численними побічними ефектами, що суттєво впливають на якість життя пацієнтів. Актуальність фармацевтичної опіки у реабілітації таких пацієнтів полягає у забезпеченні цілісного підходу до їх лікування.

Мета дослідження. Метою дослідження є вивчення ролі фармацевтичної опіки в процесі реабілітації пацієнтів із злоякісними новоутвореннями після хіміотерапії, а також оцінка її впливу на поліпшення клінічних результатів та якості життя.

Матеріали та методи дослідження. Аналіз наукової літератури: вивчення сучасних підходів до фармацевтичної опіки у онкології. Метод спостереження: моніторинг пацієнтів під час реабілітаційного періоду.

Результати. Фармацевтична опіка в рамках реабілітації пацієнтів після хіміотерапії продемонструвала свою ефективність у зниженні побічних ефектів. Завдяки індивідуальному підбору ліків вдалося значно зменшити нудоту та блювання, що є поширеними проблемами у пацієнтів після хіміотерапії. Крім того, підтримка фармацевтів допомогла покращити апетит, що позитивно вплинуло на загальний стан здоров'я. Пацієнти відмічали, що консультації фармацевта допомогли їм краще справлятися зі стресом та тривогою, що часто супроводжують лікування раку. Надання пацієнтам інформації про їхнє лікування та способи боротьби з побічними ефектами підвищило їхній комфорт і впевненість у лікуванні. В результаті всіх цих заходів пацієнти помітили загальне поліпшення якості життя.

Висновки. Фармацевтична опіка є невід'ємною частиною реабілітації пацієнтів із злоякісними новоутвореннями після хіміотерапії. Вона покращує клінічні результати, знижує побічні ефекти та підвищує якість життя. Інтеграція фармацевтичних послуг у лікувальний процес є важливою для успішної адаптації пацієнтів і має стати пріоритетом для медичних установ.

ФАРМАЦЕВТИЧНА ПРОФІЛАКТИКА ПРИ ЗАСТОСУВАННІ СТАТИНІВ ПРИ АРТЕРІАЛЬНІЙ ГІПЕРТЕНЗІЇ

Яцкова Г. Ю.

Львівський національний медичний університет імені Данила Галицького, м. Львів, Україна

Кафедра організації і економіки фармації, технології ліків та фармакоекономіки

gyatsk@ukr.net

Вступ. Статини широко використовуються в комплексному лікуванні артеріальної гіпертензії (АГ) як засоби корекції фактору ризику – гіперхолестеринемії, а також додаткового зниження артеріального тиску завдяки наявності протизапального, ангіопротекторного, антиоксидантного та інших ефектів. Застосування статинів при АГ визнається доцільним в Європейських рекомендаціях (Guidelines) з лікування АГ: ESC/ESH 2018, ESH 2023, ESC 2024 та Уніфікованому клінічному протоколі «Гіпертонічна хвороба (артеріальна гіпертензія)» (наказ МОЗ України № 1581 від 12.09.2024 р.).

Мета. Інформаційне забезпечення одного з видів фармацевтичної профілактики – попередження ускладнень фармакотерапії, пов'язаних із появою негативних побічних ефектів (ПЕ), протипоказами при застосуванні статинів, нераціональними взаємодіями лікарських засобів (ЛЗ), що вживаються одночасно. Акцентування на таких аспектах пацієнта, якому фармацевт відпускає статин, дозволить збільшити прихильність до лікування і як наслідок – досягнути необхідного клінічного результату.

Матеріали та методи. Як джерело інформаційного забезпечення були опрацьовані Державний формуляр ЛЗ 16-ого випуску, Державний реєстр ЛЗ України. Використано логічний, системно-аналітичний метод, контент-аналіз.

Результати. В результаті систематизації даних встановлено, що фармацевт, здійснюючи фармацевтичну профілактику у відповідності до вимог фармацевтичної опіки, при відпуску зареєстрованих в Україні статинів – симвастатину (С), аторвастатину (А), розувастатину (Р), пітавастатину (П) повинен наголосити на факторах ризику прояву ПЕ статинів: вік старше 65 років, жіноча стать, порушення функції нирок, гіпотиреоз, зловживання алкоголем. Ці фактори збільшують ймовірність серйозного ускладнення – міопатії та рабдоміолізу, особливо при застосуванні високих доз статинів. При появі у хворого м'язового спазму, болю, загальної слабкості він має негайно звернутися до лікаря. Фармацевт повинен акцентувати увагу пацієнта на захворюваннях, симптомах, які є протипоказами до призначення статинів або потребують застосування з обережністю під постійним наглядом лікаря. Це - захворювання печінки у гострій стадії, гіперчутливість, тяжке порушення функції нирок, наявність ВІЛ-інфекції, дефіцит лактази, галактоземія або синдром глюкозо-галактозної мальабсорбції (у складі лікарської форми є лактоза), вагітність, годування груддю (при плануванні вагітності необхідна ефективна контрацепція). Якщо заплановані опе-

ративні втручання, слід припинити прийом статинів за декілька днів до та після операції.

Пацієнт має проінформувати лікаря про ЛЗ, які він приймає одночасно: ітраконазол, кетоконазол, кларитроміцин, еритроміцин, циклоспорин та інші інгібітори CYP3A4, інгібітори ВІЛ-протеаз, фібрати, нікотинову кислоту в дозі 1 г і більше на добу, які збільшують ризик міопатії та рабдоміолізу, а остання – ще й гепатотоксичності. При цьому найменший вплив на концентрацію в крові спостерігається для П, Р. При комбінації з менш потужними інгібіторами CYP3A4 (аміодарон, амлодипін, верапаміл, дилтіазем) необхідне зменшення дози (при поєднанні з С). Тикагрелор збільшує ризик рабдоміолізу (для С та Р). У разі призначення А, Р потрібна корекція дози пероральних контрацептивів. Збільшується токсичність дигоксину (крім П, Р). Антациди на основі магнію, алюмінію гідроксиду зменшують концентрацію в крові статинів на 35-50 %, отже необхідна перерва у 2 години між прийомами цих ЛЗ. Рифампіцин як індуктор CYP3A4 при довготривалому курсі знижує концетрацію статинів у плазмі (особливо С, менше - А). При комбінації з кумариновими антикоагулянтами можливі кровотечі (особливо для С). Найбільша кількість нераціональних взаємодій спостерігається для С.

Фармацевт має повідомити пацієнта про найбільш часті ПЕ статинів: міалгії (для А, Р, П), гепатотоксичність (для А), гіперглікемію та збільшення частоти діабету (крім правастатину, який не зареєстрований в Україні), з боку ШКТ - диспепсію, нудоту, закреп/діарею (для П, Р), шкіри - алергічні реакції (для А) та ЦНС - безсоння, головний біль (для П, Р), інфекції сечових шляхів, назофарингіт (для Р). Дуже рідко можливе інтерстиціальне захворювання легень, симптомами якого є непродуктивний кашель, диспное, що потребує негайного звернення до лікаря. При появі запаморочення, сонливості (для П, рідше - С), затуманення зору (частіше для С) пацієнту слід утриматись від керування автотранспортом та діяльності, що потребує концентрації уваги.

Для зменшення ризику ПЕ фармацевт повинен рекомендувати пацієнту дотримуватися правильного способу застосування: приймати незалежно від прийому їжі в один і той же час (С та П увечері), таблетку ковтати цілою, запивати водою. Не рекомендоване вживання грейпфрутового соку. Протягом усього курсу лікування дотримуватися холестеринзнижуючої дієти: обмеження насичених жирів, трансжирів, моно- та дисахаридів, вуглеводів, алкоголю, збільшення вмісту рослинної клітковини, продуктів з фітостеролами - рослинних олій (оливкової та інших), насіння, горіхів, зернової продукції.

При тривалому курсі лікування необхідний регулярний контроль рівня трансаміназ (кожні 8-12 тижнів), глюкози (пацієнтам літнього віку, з метаболічним синдромом, ожирінням та при високодозовій терапії), функції нирок (при застосуванні Р у дозі 40 мг (ризик протеїнурії)).

Висновки. Запропоновані джерела та систематизовані дані, необхідні для інформаційного забезпечення фармацевтичної профілактики ускладнень фармакотерапії при застосуванні статинів.

ПРОБЛЕМА ВПЛИВУ ЕЛЕКТРОННИХ СИГАРЕТ НА ЗДОРОВ'Я ЛЮДИНИ

Козар В. В.[1], Кондратенко К. К.[2]

[1]Національний фармацевтичний університет, м. Харків, Україна, кафедра клінічної фармакології ІПКСФ

[2] Харківський національний медичний університет, м. Харків, Україна

kozarval@gmail.com

Електронні сигарети (ЕС) були представлені на світовому ринку в середині 2000-х років. ЕС була розроблена в Китаї у 2003 р. як альтернатива звичайній сигареті. ЕС вперше запатентовані були у Сполучених Штатах у 2007 році. З 2014 року ці вироби стали найпоширенішою формою тютюну. За даними Глобального опитування молоді щодо вживання тютюну (GYTS, 2017), 40,3% дітей пробували курити ЕС, а 18,4% дітей у віці 13–15 років (тобто фактично кожен п'ятий підліток) і тепер є курцями ЕС. В деяких містах та штатах США вже заборонили продаж ЕС. Слід також відзначити, що жодна електронна сигарета не була схвалена FDA як засіб для відмови від куріння.

Національна епідемія електронних сигарет серед молоді в основному пояснюється ароматизованими тютюновими виробами, включаючи ментол та інші ароматизатори. У США 27,5% учнів середньої школи використовують ЕС; 3 найбільші користувачі ринку походять із США, Великобританії та Японії. У 2021 році глобальний ринок ЕС оцінювався в 20,4 мільярда доларів США, очікується, що до 2027 року він зросте до 30 мільярдів доларів США. Викликає тривогу те, що використання ЕС серед молоді з 2011 по 2019 роки зросло на 1800%. Три з чотирьох користувачів віком від 15 до 21 років не знають, що ЕС містить нікотин.

Результати ряду проведених досліджень хімічного аналізу показали, що аерозолі електронних сигарет містять численні подразники та токсичні речовини (нікотин, специфічні для тютюну нітрозаміни, метали, поліциклічні ароматичні вуглеводи, леткі органічні сполуки, карбонільні сполуки та альдегіди, добавки, ароматизатори та розчинники, зокрема, пропіленгліколь та/або гліцерин, які при нагріванні можуть утворювати канцерогенні сполуки тощо).

Є документально підтверджені цитотоксичні ефекти компонентів електронної сигарети на легеневу тканину. У дослідженні впливу аерозолю ЕС на епітеліальні клітини бронхів встановлено зниження метаболічної активності та життєздатності клітин, а також збільшення вивільнення кількох маркерів запаленн [Ratajczak A. et al., 2018]. Повідомлялося, що надмірне вейпування викликає запальні реакції, включаючи активацію мітоген-активованої протеїнкінази, тирозинкінази Janus, активатора транскрипції та передачі сигналів ядерного фактора-κB, подібну до реакції, індукованої тютюновим димом. Згідно з останніми доказами, тривалий вплив деяких компонентів аерозолів електронних си-

гарет може призвести до респіраторних ускладнень, таких як астма, хронічне обструктивне захворювання легенів та запалення (Kaur G. et al., 2018).

В публікації (Rubinstein M.L. et al., 2018) про вплив на підлітків токсичних летких органічних хімічних речовин з електронних сигарет встановлено, що в організмі підлітків, які вживають електронні сигарети, виявлено у сечі щонайменше 5 потенційно шкідливих токсикантів. Багато летких органічних сполук, які виявили серед користувачів електронних сигарет, є канцерогенними, зокрема оксид пропілену, акриламід, акрилонітрил і кротоновий альдегід. За деякими винятками, ці токсичні речовини були присутні незалежно від того, містив продукт нікотин або ароматизатори.

Спостереження про те, що використання ЕС може бути пов'язано як із посиленням респіраторних симптомів, так і з симптомами астми, а також із збільшенням прогулів підлітків у школі, пов'язаних з астмою, потенційно викликає занепокоєння, оскільки зниження функції легенів у наступні роки пов'язують з астмою та хронічним бронхітом у дитячий та юнацький вік. В дослідженні (Alnajem A. et al., 2020) було показано, що на респіраторне здоров’я підлітків негативно впливають не лише використання електронних сигарет, а й пасивне вдихання їх аерозолів, які були незалежно пов’язані з симптомами астми серед підлітків.

Публікації довгострокових результатів дослідження демонструють шкідливий вплив електронних сигарет на здоров’я людини. Використання електронних сигарет серед підлітків у США зросло до рівня епідемії, виявивши новий феномен ушкодження легенів, пов’язаного з електронними сигаретами (EVALI). EVALI вперше було описано влітку 2019 року як сукупність конституційних, шлунково-кишкових і респіраторних симптомів з гіпоксемією та двосторонніми інфільтратами на візуалізації за відсутності інфекції (Layden J.E. et al., 2019). Діагностичні дані відрізнялисяся залежно від характеру ураження легенів. Проте були виявлені типові моделі загальних знахідок, у тому числі наявність матового помутніння на комп’ютерній томографії та лейкоцитоз. Основи лікування включали використання кортикостероїдів, антибіотиків і вентиляційну підтримку, в тому числі екстракорпоральну мембранну оксигенацію. Результати варіювалися від повного або майже повного відновлення функції легенів до смерті (Clarelle L. Gonsalves et al., 2021).

26 штатів і округ Колумбія в США підтвердили 52 смерті, пов’язані з вейпінгом. На додаток до захворювань, пов’язаних із легенями, деякі повідомлення про випадки включали неврологічні та шлунково-кишкові симптоми [My Hua. et al., 2020].

Так, Rothrock та ін. (2020) визначили, що серед підлітків віком від 10 до 19 років споживачі ЕС мають більший ризик для будь-якого вживання алкоголю порівняно з тими, хто не вживає ЕС, крім того, користувачі ЕС мали вищий ризик сп’яніння/запою порівняно з підлітками, які не користувалися ЕС. Загалом було виявлено, що вживання ЕС пов’язане з низкою проблем психічного

здоров’я підлітків, включаючи імпульсивність, тривогу та схильність до суїциду.

У 2022 році дохід сегмента електронних сигарет в Україні склав 47,5 млн доларів США. Зменшення доходу сегмента у 2022 році відбулося через зменшення населення в країні та окупацією територій (Pro-Consulting, 2023).

3 квітня 2014 року Європейський парламент затвердив нову Директиву 2014/40/ЄС щодо вимог до регулювання тютюнових виробів та електронних сигарет, яка набула чинності у всіх країнах Європейського союзу 20 травня 2016 року. Уряд України уже визнав зобов’язання щодо імплементації Директиви 2014/40/ЄС в українське законодавство. Законом України № 1978-IX «Про внесення змін до деяких законів України щодо охорони здоров’я населення від шкідливого впливу тютюну», редакція від 23.11.2023, внесено чимало змін до антитютюнового законодавства, зокрема, і обмежень щодо електронного паління, таких як вимоги до маркування, розширення заборонених місць використання, введення вікових обмежень тощо.

Таким чином, сучасні дані свідчать про те, що електронні сигарети не позбавлені серйозного ризику для здоров’я людини, а кількість осіб, які їх використовують значно зменшити міжнародній спільноті поки що не вдалося, тому не менш важливим є пошук лікарських засобів, які б змогли зменшити побічні ефекти ЕС, в тому числі на здоров’я дітей та підлітків.

КЛІНІКА УРАЖЕНЬ НЕРВОВОЇ СИСТЕМИ ПРИ ГОСТРИХ ОТРУЄННЯХ СИНТЕТИЧНИМИ КАНАБІНОЇДАМИ У ДІТЕЙ

Одинець І. Ю.[1], Міщенко В. А.[2], Пороша Н. С.[3], Лисенко Л. С.[1], Пушкар М. Б.[1]

[1]Навчально-науковий інститут післядипломної освіти Харківського національного медичного університету, кафедра анестезіології, інтенсивної терапії та дитячої анестезіології , м. Харків, Україна

[2]Національний фармацевтичний університет, Інститут підвищення кваліфікації спеціалістів фармації, кафедра клінічної фармакології , м. Харків, Україна

[3]Комунальне некомерційне підприємство Харківської Обласної Ради «Обласна інфекційна клінічна лікарня», м. Харків, Україна

Вступ. Високий рівень споживання та отруєння синтетичними канабіноїдами (СК) привертає до себе увагу як лікарів наркологів, так і лікарів інших спеціальностей. Швидке розповсюдженням цих наркотичних речовин (НР) спостерігається не тільки серед дорослого населення але і серед дітей та підлітків. Незважаючи на спроби посилити обмеження щодо СК, їхня популярність значно зросла з моменту появи. Вони мають відносно низьку вартість, є легкодоступними. Швидкість з якою на ринок потрапляють нові варіанти СК не дає мо-

жливості додати їх до реєстру наркотичних речовин ще до початку розповсюдження, тому вони не виявляються в сечі при використанні доступних скринінгових тестів на НР. Занепокоєність лікарів збільшилась у зв'язку з тим, що вживання СК дітьми та підлітками призводить до виникнення у більшості з них тяжких отруєнь, з подальшим розвитком критичних станів. Певні труднощі пов'язані з диференційною діагностикою між наркотичним отруєнням СК та схожими клінічними ознаками, що реєструються при деяких соматичних та інфекційних захворюваннях.

Матеріали та методи. До КНП ХОР «Обласна дитяча інфекційна клінічна лікарня» у відділення інтенсивної терапії, за період з 2017 по 2024 рр. було госпіталізовано 17 дітей віком від 7 до 17 років у критичному стані.

Результати. Ознаки хвороби виникали раптово. Діти були блідими, погано доступні контакту, збуджені, у них спостерігалися жахливі галюцинації, вони кричали, збудження чергувалося з пригніченням або втратою свідомості, у більшості дітей реєструвався судомний синдром. Мала місце блювота та гіпертермія 37,5-39 о С. Лихоманка, ознаки пошкодження нервової системи (НС) (збудження, панічні атаки, агресивність, запаморочення свідомості, дезорієнтація, гіпертонус м'язів, судомний синдром, слухові та зорові галюцинації), позитивні менінгеальні знаки дозволили діагностувати гострий менінгоенцефаліт. Вірусних або бактеріальних чинників, що викликають пошкодження НС, не виявлено. При МРТ головного мозку ознак вогнищевої патології та осередків запального процесу не знайдено. Вищезазначене дозволило припустити отруєння невідомою сполукою, можливо НР. Було прийняте рішення після отримання згоди батьків провести діагностику сечі на наявність НР. Тест був проведений 12 дітям та отриманий позитивний результат на СК. Батьки 5 пацієнтів відмовилися від проведення цього тесту. При спілкуванні з цими дітьми був встановлений факт прийому ними НР. Таким чином, дебют отруєння СК дуже схожий на початок гострого інфекційного захворювання, що перебігає з пошкодженням переважно НС з наступним розвитком коми, серцевої та дихальної недостатності. Виключення інфекційної патології у таких хворих потребує призначення дороговартісних вірусологичічних, бактеріологічних, інструментальних методів дослідження (УЗД, МРТ та ін.).

Висновки. Слід рекомендувати перед вищезазначеними дослідженнями проводити тест на наявність НР дітям з клінічними ознаками, схожими з інфекційним ураженням НС (гострі менінгоенцефаліти, енцефаліти), тяжким судомним синдромом, який виник вперше, або діагностично складним пацієнтам.

БЕЗПЕКА ВИКОРИСТАННЯ АНТИГІПЕРТЕНЗИВНИХ ПРЕПАРАТІВ ПЕРШОГО РЯДУ

Кайдаш С. П., Слєсарчук В. Ю.
Дніпровський державний медичний університет, м. Дніпро, Україна
Кафедра фармакології, загальної та клінічної фармації
skaidash1704@gmail.com

Вступ. Раціональна та безпечна фармакотерапія артеріальної гіпертензії (АГ) продовжує бути актуальною та водночас складною проблемою сьогодення. Відомо, що до антигіпертензивних препаратів першого ряду відносять: інгібітори ангіотензин-перетворюючого ферменту (АПФ), блокатори рецепторів ангіотензину II (БРА II), дигідропіридинові блокатори кальцієвих каналів тривалої дії (БКК), тіазидні та тіазидоподібні діуретики – для більшості пацієнтів у якості початкового лікування АГ та бета-адреноблокатори (БАБ) – за спеціальних показань. Також, надається перевага у комбінованій дво- або три компонентній терапії у більшості хворих з АГ. При тому, стратегічно важливо використання у одній таблетці фіксованих комбінацій кількох гіпотензивних препаратів, що спрощує алгоритми лікування АГ та покращує його комплаєнс.

Мета. З метою підвищення безпеки використання створюваних фіксованих комбінацій, був проведений аналіз даних літератури щодо побічних ефектів вищезгаданих груп гіпотензивних препаратів.

Методи дослідження. Бібліосематичний аналіз, огляд літератури, статистичний аналіз.

Результати дослідження. На основі проведеного аналізу даних літератури встановлено, що характерними побічними ефектами інгібіторів АПФ є висока частота алергічних реакцій у вигляді ангіоневротичного набряку (рідше можуть бути шкірні висипи, фотосенсибілізація), токсичні побічні ефекти – помірні, і загалом залежать від тропності, шляхів метаболізму та виведення інгібіторів АПФ, серед фармакодинамічних побічних ефектів часто зустрічається сухий кашель (покахикування, через накопичення брадикініну), електролітний дисбаланс – гіперкаліємія, артеріальна гіпотензія, також слід зазначити, що інгібітори АПФ суворо протипоказані вагітним. Група БРА II загалом досить безпечна, малотоксична, серед алергічних побічних ефектів іноді можливий набряк Квінке, кропив’янка, серед фармакодинамічних – помірна гіперкаліємія, ортостатична гіпотензія, також протипоказання для вагітних. Дигідропіридинові БКК низькоалергені, токсичність, зокрема, кардіотоксичність (уповільнення атріовентрикулярної провідності і зниження скоротливості міокарду) у терапевтичних дозах - не виражені, фармакодинамічні побічні ефекти частіше пов’язані з надмірною периферічною вазодилятацією і є дозозалежними (артеріальна гіпотензія, колапс, рефлекторна тахікардія, сердцебиття, периферичні набряки гомілок, почервоніння обличчя, головний біль, порушення зору), диспептичні прояви (нудота, біль в епігастрії, закрепи) теж пов’язані з вазодилятацією судин слизової ШКТ та брижейки. Алергічні побічні реакції тіазидних

та тіазидоподібних діуретиків є рідкими, але може бути перехресна алергія до інших груп діуретиків; тіазидні виявляють токсичність на обміни речовин: вуглеводний (порушення толерантності до глюкози, гіперглікемія), ліпідний (підвищення рівнів загального холестерину, ліпопротеїдів низької щільності та тригліцеридів, зниження рівня ліпопротеїдів високої щільності), а тіазидоподібні є метаболічно нейтральними і не впливають на ці обміни. Тіазидні та тіазидоподібні діуретики токсично впливають на пуріновий (гіперурікемія) та мінерально-електролітний (гіпокаліємія, гіпонатріємія, гіпохлоремія, метаболічний алкалоз, гіперкальціємія) обміни, також відмічають нефротоксичність (інтерстиціальний нефрит) та артеріальну гіпотензію. Бета-блокатори (БАБ) низькоалергені, ступінь токсичності залежить від селективності, ліпофільності, дози та тривалості прийому, зокрема, відмічають кардіотоксичність (пригнічення синоатріальної та атріовентрикулярної провідності, AV блокади, надмірна брадикардія з ЧСС < 50 уд/хв., зниження скоротливості міокарду з можливим проявом серцевої недостатності), нейротоксичність, як наслідок проникнення через гематоенцефалічний бар'єр (головний біль, запаморочення, втрата свідомості, депресія, жахливі сновиди, безсоння, розлади когнітивних функцій), гепатотоксичність, токсичний вплив на вуглеводний (раптова гіпоглікемія), ліпідний (зниження рівня ліпопротеїдів високої щільності), електролітний (гіперкаліємія, гіпонатріємія) обміни. Фармакодинамічними побічними ефектами БАБ є гіпотензія, синдром відміни (погіршення стану хворого при раптовій відміні БАБ у вигляді нападу стенокардії, підвищення артеріального тиску), а для неселективних БАБ характерний бронхоспазм та сексуальна дисфункція з розвитком імпотенції у чоловіків і зниженням лібідо у жінок.

Висновки. Комбінуючі між собою вищезгадані групи гіпотензивних препаратів, слід враховувати фармакодинамічну взаємодію, ефект сумації, потенціювання (підсилення гіпотензивного клінічного ефекту, кардіопротекторних, вазопротекторних, нефропротекторних властивостей груп препаратів), можливість відкоригувати електролітний калієвий дисбаланс, а також сумарне підвищення і побічних ефектів, таких як: артеріальна гіпотензія, у т.ч. ортостатична; кардіотоксичність (уповільнення атріовентрикулярної (AV) провідності, AV-блокади, зниження скоротливості міокарда, брадикардії); також, при призначенні метаболічно не нейтральних груп препаратів, слід враховувати супутню патологію у хворого (цукровий діабет, дисліпідемії, ожиріння, бронхіальну астму, ХОЗЛ, подагру, остеопороз).

ОСОБЛИВОСТІ ЗАСТОСУВАННЯ ЛІКІВ У ВАГІТНИХ

Команюк Л. В., Паламар А. О.
Буковинський державний медичний університет, м. Чернівці, Україна
Кафедра фармації
palamar@bsmu.edu.ua

Вступ. За даними ВООЗ, близько 90 % жінок приймають лікарські засоби під час вагітності, як за призначенням лікаря, так і самостійно, при цьому 75 % вживають більше чотирьох препаратів в I та II триместрах вагітності. Враховуючи, що більша частина ліків можуть виявляти ембріотоксичну, тератогенну та фетотоксичну дії на плід, є дуже актуальним питання раціонального та безпечного застосування ліків у вагітних. Оскільки, адекватна та своєчасна терапія відіграє ключову роль у збереженні життя та запобіганні негативним наслідкам як для матері, так і для дитини.

Мета. Провести аналіз безпеки та умов раціонального застосування ліків у період вагітності, включаючи вибір лікарських препаратів з мінімальним ризиком, оцінку потенційних загрозливих наслідків та умови безпечної фармакотерапії.

Матеріали та методи. Об'єкти дослідження були рекомендації, директиви ВООЗ, міжнародні та національні бази даних, які містять інформацію про безпечний прийом ЛЗ у вагітних жінок. Використано методи аналізу, систематизації та узагальнення інформації.

Результати. Наслідки прийому ліків вагітними залежать не лише від препарату, а й терміну вагітності, дози і тривалості прийому, віку вагітної, стану її здоров'я та генетичної схильності до розвитку тієї чи іншої вади.

Згідно з літературними даними відомо, що найчастіше вагітні у I триместрі приймають протиблювотні лікарські засоби, у II – антигістамінні та антибактеріальні, у III – седативні та діуретичні препарати. Значна частина жінок протягом усього терміну вагітності застосовують препарати заліза та вітаміни, а також часто анальгетики, спазмолітики, антигіпертензивні, гормональні та дерматологічні засоби. Доведено необхідність приймання фолієвої кислоти вагітними жінками у дозі 400-800 мкг до 12 тижнів вагітності, проте слід пам'ятати, що є низка препаратів, які порушують метаболізм гормонів та обмін фолієвої кислоти (синтетичні естрогени, гестагени, антитиреоїдні засоби, метотрексат та ін.)

При застосуванні лікарських засобів у вагітних важливо враховувати термін вагітності, особливо зважити необхідність у прийомі тих чи інших ЛЗ під час I триместру, оскільки під час цього періоду відбувається ембріогенез. Ембріотоксичну дію можуть викликати естрогени, гестагени, анаболічні стероїди, дезоксикортикостерону ацетат, соматотропний гормон, саліцилати, барбітурати, сульфаніламіди, цитостатики, ізоніазид, нікотин та ряд інших ксенобіотиків. Фетотоксична дія спостерігається при застосуванні таких препаратів, як антикоагулянти, які провокують виникнення кровотечі у новонарод-

женого, алкоголь, снодійні та транквілізатори пригнічують функції ЦНС, похідні сульфонілсечовини зумовлюють гіпоглікемію, нестероїдні протизапальні засоби у III триместрі призводять до передчасного закриття артеріальної протоки.

Перевагу слід надавати лікарським препаратам заліза із категорії А (безпечні ЛЗ) та В (умовно безпечні ЛЗ), до яких належать заліза гідроксид (III) полімальтозат, заліза гідроксид (III) сахарозний комплекс, заліза глюконат (II), заліза фумарат та заліза амонійного цитрат. Антипіретиком та анальгетиком першого вибору у вагітних є ацетамінофен (парацетамол), який відноситься до категорії В. Проходить крізь плаценту, але вважається безпечним для плода у терапевтичних дозах. При артеріальній гіпертензії використовують антигіпертензивні препарати з доведеною ефективністю щодо зниження ризику розвитку ускладнень, а саме метилдопа (категорія В). Також застосовують дилтіазем, верапаміл, ніфедипін, які віднесені до категорії С (потенційно небезпечні), бісопролол та комбінації цих препаратів.

Протипоказані інгібітори АПФ (каптоприл, еналаприл) тому, що спричиняють пошкодження нирок у плода та анурію новонародженого. Вони також гальмують нормальний гестаційний розвиток судинної системи. Блокатори ангіотензинових рецепторів (валсартан, ірбесартан, кандесартан) також не можна застосовувати під час вагітності, адже вони, так само як і інгібітори АПФ належать до категорії D (небезпечні ЛЗ).

Серед антибактеріальних препаратів найбезпечнішими у вагітних є пеніциліни, за умов відсутності алергічних реакцій. Також можуть бути застосовані цефалоспорини I-III покоління, азтреонам та окремі макроліди (еритроміцин). Протипоказане застосування таких препаратів з категорії D, як тетрацикліни, що можуть призвести до порушення прорізування зубів та росту кісток, аномалії кінцівок, катаракти, аміноглікозиди – до глухоти, левоміцетин (у III триместрі) – синдрому «сірого немовляти», а прийом нітрофуранів та сульфаніламідів викликає гемоліз та жовтяницю.

Потрібно повністю уникати застосування препаратів з категорії Х (абсолютно протипоказані): андрогени (призводять до вірилізації, укорочення кінцівок, аномалії трахеї, стравоходу, порушень серцево-судинної системи), естрогени (вроджені вади серця, фемінізація чоловічого плода), дисульфірам (спонтанні аборти, розщеплення кінцівок), йод (кретинізм, гіпотиреоз), ретиноїди (аномалії кінцівок, черепно-лицьових відділів, аномалії серця і центральної нервової системи, сечостатевої системи) та ін.

Висновки. Отже, прийом лікарських засобів під час вагітності має бути ретельно зважений і обґрунтований, оскільки він може мати різні наслідки для здоров'я матері та розвитку плода. Слід дотримуватися таких рекомендацій – обмежити прийом ліків у I триместрі вагітності, застосовувати мінімальні дози найбільш безпечного препарату, надавати перевагу монотерапії, застосовувати лікарський засіб тільки в тому випадку, якщо його користь перевищує всі потенційні ризики для плоду.

THE SCIENTIFIC TALKS OF MANIFESTATION OF TOPICAL ISSUES OF CHARACTERISTICS OF COGNITIVE-ENHANCING STIMULANT DRUGS ACCORDING THE SAFETY OF PHARMACOTHERAPY AND PHARMACOVIGILANCE SYSTEM IN GENERAL

Luiza Gabunia[1], Ketevan Ghambashidze[2], Levan Ratiani[3], Nana Gorgaslidze[4], Shorena Khetsuriani[5], Natia Gamkrelidze[6], Elena Varazi[7], Londa Rukhadze[8], Nana Peikrishvili[9], Nodar Sulashvili[10]

[1]Tbilisi State Medical University, Tbilisi, Georgia; Scientific Research-Skills Center.
[2]Tbilisi State Medical University, Tbilisi, Georgia; Pathophysiology Department.
[3]Tbilisi State Medical University, Tbilisi, Georgia; The First University Clinic, Department of Emergency Medicine, Reanimatology and Anesthesiology.
[4]Tbilisi State Medical University, Tbilisi, Georgia; Department of Social and Clinical Pharmacy.
[5]Tbilisi State Medical University, Tbilisi, Georgia; Scientific Research Skills Center, Microbiology Department.
[6]Tbilisi State Medical University, Tbilisi, Georgia; Scientific Research Skills, Pathophysiology Department.
[7]Tbilisi State Medical University, Tbilisi, Georgia; Research Skills Center.
[8]Faculty of Healthcare of the University of Porto, Portugal; Scientific Research Skills Centre at Tbilisi State Medical University; Tbilisi, Georgia.
[9]Scientific Research Skills Centre at Tbilisi State Medical University; Tbilisi, Georgia.
[10]Scientific Research-Skills Center at Tbilisi State Medical University, Faculty of Medicine at Georgian National University SEU, Faculty of Medicine at Sulkhan-Saba Orbeliani University, Division of Pharmacology of International School of Medicine at Alte University; Shota Meskhia Zugdidi State University; School of Medicine at David Aghmashenebeli University of Georgia, Associate Professor of Biochemistry and Pharmacology Direction at the University of Georgia, School of Health Sciences. Associate Professor of Pharmacology of Faculty of Medicine at East European University, Associate Professor of Pharmacology of Faculty of Dentistry and Pharmacy at Tbilisi Humanitarian Teaching University; Tbilisi, Georgia; Researcher of Department of Pharmaceutical Management of Yerevan State Medical University after Mkhitar Heratsi, Yerevan, Armenia.
n.sulashvili@ug.edu.ge

Abstract. The aim of the research was to study and analyze the key issue aspects related to features of effects of using psychostimulant substances. This article explores the current scientific discourse surrounding cognitive-enhancing stimulant drugs. It examines their characteristics, mechanisms of action, potential benefits, risks, and societal implications. The analysis aims to provide a comprehensive overview of these substances, focusing on their impact on cognitive functions and their broader implications for individuals and society. Generally, amphetamines are characterized by high abuse potential; They represent recreational drugs and are characterized mainly by central rather than peripheral effects; Frequently, an ecstasy tablet with MDMA contains antihistamines and caffeine, and in the worst case, cocaine or ketamine. MDMA enhances not only the release of serotonin, but also increases its consumption in the body; It is also worth noting that hangover is usually accompanied by bad mood or depression, because, as mentioned above, high levels of serotonin are consumed throughout consumption of MDMA and during hangover the body suffers from lack of serotonin. Serotonin is known as a transmitter of good mood and pleasant feeling, which is mainly concentrated in the central nervous system and intestines. The relationship between sleep disorders and substance abuse is only loosely understood, but shows some relation with sleep problems reinforcing substance use disorders, as well as substance use leading to sleep disturbances. In the review, characterized several sleep disturbances produced by cocaine, cannabis, and opioid short-

term and long-term abstinence, suggesting that substantial research into the effectiveness of sleep agents for addiction treatment is needed. The use of cognitive-enhancing stimulant drugs has become increasingly prevalent in both clinical and non-clinical settings. These substances, which include a variety of pharmacological agents, are designed to improve cognitive function, particularly in domains such as memory, attention, and executive function. The scientific community has been actively discussing these drugs' mechanisms, efficacy, and ethical considerations. This article aims to synthesize current scientific discussions and provide a thorough understanding of the topical issues related to these drugs.

Key words: *Features, effects, using, psychostimulant, substances pharmaceutical, pharmacodynamics.*

Introduction. The quest to enhance cognitive functions through pharmacological means has intrigued scientists and clinicians for decades. Cognitive-enhancing stimulant drugs, ranging from prescription medications to over-the-counter substances, have garnered significant attention for their potential to improve various aspects of mental performance. These drugs, often designed to target specific neurotransmitter systems, offer a promise of enhanced memory, attention, and executive function. However, the use and impact of such substances raise important questions regarding their efficacy, safety, and ethical implications [1-2]. In this article, we delve into the intricate details of cognitive-enhancing stimulant drugs, exploring their mechanisms of action, efficacy, and side effects. We also examine their applications in clinical settings and the broader societal implications of their use. By synthesizing current scientific discussions, this article aims to provide a comprehensive overview of the topical issues surrounding cognitive enhancers, offering insights into both their potential benefits and risks [3-4].

Amphetamines are substances that have a stimulating effect on the central nervous system. Although they are widely known as illegal drugs that cause physical and mental harm, they are also legal drugs with beneficial properties. The main difference lies in the composition and use of drugs, because amphetamine drugs must be taken under strict medical supervision. Although they are used as medicines and not for abuse, these substances can cause a number of side effects on the body. In the therapeutic field, amphetamine is used to synthesize drugs such as fenproprox or clobenzox, which increase brain stimulation. In contrast, in the field of drugs, amphetamine is used to synthesize narcotic substances such as methamphetamine. Generally, all amphetamine preparations (medical and recreational) are taken by mouth, with effects beginning in about 30 minutes. However, in some cases, especially with drug abuse, the drug can be taken parenterally (injection), which leads to an increase in the hallucinogenic effect of the compound. The main feature of this drug is that its distribution in the body is general, it easily crosses the blood-brain barrier and, despite the size of the molecules, it penetrates the brain very easily. Usually, these substances have multiple effects on the brain, which is why many neurotransmitters are involved in the use amphetamine. So, dopamine and serotonin, adrenaline or norepinephrine

change in their function when amphetamine enters the brain, so the effects caused by this drug are very diverse. This fact partly explains that the same substance can be used as a therapeutic measure, as well as for narcotic and recreational use, since a controlled action can benefit the brain, but too much action can cause obvious addiction and various psychological changes [5-7].

Side effects of amphetamines are the most common: nervousness, trouble sleeping, uncontrollable fear, headache, changes in sexual sensitivity, dry mouth, constipation or diarrhea, loss of appetite, weight loss, and addiction. Amphetamines are central nervous system stimulants, so taking these substances usually increases brain activity. Thus, while the use of other drugs, such as cannabis or alcohol, can suppress or slow down certain types of brain activity, amphetamine is mainly characterized by excitation and stimulation of this area of the body. This fact leads to the manifestation of symptoms after methamphetamine use, such as violent and violent behavior, increased energy and movement, feelings of well-being, or increased safety and confidence. Similarly, methamphetamine use reduces sleep and fatigue, increases the speed of thought and speech, suppresses appetite, increases blood pressure and heart rate, and raises body temperature. Amphetamine is a substance that can be used to synthesize designer drugs that achieve these pleasurable effects and increase energy and arousal [8-10].

Amphetamine causes a general stimulation of the central nervous system, so an additional increase in arousal can sometimes be expressed in a feeling of nervousness. This is probably one of the most discussed side effects of using this substance to treat ADHD. Children experiencing this psychological change suffer from attention deficits, so amphetamine use is very suitable for increasing the attention span of these children. However, amphetamine makes the body too healthy, so it can increase a person's anxiety and nervousness, which is not good for children with high hyperactivity [11-12].

Cognitive-enhancing stimulant drugs, designed to improve cognitive functions such as memory, attention, and executive function, are increasingly prevalent in both clinical and non-clinical settings. This article explores the scientific discussions surrounding the characteristics of these drugs, focusing on their efficacy and safety. Emphasis is placed on the safety of pharmacotherapy and the role of pharmacovigilance systems in monitoring and managing potential risks associated with cognitive enhancers. By examining current research and regulatory practices, the article aims to provide a comprehensive overview of the challenges and considerations in ensuring the safe use of these substances [13-14].

Another of the most common side effects of amphetamines is difficulty falling asleep and staying asleep. The reason for this effect is the same as the previous one, that is, the overstimulation that these substances create on the brain. Amphetamine use means that brain functions are increased and become more agitated than usual, so if the doses taken are not adapted to the characteristics of the person, they can cause insomnia relatively easily. In these cases, stimulation to increase attention and concentration makes the brain more alert when it should be resting, making it harder to

fall asleep. Likewise, after you fall asleep, the brain may continue to work over actively, so early awakenings may also occur [15-16].

It is already known that ecstasy, which includes mainly MDMA (methylenedioxyamphetamine) and other amphetamines represent recreational drugs with high potential for abuse worldwide and in Georgia. Ecstasy is manufactured in iillegal laboratories and it is never known how pure its chemical composition is (it may contain cocaine, ketamine, marijuana, methamphetamine or synthetic cathinones and etc.), which further increases the risk of health risks posed by ecstasy consumption [17-18].

Amphetamines, including MDMA, are products of receptive phenylisopropilamines which have a strong stimulating effect on the CNS; They represent sympathomimetics of indirect actions, that release serotonin, dopamine and other monoamines in the CNS; The action of these excess neurotransmitters and their increased consumption are the basis for the development of MDMA's life-threatening side effects and toxicity in both episodic and prolonged, chronic use. The main target for MDMA is brain, in particular, the short- and long-term effects of ecstasy on the brain: short-term effects of ecstasy on the brain include chemical and behavioral changes of the brain, and the long-term effects of ecstasy include changes of brain structure, mood, memory and behavior. Unfortunately, there is no FDA-recommended medication or specific treatment available to deal with MDMA addiction [19-20].

The most effective current treatment for patients with MDMA addiction is cognitive-behavioural interventions designed to alter the patient's thinking, expectations and behaviour, as well as increase their ability to cope with life's stressful situations. Research and clinical perspectives differ in how addictive ecstasy is or whether it is addictive at all. There are many people who claim to be addicted to ecstasy, but clinical research on the subject is less extensive than many other medications. A number of studies have shown that ecstasy is addictive, albeit to a lesser extent than many other drugs. However, this does not mean that developing ecstasy addiction is not a serious problem. Detecting an addiction to ecstasy can be difficult because the drug has no obvious withdrawal symptoms and is often used by young people in social settings. Regardless, continued use of ecstasy can lead to psychological and physical dependence. If someone is addicted to ecstasy, they will continue to take the drug despite knowing the consequences. Even if ecstasy isn't the only drug that causes addiction, it can be part of a multi-drug patchwork of addiction [21-23].

Cognitive-enhancing stimulant drugs have gained significant attention for their potential to improve cognitive functions such as memory, attention, and executive functioning. This review provides an in-depth analysis of the characteristics, mechanisms, efficacy, and societal implications of these drugs. By examining both prescription and non-prescription stimulants, we aim to offer a comprehensive understanding of their role in cognitive enhancement, their benefits and risks, and the ethical and societal issues associated with their use [24-26].

Aim. The aim of the research was to study and analyze the key issue aspects related to features of effects of using psychostimulant substances.

Methods. The material of the article was the data from scientific publications, which were processed, analyzed, overviewed and reviewed by generalization and systematization. research studies are based on a review/overview assessment of the development of critical visibility and overlook of the modern scientific literature. use the following databases: (for extensive literature searches to identify the key issue aspects related to features of effects of using psychostimulant substances). PubMed, Web of Science, Clinical Key, Tomson Reuters, Google Scholar, Cochrane library, and Elsevier foundations. national and international policies and guidelines were also reviewed.

Results and Discussion. Cognitive-enhancing stimulant drugs have garnered significant attention for their potential to improve cognitive performance. These substances, including prescription medications and over-the-counter nootropics, are used to enhance various cognitive functions. Despite their promise, there are growing concerns about their safety and the need for effective pharmacovigilance to monitor and mitigate risks. This article reviews the scientific discussions on the characteristics of cognitive-enhancing stimulants, their safety profiles, and the role of pharmacovigilance systems in ensuring safe pharmacotherapy [27-28].

The pursuit of cognitive enhancement through pharmacological means has seen considerable advancements with the development of various stimulant drugs. These substances, ranging from well-established prescription medications to emerging nootropics, promise to enhance cognitive functions beyond normal levels. This review synthesizes current scientific knowledge regarding cognitive-enhancing stimulants, providing insights into their mechanisms, efficacy, safety, and the broader societal implications of their use [29-30].

Cognitive-enhancing stimulant drugs offer significant potential for improving cognitive functions, yet their use is accompanied by important safety considerations. Ensuring the safe use of these substances requires a thorough understanding of their characteristics, efficacy, and potential risks. Pharmacovigilance systems play a crucial role in monitoring drug safety and addressing emerging concerns. Ongoing research and advancements in safety monitoring will be essential in managing the risks associated with cognitive enhancers and ensuring their safe and effective use [31-32].

The pharmacokinetics and pharmacodynamics of psychostimulants are discussed elsewhere in this issue. In brief, methamphetamine is structurally very similar to amphetamine and related agents, such as MDMA (3,4-methylenedioxy-*N*-methylamphetamine), which is widely known as “ecstasy,” and designer drugs, such as cathinone derivatives (including “bath salts”). The amphetamines, including methamphetamine, are similar to cocaine in causing an increase in synaptic levels of the monoamine neurotransmitters dopamine (DA), norepinephrine (NE), and serotonin, in both the central nervous system (CNS) and peripheral nervous system (PNS) [33-34].

Methamphetamine has two differentially active stereoisomers. The L-enantiomer has peripheral alpha-adrenergic activity and has been used as a nasal decongestant. The D-enantiomer is a powerful stimulant with 3–5 times the CNS activity of the L-enantiomer. It can be insufflated, smoked, injected, or taken orally or per rectum and is highly lipophilic, so it enters the brain faster than other psychostimulants (including amphetamine) and is less vulnerable to enzymatic degradation in the brain; its half-life is 10–12hours [35-36].

Regardless of the route of administration, or even the specific drug used, psychostimulants produce the same basic spectrum of acute CNS effects: euphoria, increased energy/decreased fatigue, reduced need for sleep, decreased appetite, decreased distractibility, increased self-confidence and alertness, increased libido, and prolonged orgasm. Peripheral effects may include tremor, diaphoresis, increased muscle tension, tachypnea, hyperreflexia, and hyperpyrexia [37-38].

Many of the effects are biphasic; for example, low doses improve psychomotor performance while higher doses may cause tremors or convulsions. Cardiovascular effects (alpha-adrenergically mediated) are also biphasic, with low doses decreasing heart rate via the vagus nerve and high doses causing increased heart rate and vasoconstriction, leading to increased blood pressure. Over several hours, days, or weeks, psychostimulant use can result in restlessness, irritability, and insomnia and at higher doses, suspiciousness, repetitive stereotyped behaviors, and bruxism [39-40].

The therapeutic goal of amphetamine-based drugs is to control the actions of only certain areas of the brain, this is quite difficult to do. As we have seen, amphetamine is a substance that spreads easily to areas of the brain, so although it has its greatest effect in areas where it has a therapeutic effect, amphetamine has other areas of the brain as well. Although amphetamine acts on the cerebral cortex or higher brain structures, it can also have an effect, albeit a minor one, on the lower brain regions, while being useful for increasing attention and concentration. Many of these areas are known as subcortical because they are below the cerebral cortex and control body movements, so if amphetamines hit these areas of the brain too hard, it can cause tremors in various parts of the body [41-42].

Once in our brain, amphetamines can cause headaches and discomfort in that area of the body. In general, amphetamine abuse, i.e methamphetamine use, causes persistent headaches for several hours after use (just like alcohol or other drugs). Changes in sexual desire or the capacity for sexual impulses, desires and processes in the brain and the functioning of various areas of the brain begin. In particular, sexual behavior is associated with the action of serotonin, a substance presents in many parts of the brain. As we have seen, amphetamine regulates the activity of serotonin by inhibiting the return of this neurotransmitter from the brain. Therefore, the action of amphetamine tries to achieve other effects by modifying serotonin, such as increasing well-being or improving concentration (functions performed by this neurotransmitter), which can cause negative consequences of a sexual nature [43-44].

The effects produced by amphetamines are not limited to the mental component, but also change the physical functioning of the body. When the brain is

“overstimulated” by this substance, it stops doing other things, like preparing the body for food. This fact can cause many symptoms, but one of the most typical is a feeling of dryness in the mouth, much like when we experience anxiety. Constipation or diarrhea. Thus, constipation, diarrhea or other digestive disorders can be detected. This can be explained by the fact that although amphetamines regulate cognitive (such as the ability to concentrate) or psychological (such as feelings of well-being), they also modulate parts of the brain that are responsible for basic functional processes, such as Loss of appetite is one of the effects of amphetamine use. That - is one of the main symptoms, to the point that these substances are sometimes used to treat problems associated with amphetamine use or being overweight. This fact, as with sexual changes, is largely due to the action of serotonin in our body [45-46].

Amphetamine mainly affects this substance, since it is closely associated with attentional processes and other cognitive activities, which allows the drug to have a therapeutic effect. However, serotonin does other things as well, such as balancing sexual desire, regulating appetite through satiety, or controlling body temperature. Thus, when amphetamines are taken, the function of this neurotransmitter is altered, and although its stronger effects are beneficial, it is often difficult to control these negative effects [47-48].

Loss of appetite - this symptom is closely related to amphetamines, since amphetamines cause loss of appetite continuously, weight loss occurs gradually. In these cases, it is very important to consult a doctor, because if the drug is not used for weight loss, if the appetite is excessively reduced, the dose and course of treatment with amphetamine should be restored. Addiction Finally, the last symptom we'll look at is amphetamine addiction. Although found in small amounts and strictly controlled by a physician when consumed, amphetamine has a high addictive component. This action is mediated primarily by dopamine, a neurotransmitter that is much more effective in recreational amphetamines because the formulations of these drugs are addictive to the user. In amphetamines for medical use, the narcotic component is much more controlled and they try to reduce it as much as possible, and not as much as possible, although dopamine is somewhat modified, so its use can be addictive. Condition, processes of attention and the ability to concentrate. The controlled and therapeutic use of amphetamines can increase higher cognitive functions (attention and memory) and improve the executive functions of a person. Similarly, amphetamines, when used therapeutically, also have a reinforcing effect, as they induce pleasant emotions and strengthen the systems involved in the regulation of specific emotional responses: reducing impulsivity and increasing self-control. On the other hand, since these substances suppress appetite, the use of amphetamines in very controlled amounts can be useful in the treatment of eating disorders and obesity. However, despite the controlled use of these substances, the use of amphetamines can cause side effects, so these drugs should be used very carefully [49-50].

Lysergic acid diethylamide, or LSD, is a product of processing ryegrass, a fungus that parasitizes on ryegrass. For example, dihydroergotamine has a great effect on the human psyche.

LSD-hallucinogenic effect begins 30-60 minutes after ingestion. The action lasts 8-12 hours. Effects: visual hallucinations, increased hearing, disorientation in time, a feeling of lightness or panic fear, clouding of consciousness, aggression, auto-aggression, suicidal thoughts. Overdose signs: increased sensitivity to light, touch and sounds; burning on the face; the so-called goosebumps; dry mouth; severe sweating; nausea, diarrhea; dizziness, drowsiness; severe general weakness; a sharp expansion of the pupils; increased blood pressure, increased body temperature. Signs of severe poisoning: respiratory failure; heartbeat; vomit; arrhythmia; hyperemia; coma (rare) hemorrhage in the brain [51-52].

Prescription stimulants – medications such as amphetamines (e.g., adderall) and modafinil, used primarily for conditions like ADHD and narcolepsy.

Non-prescription nootropics: over-the-counter substances like caffeine and various nootropic supplements claimed to improve cognitive function.

Mechanisms of action – pharmacodynamics: interaction with neurotransmitter systems such as dopamine and norepinephrine, and effects on brain regions associated with cognition. Clinical benefits: evidence of cognitive enhancement in individuals with specific conditions like ADHD or narcolepsy. Variability in efficacy: differences in responses among individuals, influenced by genetic, psychological, and environmental factors [53-54]. Common adverse effects: insomnia, jitteriness, cardiovascular issues, and gastrointestinal problems. Long-term risks: potential for dependence, cognitive impairment with prolonged use, and other health concerns.

Non-medical use: risks associated with the use of cognitive enhancers by healthy individuals, including overuse and potential for abuse. Lack of long-term data: insufficient research on long-term effects and safety in healthy populations [55-56].

Cannabis, also called marijuana or plana among other names, is a psychoactive substance derived from the hemp plant and used for medical and recreational purposes. at least 65 other cannabinoids - elements that cause euphoria /narcotic effect. Marijuana is often taken for its mental and physical effects, such as feeling "high" and "dizzy". Effects also include a general change in perception, euphoria (pleasure and happiness), appetite. These effects usually begin within minutes of smoking or about 30 to 60 minutes after food preparation and ingestion and typically last two to six hours. Short-term side effects include impaired short-term memory, dry mouth, slow motor skills, red eyes, and feeling paranoid or agitated [57-58].

In general, the medicinal use of the cannabis plant has a long history across cultures; According to one of the projects of the US National Library of Medicine, there is information about its alleged use in materials 5000 years old. As of 2015, 23 US states, including Washington, allow the sale of medical marijuana; In the same year, the country's medical cannabis industry topped a billion dollars. With the rise of cannabis use, scientists are trying to delve deeper into the benefits and risks of its medical use; As the US National Institute on Drug Abuse explains, medical cannabis is sometimes used to treat the symptoms of an illness; Some studies conclude that

cannabis can reduce nausea and vomiting during chemotherapy, improve appetite in people with HIV/AIDS, relieve chronic pain, muscle spasms, etc.

Cannabis use is accompanied by psychoactive and physiological effects. The immediate effects of cannabis use include relaxation and euphoria, a general change in conscious perception, heightened sensitivity, increased libido, and distortions in the perception of time and space. At higher doses, body image changes, auditory and/or visual illusions, pseudo hallucinations and ataxia may occur due to selective impairment of polysynaptic reflexes. In some cases, marijuana can cause dissociative states such as depersonalization and derealization [59-60].

Ongoing research is needed to address gaps in our understanding of cognitive-enhancing stimulants. Long-term studies are essential to assess the safety and efficacy of these drugs over extended periods. Research should also focus on different populations, including healthy individuals and those with varying health conditions, to better understand the broader implications of cognitive enhancers.

Cognitive-enhancing stimulant drugs offer promising benefits for improving cognitive functions, but their use is accompanied by significant safety considerations. A thorough understanding of their characteristics, efficacy, and potential risks is essential for ensuring their safe use. Pharmacovigilance systems play a critical role in monitoring drug safety and addressing emerging concerns. Continued research and advancements in safety monitoring are vital for managing the risks associated with cognitive enhancers and ensuring their responsible use.

Recent advancements in technology offer new opportunities for improving pharmacovigilance. Innovations such as electronic health records and real-time monitoring systems can enhance the detection and management of adverse effects. Developing more comprehensive surveillance systems will be crucial for addressing emerging safety concerns.

Pharmacovigilance systems face several challenges, including underreporting of adverse events and difficulties in data interpretation. Incomplete reporting and lack of awareness among users and healthcare providers can hinder the detection of safety concerns. Enhancing reporting practices and improving data analysis are critical for addressing these challenges.

Pharmacovigilance practices are governed by global standards set by organizations such as the FDA, EMA, and WHO. These regulations require rigorous monitoring and reporting of adverse effects. However, differences in regulatory practices across countries can affect the consistency and effectiveness of safety monitoring.

Pharmacovigilance systems are essential for monitoring the safety of cognitive enhancers. These systems collect and analyze data on adverse effects, enabling timely detection and management of potential risks. Effective pharmacovigilance helps ensure that the benefits of cognitive enhancers outweigh their risks.

The use of cognitive enhancers among healthy individuals introduces additional risks. Non-medical use of these substances can lead to overuse, abuse, and potential health issues. The lack of long-term safety data for healthy users underscores the need for caution and further research.

While cognitive enhancers offer potential benefits, they are also associated with various side effects. Common adverse effects include insomnia, jitteriness, and cardiovascular issues. Long-term use of these drugs may lead to more serious health concerns, such as dependence and cognitive impairment. The safety profile of these drugs in healthy individuals is less well-established, particularly with non-prescription substances.

Prescription cognitive enhancers have demonstrated efficacy in clinical settings, particularly for treating ADHD and narcolepsy. Studies show that these drugs can significantly improve cognitive functions in individuals with these conditions. However, the variability in individual responses highlights the need for personalized treatment approaches.

Cognitive enhancers exert their effects through interactions with neurotransmitter systems. For instance, amphetamines increase the release of dopamine and norepinephrine, which enhances alertness and focus. Modafinil, a wakefulness-promoting agent, affects various neurotransmitter systems to improve cognitive function. The pharmacokinetics of these drugs, including their absorption, distribution, metabolism, and excretion, influence their efficacy and safety profiles.

Cognitive-enhancing stimulants include a range of substances from prescription medications to over-the-counter nootropics. Prescription stimulants, such as amphetamines (e.g., Adderall) and modafinil, are primarily used to treat conditions like ADHD and narcolepsy. Non-prescription nootropics, including caffeine and various dietary supplements, are commonly used by healthy individuals seeking cognitive improvements.

Cognitive-enhancing stimulant drugs, including both prescription medications and non-prescription substances, are increasingly used to improve cognitive performance. These drugs promise enhancements in various cognitive domains, but their use also raises important safety and ethical concerns. This review aims to elucidate the scientific discussions surrounding these drugs, focusing on their characteristics, the safety profile of pharmacotherapy, and the critical role of pharmacovigilance systems.

Cognitive-enhancing stimulant drugs have emerged as significant tools for improving cognitive functions such as memory, attention, and executive function.

Other unwanted side effects include short-term memory loss, dry mouth, dysmotility and eye redness. In addition to subjective changes in perception and mood, the most common short-term physical and neurological effects include increased heart rate, increased appetite, decreased blood pressure, impaired short-term and working memory, psychomotor coordination and concentration. Some users may experience an acute episode of psychosis that usually resolves within six hours, but in rare cases, long-term drug users may experience symptoms for many days.

Marijuana is used in a variety of ways: During smoking, Cannabis burns and as a result a person inhales its smoke. Currently, many ways and techniques for smoking cannabis have been invented: bong, joint, pipe, paper wrap, vape (electronic cigarette), “bulbulator”, etc. Tablets or capsules are also often used by addicted peo-

ple. Capsules have become a popular method of consuming marijuana. The tablets contain marijuana. The benefits of this path are suitable for people with breathing problems. It also provides accurate dosage and can be used to treat many conditions [61-62].

Physiological effects that occur after consuming marijuana: inflammation of the mucous membrane of the eye, inflammation of the eye; increased pulse rate, increased heart rate; an increase or decrease in blood pressure; disrupts movement functions; drowsiness; changes in body temperature; headaches; vertigo; nausea; feelings of hunger.

The effects that marijuana causes on human behavior: promotes the relaxation process; slows down psychomotor activity; worsens the concentration of attention, interferes with the ability to assess the situation adequately; reduces the ability to drive a vehicle and all activities that require attention; causes acceleration of speech and unrestrained speech;

Emotional effects caused by marijuana: euphoria, a feeling of well-being; a state of carelessness that alternates between restlessness and inner anxiety.

Neuropsychological effects caused by marijuana: deterioration of short-term memory, understanding and ability to perform any task; loss of sense of time and space; apathy; visual, auditory and olfactory sensations; lack of motivation; lowers the pain threshold; changes in sexual emotions; hallucinations; the loss of the ability to perceive oneself as a person; psychosis (especially a large dose).

Marijuana affects thinking, the ability to understand, abstract thinking, the ability to learn. Marijuana, depending on the dose and the user's ability to absorb it, can cause panic, anxiety, and toxic psychosis. Its prolonged use can cause structural and functional changes. The use of marijuana causes a number of harmful effects characteristic of tobacco smokers: bronchitis, pharyngitis, sinusitis, contributes to diseases of the upper respiratory tract and canals.

In case of taking ecstasy on fasting, it needs 15 minutes to reach the brain. MDMA enhances the activity of three transmitters in the brain: dopamine, norepinephrine, serotonin. The latter is responsible for the "fight or run" reaction. In addition, they stimulate the ways of so-called "self-preservation". These transmitters are characterized by the following effects: dopamine–produces an increased energy/activity and acts to enhance behavioural activities in the reward system; norepinephrine - increases heart rhythm and blood pressure, which is especially dangerous for people with cardiovascular problems; serotonin - it affects the mood, appetite, sleep, also stimulates sexual arousal and strengthen confidence. High levels of serotonin can cause emotional closeness, elevated mood, and empathy in those who use MDMA [63-64].

Generally, amphetamines are characterized by high abuse potential. They represent recreational drugs and are characterized mainly by central rather than peripheral effects; Frequently, an ecstasy tablet with MDMA contains antihistamines and caffeine, and in the worst case, cocaine or ketamine. MDMA enhances not only the release of serotonin, but also increases its consumption in the body; It is also

worth noting that hangover is usually accompanied by bad mood or depression, because, as mentioned above, high levels of serotonin are consumed throughout consumption of MDMA and during hangover the body suffers from lack of serotonin. Serotonin is known as a transmitter of good mood and pleasant feeling, which is mainly concentrated in the central nervous system and intestines [36-65].

Short-term effects of ecstasy on the brain include chemical and behavioural changes of the brain. Long-term effects of ecstasy include changes of brain structure, mood, memory, and behaviour. Ecstasy, like other drugs, have an effect on the so-called "reward pathway" located in the limbic system of the brain, which explains its psychostimulant and hallucinogenic action. The effect of MDMA lasts from 3 to 6 hours. People after taking of MDMA: have a positive attitude towards themselves and those around them; energy increases; self-limitations are diminished; the ego softens and emotional bonding - empathogenic establishment even with a stranger becomes easier [38,39,46].

So, the majority of those taking this drug are euphorically happy, consequently, open and friendly towards the outside world. The main basis of the short-term effect of ecstasy, i.e., on the target in brain, is the serotonin system, in particular, it transports serotonin from neuronal terminals to the synaptic space along with other neurotransmitters, leading to an elevated mood, a feeling of empathy, and a sense of pleasure that may be accompanied by a vague thinking, agitating, and disruptive behaviour. Then there is the feeling of withdrawal from the body of drug and longing for repeated use of ecstasy and consequently, increasing the threshold of pleasure (i.e., decreased reaction). After 2 weeks of taking ecstasy, level of serotonin in the brain decreases dramatically, and after years, its level will not be completely restored, which disrupts the neuronal connections and it actually causes an abnormal change in mood, behaviour and memory [40, 58, 62].

During a week of moderate use of the drugs, a person may experience: anxiety, sleep problems, impulsivity, aggression, and depression; decreased appetite; memory and attention problems; diminished interest and pleasure in sex, excessive thirst. persistent or prolonged use of ecstasy and after its withdrawal from the body causes: degeneration of nerve terminals which is primarily reflected in in impaired memory (impaired verbal and visual memory) and behavioural impairment due to a sharp decrease in serotonin and its metabolites in the neocortex and hippocampus, sleep disturbance and depression [15, 28].

Life-threatening effects after taking multiple doses: some people take multiple doses of ecstasy overnight; increased doses of ecstasy increase the risk of developing side effects. Clinicians have observed that episodic use of ecstasy in young people in the future develops alcohol dependence and chronic alcoholism, especially among young people with genetic predisposition. Taking of repeated or high doses of ecstasy can result in hyperthermia, hypertension, cardiac arrhythmia, impairment of functional activity of muscles and kidney failure due to salt and fluid reduction. The development of these dangerous side effects may primarily be due to the impact of

ecstasy on the brain, in particular the hypothalamus; severe hyperthermia can cause brain damage or even death [4-27].

So, the brain predominantly is affected by ecstasy. For people taking ecstasy is especially dangerous: a sharp rise in body temperature (hyperthermia, even up to 42 degrees Celsius), dehydration, hypercoagulation and severe cardiovascular events. Studies conducted on monkeys made it clear that ecstasy mainly affects the brain. Overdose causes: serotonin syndrome (muscle rigidity, hyperthermia, cardiovascular disorders, seizures), psychosis, hypertensive crisis [34, 38].

Ecstasy tablets, capsules, or powder sold as MDMA-containing are likely to consist of other drugs instead of MDMA, or may be combined with MDMA; for example, such as cocaine, ketamine, methamphetamine, or synthetic cathinones (so-called "sea salt"). These substances can be very dangerous if the person does not know what he/she is taking. People who purposely or unknowingly mix such a mixture with other substances, such as marijuana and alcohol, may pose an even greater risk for their health. Active self-administration of MDMA by experimental animals - was an important indicator of the potential for drug addiction, although it was expressed to a lesser degree than in the case of other drugs, e.g., such as cocaine. Some people observed symptoms of the drug addiction, which include the following symptoms: fatigue, loss of appetite, depression, difficulty in concentrating [12, 33].

MDMA causes increased production of prolactin, oxytocin, adrenocorticotropic hormone, Vasopressin. Thus, the consumption of MDMA includes both pleasure and discomfort (rise in body temperature, abrupt change in blood pressure and heart rate, dry mouth and shivering, blurred vision, involuntary movement of muscles, especially the lower jaw muscle – causing clenching and damaged teeth, in some cases panic attack and hallucination). Unfortunately, there is no FDA-recommended medication or specific treatment available to deal with MDMA addiction. The most effective current treatment for patients with MDMA addiction is cognitive-behavioural interventions designed to alter the patient's thinking, expectations and behaviour, as well as increase their ability to cope with life's stressful situations. Supportive healing groups can be effective in combination with behavioural interventions to support long-term recovery [50, 51, 63].

Research and clinical perspectives differ in how addictive ecstasy is or whether it is addictive at all. There are many people who claim to be addicted to ecstasy, but clinical research on the subject is less extensive than many other medications. A number of studies have shown that ecstasy is addictive, albeit to a lesser extent than many other drugs. However, this does not mean that developing ecstasy addiction is not a serious problem. Detecting an addiction to ecstasy can be difficult because the drug has no obvious withdrawal symptoms and is often used by young people in social settings. Regardless, continued use of ecstasy can lead to psychological and physical dependence. If someone is addicted to ecstasy, they will continue to take the drug despite knowing the consequences. Even if ecstasy isn't the only drug that causes addiction, it can be part of a multi-drug patchwork of addiction.

Cocaine is a powerful stimulant. It enhances thinking, causes euphoria, and at this time people experience a feeling of omnipotence. High doses can cause serious, life-threatening disorders such as heart attacks and strokes. Diagnosis is confirmed by urinalysis. Use of intravenous sedatives such as lorazepam may relieve symptoms. People who have stopped taking medication need constant monitoring as they are suicidal and also need extra help to get rid of the need for medication.

The effects of cocaine are similar to those of amphetamines. It is used as a powder for inhalation through the nose, for injection into a vein or for inhalation of heated steam. When cocaine is boiled with sodium bicarbonate, it turns into its base form, called refined crystallized cocaine. When heated, purified cocaine releases a vapor that can be inhaled. Inhaling purified cocaine is commonly referred to as smoking, although it is not actually burned. The effect of refined cocaine is almost as fast as that of intravenous cocaine. Heavy cocaine users who inject or smoke cocaine are more likely to become addicted. Occasional and occasional users of cocaine, as well as people using it as a nasal powder, are less likely to become addicted to cocaine. Cocaine can contain excipients, impurities, and infectious agents that, when injected, can cause complications such as infections. Symptoms- Quick effect: cocaine causes a sharp change in consciousness, euphoria. It is especially effective when administered intravenously or inhaled. These phenomena are expressed with less intensity when using cocaine nasally, in the form of a powder. Because the effects of cocaine are short-lived, users may inject, smoke, or snort every 15 to 30 minutes. For users who frequently use cocaine for only a few days or more, it can cause fatigue, exhaustion, and a desire to sleep. Complications**:** An acute toxic reaction caused by cocaine can be fatal. Cocaine causes an increase in blood pressure, an increase in heart rate, and constriction of blood vessels. There may be a heart rhythm disorder (arrhythmia). The effects of cocaine on the heart can lead to chest pain, heart attack (even in healthy young athletes), or sudden death. Cocaine also causes kidney failure, stroke, and lung problems [52, 55, 58].

High doses (overdose) can impair thinking, cause tremors, extreme nervousness, seizures, hallucinations, insomnia, paranoia, anxiety, and aggressive behavior. People experience profuse sweating and dilated pupils. An overdose can cause a sudden rise in temperature (hyperthermia) to a life-threatening level. Later effects: Long-term users of cocaine may develop a tolerance to it, and more and more doses will be needed to achieve the initial effect. Long-term use can lead to damage to the tissues of the nasal septum with the development of ulcers requiring surgical intervention. Heavy users may experience impairment of mental functions, including attention and memory.

The addictive properties of psychostimulants such as cocaine, amphetamine, methamphetamine, and methylphenidate are based on their ability to increase dopaminergic neurotransmission in the reward system. While cocaine and methamphetamine are predominately used recreationally, amphetamine and methylphenidate also work as effective therapeutics to treat symptoms of disorders including attention deficit and hyperactivity disorder (ADHD) and autism spectrum disorder (ASD). Alt-

hough both the addictive properties of psychostimulant drugs and their therapeutic efficacy are influenced by genetic variation, very few genes that regulate these processes in humans have been identified. This is largely due to population heterogeneity which entails a requirement for large samples. Drosophila melanogaster exhibits similar psychostimulant responses to humans, a high degree of gene conservation, and allow performance of behavioral assays in a large population. Additionally, amphetamine and methylphenidate reduce impairments in fly models of ADHD-like behavior. Treatment of psychostimulant1 addiction has been a major, and not fully met, challenge. For opioid addiction, there is strong evidence for the effectiveness of several medications, including methadone, buprenorphine, and naltrexone. For psychostimulants, there is no corresponding maintenance medication that has met criteria for regulatory approval or generally accepted use. The mainstay of treatment has been behavioral interventions such as cognitive-behavioral therapy (CBT). In this paper, we will discuss advances in pharmacological as well as behavioral treatments for psychostimulant addiction [22, 29, 57].

If a woman uses cocaine during pregnancy, the fetus is more likely to have problems. However, it should be noted that there are many other risk factors in pregnant women, such as tobacco smoking and alcohol use, nutritional deficiencies, and problems with prenatal care. Doctors believe that these risk factors are sometimes more important than cocaine in developing problems. Withdrawal Syndrome: The reactions that develop as a result of stopping cocaine use (withdrawal symptoms) include severe weakness and depression - in contrast to reactions while taking the drug. Appetite increases, people have problems with the ability to concentrate. With the abolition of the drug, suicidal urges increase. Even after a few days, when the mental and physical state returns to its original level, drug addicts may try to commit suicide.

Emergency care: Since the effects of cocaine are short-lived, treatment of existing unpleasant symptoms is usually not required. People who are agitated, anxious, develop seizures, or have high blood pressure are given benzodiazepines (sedatives) such as lorazepam intravenously. Nitrates and other antihypertensive drugs can be given intravenously to lower blood pressure and decrease heart rate. Antihyperthermic treatment may also be required.

Detoxification and rehabilitation. Long-term cocaine use may require constant monitoring as people become depressed and suicidal. It may be necessary to place them in medical facilities or drug treatment centers. Psychotherapy is the most effective treatment for cocaine addiction. There are many support groups and telephone hotlines that help people recover from cocaine use.

Occasionally, cocaine addicts, who often have psychiatric disorders such as depression, may be put on appropriate drug addiction treatment.

3,4-methylenedioxy-methamphetamine (MDMA) is a synthetic drug that alters mood and perception (awareness of surrounding objects and conditions). It is chemically similar to both stimulants and hallucinogens, producing feelings of increased energy, pleasure, emotional warmth, and distorted sensory and time perception.

MDMA was initially popular in the nightclub scene and at all-night dance parties ("raves"), but the drug now affects a broader range of people who more commonly call the drug ecstasy or molly.

MDMA is an abbreviation for methylenedioxy-methylamphetamine. The formal (IUPAC) name is N-methyl-1-(3,4-methylenedioxyphenyl)propan-2-amine, but MDMA (CAS-42542-10-09) is commonly known as 3,4-methylenedioxy-methamphetamine or methylenedioxy methylamfetamine. Other chemical names include N, α-dimethyl-3,4-methylenedioxyphenethylamine or, less usually, N-methyl-1-(1,3-benzodioxol-5-yl)-2-propanamine. MDMA is a member of the larger group of ring-substituted phenethylamines. As with other phenethylamines, and like its close relative methamphetamine, MDMA also exists in two enantiomeric forms (R and S) [5, 36, 57].

MDMA increases the activity of three brain chemicals: dopamine (produces increased energy/activity and acts in the reward system to reinforce behaviors); norepinephrine (increases heart rate and blood pressure, which are particularly risky for people with heart and blood vessel problems); serotonin (affects mood, appetite, sleep, and other functions). It also triggers hormones that affect sexual arousal and trust. The release of large amounts of serotonin likely causes the emotional closeness, elevated mood, and empathy felt by those who use MDMA.

Whereas phenethylamines without ring substitution usually behave as stimulants, ring substitution (as in MDMA) leads to a modification in the pharmacological properties. Ingestion of MDMA causes euphoria, increased sensory awareness and mild central stimulation. It is less hallucinogenic than its lower homologue, methylenedioxyamphetamine (MDA). The terms empathogenic and entactogenic have been coined to describe the socializing effects of MDMA. Following ingestion, most of the dose of MDMA is excreted in the urine unchanged. Major metabolites are 3,4-methylenedioxyamphetamine (MDA) and *O*-demethylated compounds. Following a dose of 75 mg, the maximum plasma concentration of around 0.13 mg/L is reached within two hours. The plasma half-life is 6–7 hours. In animals, MDMA causes neurotoxicity, as evidenced by anatomical changes in axon structure and a persisting reduction in brain serotonin levels. The significance of these findings to human users is still unclear, although cognitive impairment is associated with MDMA use. Some of the pharmacodynamic and toxic effects of MDMA vary, depending on which enantiomer is used. However, almost all illicit MDMA exists as a racemic mixture. Fatalities following a dose of 300 mg have been noted, but toxicity depends on many factors, including individual susceptibility and the circumstances in which MDMA is used.

Ecstasy is considered as heavy drug and nowadays, its manufacturing, storage or realization is forbidden in all countries of the world, including Georgia. Internationally, it is regulated by the UN Convention on Psychotropic Substances. Ecstasy is produced in illegal laboratories and it is never known how pure its chemical composition is. Often, ecstasy contains various additives that modify MDMA's activity in different ways.

It is already known that ecstasy, which includes mainly MDMA (methylenedioxyamphetamine) and other amphetamines represent recreational drugs with high potential for abuse worldwide and in Georgia. Ecstasy is manufactured in illegal laboratories and it is never known how pure its chemical composition is (it may contain cocaine, ketamine, marijuana, methamphetamine or synthetic cathinones and etc.), which further increases the risk of health risks posed by ecstasy consumption. Amphetamines, including MDMA, are products of receptive phenylisopropilamines which have a strong stimulating effect on the CNS. They represent sympathomimetics of indirect actions, that release serotonin, dopamine and other monoamines in the CNS. The action of these excess neurotransmitters and their increased consumption are the basis for the development of MDMA's life-threatening side effects and toxicity in both episodic and prolonged, chronic use. The main target for MDMA is brain, in particular, the short- and long-term effects of ecstasy on the brain: short-term effects of ecstasy on the brain include chemical and behavioural changes of the brain, and the long-term effects of ecstasy include changes of brain structure, mood, memory and behaviour [27, 41, 48].

Unfortunately, there is no FDA-recommended medication or specific treatment available to deal with MDMA addiction. The most effective current treatment for patients with MDMA addiction is cognitive-behavioural interventions designed to alter the patient's thinking, expectations and behaviour, as well as increase their ability to cope with life's stressful situations. The word ecstasy is used to describe a strong, euphoric feeling in a conversational language. According to psychologists definition, it is a special mental state, when a person is completely focused on the subject of interest and beyond it, he or she sees no time, no pain, no fatigue [27, 29, 64].

More often, ecstasy contains MDMA (Methyl-enedioxy-methamphetamine), which is a synthetic substance with a hallucinogenic and psychostimulant effect. As a result of its chemical and pharmaceutical processing, colored tablets and capsules are made. Other slang names of ecstasy are the following: X, XTC, Adam, Molly, E, etc. There are various methods of achieving ecstasy as a special mental state: dance, meditation, music, religious ritual, sex, and in the last 30 years, amphetamine, called MDMA. Ecstasy is taken by perorally, rarely by inhalation; Its excitationic and hallucinogenic action are manifested in a feeling of love, closeness and happiness; Warm and close relationships with those around them; In a sense of calm and complacency.

Treatment of psychostimulant addiction has been a major, and not fully met, challenge. For opioid addiction, there is strong evidence for the effectiveness of several medications. For psychostimulants, there is no corresponding form of agonist maintenance that has met criteria for regulatory approval or generally accepted use. Stimulant-use disorders remain prevalent and can result in both short-term and long-term adverse consequences. The mainstay of treatment remains behavioral interventions. In this paper, we discuss those interventions and some promising candidates in the search for pharmacological interventions.

The treatment of psychostimulant addiction has been a major, and not fully met, challenge. For opioid addiction, there is strong evidence for the effectiveness of several medications. For psychostimulants, there is no corresponding form of agonist maintenance that has met criteria for regulatory approval or generally accepted use. Stimulant-use disorders remain prevalent and can result in both short-term and long-term adverse consequences. The mainstay of treatment remains behavioral interventions. In this paper, we discuss those interventions and some promising candidates in the search for pharmacological interventions.

Cognitive-enhancing stimulant drugs represent a significant advancement in the pursuit of improved cognitive function, offering potential benefits across various domains including memory, attention, and executive function. These substances, which range from well-established prescription medications to emerging nootropics, have demonstrated varying degrees of efficacy and safety. The scientific evidence indicates that while cognitive enhancers can offer substantial improvements in specific contexts, their effects can be inconsistent and vary widely among individuals.

Prescription stimulants like amphetamines and modafinil have shown promise in clinical settings, particularly for conditions such as ADHD and narcolepsy. However, their non-medical use, particularly among healthy individuals seeking cognitive enhancement, raises important ethical and societal questions. The benefits of these drugs in healthy populations are often less clear-cut, with potential risks including side effects, dependency, and long-term health implications.

The ethical considerations surrounding cognitive enhancers are complex. Issues of fairness and equity, as well as the potential for coercion in competitive environments, highlight the need for careful regulation and ethical oversight. Societal impacts include the growing trend of cognitive enhancer use among students and professionals, which necessitates ongoing dialogue and policy development to address these emerging challenges.

Conclusion. In summary, while cognitive-enhancing stimulant drugs offer significant potential for improving cognitive function, their use must be approached with caution. A nuanced understanding of their mechanisms, efficacy, and ethical implications will be essential in harnessing their benefits while mitigating potential risks. Future research is essential to fill existing gaps in our understanding of cognitive-enhancing stimulants. Longitudinal studies, real-world evidence, and exploration of new drugs and technologies will be crucial in assessing the long-term effects and refining the applications of these substances. As the field of cognitive enhancement continues to evolve, a balanced approach that considers both the potential benefits and the associated risks will be vital in guiding future developments and ensuring the responsible use of these drugs.

References:

1. National Institute on Drug Abuse (NIDA). Monitoring the Future Study: Trends in Prevalence of Various Drugs for 8th Graders, 10th Graders, and 12th Graders; 2014 - 2017. Accessed Oct. 1, 2018 at https://www.drugabuse.gov/trends-statistics/monitoring-future/monitoring-future-study-trends-in-prevalence-various-drugs.

2. Ecstasy May Help Some PTSD Sufferers, but Safety Issues Remain. May 1, 2018. Drugs.com. Consumer News. Accessed Oct. 1, 2018 at https://www.drugs.com/news/ecstasy-may-help-some-ptsd-sufferers-but-safety-issues-remain-74011.html
3. National Institute on Drug Abuse (NIDA). MDMA (Ecstasy/Molly). Drug Facts. National Institute on Drug Abuse website. June 6, 2018. Accessed Oct. 1, 2018 at https://www.drugabuse.gov/publications/drugfacts/mdma-ecstasymolly.
4. Substance Abuse and Mental Health Services Administration. (2017). Key substance use and mental health indicators in the United States: Results from the 2016 National Survey on Drug Use and Health (HHS Publication No. SMA 17-5044, NSDUH Series H-52). Rockville, MD: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration. Retrieved Oct. 1, 2018. https://www.samhsa.gov/data/
5. Halpern, J. H., Sherwood, A. R., Hudson, J. I., et al. (2011). Residual neurocognitive features of long-term ecstasy users with minimal exposure to other drugs. Addiction,106: 777-86.
6. Newman A.H., Ku T., Jordan C.J., Bonifazi A., Xi Z.X. (2021). New drugs, old targets: tweaking the dopamine system to treat psychostimulant use disorders. Annu Rev Pharmacol Toxicol. 6;61:609-628.
7. Phillips K.A., Epstein D.H., Preston K.L. (2014). Psychostimulant addiction treatment. Neuropharmacology. 87:150-60.
8. Coffin P.O., Santos G.M., Das M. et al. (2013). Aripiprazole for the treatment of methamphetamine dependence: a randomized, double-blind, placebo-controlled trial. Addiction, 108(4): 751–761.
9. Cruickshank C.C., Dyer K.R. (2009). A review of the clinical pharmacology of methamphetamine. Addiction; 104(7): 1085–1099.
10. Cruickshank C.C., Montebello M.E., Dyer K.R. et al. (2008). A placebo-controlled trial of mirtazapine for the management of methamphetamine withdrawal. Drug and Alcohol Review. 27(3): 326–333.
11. Dackis C.A., Kampman K.M., Lynch K.G. et al. (2012). A double-blind, placebo-controlled trial of modafinil for cocaine dependence. Journal of Substance Abuse Treatment. 43(3):303–312.
12. Dackis CA, Lynch KG, Yu E, et al. (2003). Modafinil and cocaine: a double-blind, placebo-controlled drug interaction study. Drug and Alcohol Dependence.70(1):29–37.
13. Darker CD, Sweeney B, El Hassan H, et al. (2012). Cognitive behavioural coping skills therapy in cocaine using methadone maintained patients: a pilot randomised controlled trial. Heroin Addiction and Related Clinical Problems.14(3):101–110.
14. De Sousa A., De Sousa A. (2008). An open randomized trial comparing disulfiram and naltrexone in adolescents with alcohol dependence. Journal of Substance Use.13(6): 382–388.
15. Deas D., May K., Randall C., et al. (2005). Naltrexone treatment of adolescent alcoholics: An open-label pilot study. Journal of Child and Adolescent Psychopharmacology.15(5): 723–728.
16. Devoto P., Flore G., Saba P. et al. (2012). Disulfiram stimulates dopamine release from noradrenergic terminals and potentiates cocaine-induced dopamine release in the prefrontal cortex. Psychopharmacology. 219(4):1153–1164.
17. Dopheide J.A., Pliszka S.R. (2009). Attention-Deficit-Hyperactivity Disorder: An Update. Pharmacotherapy. 29(6):656–679.
18. Dutra L., Stathopoulou G., Basden S.L. et al. (2008). A meta-analytic review of psychosocial interventions for substance use disorders. American Journal of Psychiatry.165(2):179–187.
19. Eiler K., Schaefer M.R., Salstrom D., Lowery R. (1995). Double-blind comparison of bromocriptine and placebo in cocaine withdrawal. American Journal of Drug and Alcohol Abuse. 21(1):65–79.
20. Elkashef A.M., Rawson R.A., Anderson A.L. et al. (2008). Bupropion for the treatment of methamphetamine dependence. Neuropsychopharmacology. 33(5):1162–1170.

21. Ellinwood E.H., Jr, Lee T.H. (1989). Dose- and time-dependent effects of stimulants. NIDA research monograph. 94:323–340.
22. Epstein D.H., Hawkins W.E., Covi L. et al. (2003). Cognitive-behavioral therapy plus contingency management for cocaine use: Findings during treatment and across 12-month follow-up. Psychology of Addictive Behaviors. 17(1):73–82.
23. Fasano A., Barra A., Nicosia P. et al. (2008). Cocaine addiction: From habits to stereotypical-repetitive behaviors and punding. Drug and Alcohol Dependence. 96(1–2): 178–182.
24. Foltin R.W., Fischman M.W. (1997). A laboratory model of cocaine withdrawal in humans: Intravenous cocaine. Experimental and Clinical Psychopharmacology. 5(4): 404–411.
25. Galloway G.P., Buscemi R., Coyle J.R. et al. (2011). Randomized, Placebo-Controlled Trial of Sustained-Release Dextroamphetamine for Treatment of Methamphetamine Addiction. Clinical Pharmacology & Therapeutics. 89(2): 276–282.
26. Garbutt J.C. (2010). Efficacy and tolerability of naltrexone in the management of alcohol dependence. Current Pharmaceutical Design.16(19):2091–2097.
27. Gardner T.J., Kosten T.R. (2007). Therapeutic options and challenges for substances of abuse. Dialogues in clinical neuroscience. 9(4): 431–445.
28. Gawin F.H., Kleber H.D. (1986). Abstinence symptomatology and psychiatric-diagnosis in cocaine abusers - clinical observations. Archives of General Psychiatry; 43(2):107–113.
29. Gay G.R. (1982). Clinical management of acute and chronic cocaine poisoning. Annals of Emergency Medicine.11(10): 562–572.
30. George T.P., Chawarski M.C., Pakes J. et al. (2000). Disulfiram versus placebo for cocaine dependence in buprenorphine-maintained subjects: A preliminary trial. Biological Psychiatry. 47(12):1080–1086.
31. Giannini A.J., Malone D.A., Giannini M.C. et al. Treatment of depression in chronic cocaine and phencyclidine abuse with desipramine. Journal of Clinical Pharmacology. 2016; 26(3):211–214.
32. Goodwin J.S., Larson G.A., Swant J. et al. (2009). Amphetamine and methamphetamine differentially affect dopamine transporters in vitro and in vivo. Journal of Biological Chemistry. 284(5): 2978–2989.
33. Sopho Alavidze, Lela Sturua, Mzia Tabatadze, Davit Otiashvili (2016). Drug abuse in Georgia (Annual Report). Tbilisi.
34. Garcia V. A., De Freitas B. S., Busato S. B. et al. (2013). Differential effects of modafinil on memory in naïve and memory-impaired rats. Neuropharmacology. 75: 304–311.
35. García-Pardo M. P., De La Rubia, Ortí J. E., Calpe M. A. A. (2017). Differential effects of MDMA and cocaine on inhibitory avoidance and object recognition tests in rodents. Neurobiol. Learn. Mem. 146: 1–11.
36. Bourin, M., & Hascoët, M. (2003). The role of serotonin in cognitive processes. Psychopharmacology. 167(3): 319-328.
37. Kirkpatrick, B., & Buchanan, R. W. (2007). Cognitive enhancers in schizophrenia. Schizophrenia Research. 90(1-3): 147-155.
38. Ling, C. (2020). Modafinil: A review of its efficacy and safety for cognitive enhancement. Journal of Clinical Psychopharmacology. 40(2): 112-120.
39. Miller, R., & McGaugh, J. L. (2019). Cognitive enhancers: The pharmacology and psychology of memory improvement. Pharmacology Biochemistry and Behavior. 181: 20-29.
40. Neylan, T. C., & Metzler, T. J. (2004). The impact of cognitive enhancers on memory and attention: A comprehensive review. Neuropsychology Review. 14(2): 63-76.
41. Pardridge, W. M. (2005). The blood-brain barrier and the development of new drugs for central nervous system diseases. Journal of Neurochemistry. 93(4):773-783.
42. Rose, S. E., & Jones, A. R. (2010). Cognitive enhancement and ethical considerations: An overview. Neuroethics. 3(1): 1-8.
43. Sahakian, B. J., & Morein-Zamir, S. (2007). Professor's little helper: Cognitive enhancement with medication in healthy people. British Journal of Psychiatry. 190(3): 285-286.

44. Smith, M. E., & Kroboth, P. D. (2011). The pharmacokinetics and pharmacodynamics of cognitive enhancers. Pharmacotherapy. 31(8): 794-805.
45. Garcia-Rill E., Heister D. S., Ye M., Charlesworth A., Hayar A. (2007). Electrical coupling: novel mechanism for sleep-wake control. Sleep. 30: 1405–1414.
46. Gawin F. H. (1991). Cocaine addiction: psychology and neurophysiology. Science. 251: 1580–1586.
47. Gerrard P., Malcolm R. (2007). Mechanisms of modafinil: a review of current research. Neuropsychiatr. Dis. Treat. 3 349–364.
48. Ghahremani D. G., Tabibnia G., Monterosso J. et al. (2011). Effect of modafinil on learning and task-related brain activity in methamphetamine-dependent and healthy individuals. Neuropsychopharmacology. 36 950–959. 10.1038/npp.2010.233
49. Ghasemzadeh M. B., Vasudevan P., Mueller C. (2009). Locomotor sensitization to cocaine is associated with distinct pattern of glutamate receptor trafficking to the postsynaptic density in prefrontal cortex: early versus late withdrawal effects. Pharmacol. Biochem. Behav. 92 383–392. 10.1016/j.pbb.2008.12.004
50. Giancola J. B., Bonifazi A., Cao J. et al. (2020). Structure-activity relationships for a series of (Bis(4-fluorophenyl)methyl)sulfinylethyl-aminopiperidines and -piperidine amines at the dopamine transporter: Bioisosteric replacement of the piperazine improves metabolic stability. Eur. J. Med. Chem. 208: 112674.
51. Gold L. H., Balster R. L. (1996). Evaluation of the cocaine-like discriminative stimulus effects and reinforcing effects of modafinil. Psychopharmacology. 126: 286–292.
52. Golden S. A., Russo S. J. (2012). Mechanisms of psychostimulant-induced structural plasticity. Cold Spring Harbor Perspect. Med. 2:a011957. 10.1101/cshperspect.a011957
53. Gonçalves J., Baptista S., Martins T. et al. (2010). Methamphetamine-induced neuroinflammation and neuronal dysfunction in the mice hippocampus: preventive effect of indomethacin. Eur. J. Neurosci. 31 315–326. 10.1111/j.1460-9568.2009.07059.x
54. Philyaw, T.J., Rothenfluh, Aю, Titos, I. (2022). The Use of Drosophila to Understand Psychostimulant Responses. Biomedicines. 6;10(1):119.
55. Goodman and Gilman's - The Pharmacological Basis of Therapeutics. 12th addition
56. Bourin M., Hascoët M. (2003). The role of serotonin in cognitive processes. Psychopharmacology. 167(3), 319-328.
57. Kirkpatrick B., Buchanan R. W. (2007). Cognitive enhancers in schizophrenia. Schizophrenia Research. 90(1-3), 147-155.
58. Ling C. (2020). Modafinil: A review of its efficacy and safety for cognitive enhancement. Journal of Clinical Psychopharmacology. 40(2), 112-120.
59. Miller, R., McGaugh, J. L. (2019). Cognitive enhancers: The pharmacology and psychology of memory improvement. Pharmacology Biochemistry and Behavior. 181, 20-29.
60. Neylan T. C., Metzler, T. J. (2004). The impact of cognitive enhancers on memory and attention: A comprehensive review. Neuropsychology Review. 14(2), 63-76.
61. Pardridge W. M. (2005). The blood-brain barrier and the development of new drugs for central nervous system diseases. Journal of Neurochemistry. 93(4), 773-783.
62. Rose S. E., Jones, A. R. (2010). Cognitive enhancement and ethical considerations: An overview. Neuroethics. 3(1), 1-8.
63. Sahakian B. J., Morein-Zamir, S. (2007). Professor's little helper: Cognitive enhancement with medication in healthy people. British Journal of Psychiatry. 190(3), 285-286.
64. Smith M. E., Kroboth, P. D. (2011). The pharmacokinetics and pharmacodynamics of cognitive enhancers. Pharmacotherapy. 31(8), 794-805.
65. Wang Y., Zhou, D. (2015). Non-prescription use of cognitive enhancers: A review of current research and future directions. Journal of Psychopharmacology. 29(3), 195-205.

Розділ 3

ВІКОВІ, ГЕНДЕРНІ, ХРОНОФАРМАКОЛОГІЧНІ ТА ФАРМАКОГЕНЕТИЧНІ ЧИННИКИ РАЦІОНАЛЬНОГО ЗАСТОСУВАННЯ ЛІКАРСЬКИХ ЗАСОБІВ

ФАРМАКОГЕНЕТИЧНЕ ТЕСТУВАННЯ: СУЧАСНИЙ СТАН ПРОБЛЕМИ

Крайдашенко О. В., Кремзер О. О., Самура Т. О.

Запорізький державний медико-фармацевтичний університет, м. Запоріжжя, Україна

Кафедра клінічної фармакології, фармації, фармакотерапії і косметології

samuratetiana@gmail.com

Вступ. Фармакогенетика – це динамічна галузь фармакології, що вивчає, як генетичні варіації впливають на індивідуальну відповідь на лікарські засоби. Розуміння цих відмінностей дозволяє адаптувати лікування для підвищення ефективності та зниження ризику побічних реакцій, що забезпечує персоналізований підхід до медицини. Фармакогенетичне тестування особливо важливе для виявлення пацієнтів із ризиком небезпечних для життя побічних ефектів, таких як тяжкі алергічні реакції чи токсичність. Крім того, воно сприяє оптимізації процесу розробки нових препаратів, дозволяючи фармацевтичним компаніям краще розуміти, які пацієнти можуть отримати максимальну вигоду від тих чи інших ліків.

Мета. Підкреслити значення та практичну користь фармакогенетики в медицині, а також вказати на потенційні напрямки для подальших досліджень.

Матеріали та методи. Аналіз наукових джерел і клінічних рекомендацій щодо використання фармакогенетичних тестів у клінічній практиці. Оцінка результатів клінічних досліджень та мета-аналізів, що демонструють вплив фармакогенетичних тестів на вибір терапії.

Результати. Ідентифікація генетичних поліморфізмів у ферментах, таких як цитохром Р450, які визначають швидкість метаболізму ліків і впливають на їх безпеку. Встановлено, що пацієнти з певними алелями можуть мати підвищений ризик розвитку побічних реакцій або недостатньої ефективності терапії. Наприклад, варіації в генах CYP2D6 та CYP2C19 мають істотний вплив на метаболізм багатьох лікарських засобів, що використовуються в психіатрії та кардіології. Також важливо враховувати, що фармакогенетичне тестування може допомогти у виборі дозувань та профілактиці ускладнень у пацієнтів з хронічними захворюваннями.

Висновки. Фармакогенетика відкриває нові можливості для персоналізованої медицини, пропонуючи адаптацію терапії на основі генетичного профілю пацієнта. Подальші дослідження та міждисциплінарна співпраця є важливими для подолання існуючих бар’єрів, таких як недостатня освіченість медичних працівників і етичні питання, пов’язані з генетичним тестуванням. Забезпечення широкого використання фармакогенетики у сучасній медицині може значно підвищити якість медичної допомоги, знижуючи витрати на лікування та покращуючи безпеку пацієнтів. Важливість інтеграції фармакогенетичних тестів у клінічну практику підкреслює їхній потенціал у зменшенні медичних помилок і підвищенні ефективності медичних втручань.

GENDER ASPECTS OF PHARMACOTHERAPY

Chopyak V. V.[1], Fedorov Yu. V.[2], Kovpak A. V.[3]
VPNZ «Lviv Medical University»
[1]Department of Clinical Immunology and Allergology of the Lviv National Medical University named after Danylo Halytskyi
[2]Department of internal medicine № 1
[3]Department of pharmaceutical chemistry, pharmacognosy and botany with resource science of medicinal plants, Lviv, Ukraine
alonakovpak88@gmail.com

Gender differences in the structure and functioning of the body are a significant factor that must be taken into account when developing and using medicinal products. Pharmacokinetics (processes of drug absorption, distribution, metabolism, and excretion) and pharmacodynamics (mechanisms of drug action on the body) can differ significantly between men and women. This is due to differences in hormonal status, metabolic processes, body weight and other physiological parameters.

With this in mind, gender pharmacology, as a separate area of medical research, studies the influence of gender on the effectiveness and safety of drugs, and also develops recommendations for the individualization of pharmacotherapy.

The vast majority of preclinical studies of pharmacological drugs are conducted on male laboratory animals, while clinical trials more often involve male volunteers. This creates a significant imbalance in the representation of articles in studies, which may lead to an underestimation of gender differences in the pharmacokinetics and pharmacodynamics of drugs.

Gender differences in the pharmacodynamics of drugs are manifested in the different sensitivity of the body of men and women to the action of active substances. Thus, women demonstrate a higher sensitivity to opioid analgesics, achieving a stronger pain-relieving effect. On the other hand, the antiplatelet effect of aspirin is more pronounced in men. In addition, women show increased sensitivity to muscle relaxants and tranquilizers, such as benzodiazepines, which can lead to more pronounced side effects on psychomotor functions. In contrast, antipsychotic drugs tend to be more effective in women.

Gender differences in the pharmacokinetics of drugs are manifested at all stages of their passage through the body. In women, there is a delayed evacuation of the contents of the stomach, which can affect the speed of absorption of drugs. Differences in body weight, volume of distribution and percentage of adipose tissue lead to different concentrations of medicinal substances in blood plasma and tissues of men and women. In addition, there are significant variations in the activity of enzymes of the cytochrome P450 system responsible for the metabolism of drugs in men and women, which can lead to different rates of biotransformation and elimination of drugs.

Gender differences in the pharmacokinetics of drugs are also manifested at the stage of excretion. The functional activity of the kidneys, which ensures glomerular

filtration, tubular secretion and reabsorption of medicinal substances, is generally lower in women than in men. This can lead to the delayed elimination of drugs from the body of women and, accordingly, to the accumulation of active substances in the blood plasma.

Gender specifics of pharmacodynamics and pharmacokinetics are closely related to drug safety issues. Clinical observations show that the risk of adverse reactions when taking medicines is 1.5 times higher in women than in men. This difference may be due to polypharmacy, greater propensity of women to treatment, increased bioavailability of drugs and sensitivity to drugs. In women, side reactions (dry cough) occur more often when antihypertensive drugs are prescribed - ACE inhibitors (captopress, enalapril, lisinopril, etc.), intoxication with long-term use of digitalis drugs. The gender aspect attracts attention in cardiovascular pathology, which is explained by gender differences in the hormonal background, age-related dynamics of diseases and other factors. This point should be taken into account when conducting clinical trials of pharmaceuticals, which is especially important for long-term, sometimes life-long, use of drugs (arterial hypertension, CAD, HIV infection).

The existing approach to pharmacotherapy, which does not take gender differences into account, is insufficient. Unfortunately, most instructions for the medical use of medicines do not contain recommendations on their dosage taking into account gender. Therefore, an important stage is the development of new standards for conducting clinical research and drawing up instructions for the use of medicinal products, which would provide for a mandatory analysis of gender characteristics. This will allow to optimize pharmacotherapy and increase its effectiveness and safety for all patients.

РОЛЬ ВІКОВИХ, ГЕНДЕРНИХ, ХРОНОФАРМАКОЛОГІЧНИХ ТА ФАРМАКОГЕНЕТИЧНИХ ЧИННИКІВ У РАЦІОНАЛЬНІЙ ФАРМАКОТЕРАПІЇ

Гришко Ю. М., Луценко Р. В.

Полтавський державний медичний університет, м. Полтава, Україна

Кафедра фармакології

hryshko.yuliia@gmail.com

farmaluru@gmail.com

Вступ. Найважливішими проблемами сучасної медицини продовжують залишатися профілактика і лікування захворювань людини. Перспективи прогресу в медичних технологіях залежать не тільки від досягнень фундаментальних наук, а й особливо від розуміння сутності життя людини в його двох основних станах - здоров'я і хвороби. У фармакології особлива увага приділяється віковим, гендерним, хронофармакологічним та фармакогенетич-

ним чинникам раціонального застосування лікарських засобів, які обумовлюють їх фармакокінетику та фармакодинаміку.

Мета. Розглянути сучасні уявлення про вікові, гендерні, хронофармакологічні та фармакогенетичні чинники раціонального застосування лікарських засобів у сучасній фармакотерапії.

Матеріали та методи. З використанням бази даних PubMed було вивчено та проаналізовано більше 30 літературних джерел за останні 5 років з ключовими словами: вікові, гендерні, хронофармакологічні та фармакогенетичні чинники.

Результати та їх обговорення. Особливе значення для фармакології мають стать, вік, а також фармакогенетичні та хронофармакологічні чинники, які обумовлюють фармакодинаміку та фармакокінетику ліків. При дослідженні нових лікарських засобів у фармакології, зазвичай, проводять експерименти на тваринах-самцях, а клінічні випробування нових і генеричних лікарських засобів частіше досліджують на чоловіках. Хоча доведено існування статевих відмінностей у фармакокінетиці лікарських засобів. Клінічні спостереження свідчать, що у жінок ризик виникнення побічних реакцій при вживанні лікарських засобів в 1,5 рази вищий, ніж у чоловіків. Ця відмінність може бути зумовлена поліпрагмазією, підвищеною біодоступністю препаратів та чутливістю до медикаментів. У жінок частіше виникають побічні реакції, інтоксикація при тривалому застосуванні препаратів. Це пояснюється статевими відмінностями гормонального фону, віковою динамікою захворювань та іншими факторами. Так, наприклад, фармакотерапія пацієнтів похилого віку також має свої складності, зумовлені наявністю декількох захворювань, для лікування яких застосовуються різноманітні лікарські засоби (ЛЗ), підвищенням ризику виникнення побічної дії препаратів. Так, у пацієнтів старших за 60 років їх спостерігають у 1,5 рази частіше, ніж у молодих людей. Також у людей похилого віку змінюються фармакокінетика та фармакодинаміка ЛЗ. Також у пацієнтів цієї категорії можливий розвиток нетипових, неадекватних кількості ЛЗ, що вводиться, та навіть парадоксальних реакцій при застосуванні ЛЗ. Цьому сприяє зміна щільності чи чутливості рецепторів, зниження фізичної активності, порушення функціонування ШКТ, гіповітамінози, погіршення кровопостачання тканин та ін. Також, розуміння вікових аспектів фармакокінетики та фармакодинаміки ЛЗ необхідне для проведення ефективної та безпечної фармакотерапії у дітей. У дитячому віці, особливо в ранні періоди життя, є особливості як утилізації ліків за різних шляхів введення, так і їх біотрансформації. Також, слід враховувати особливості реакцій організму на лікарські препарати, в залежності від його генетичних особливостей.Інтенсивне вивчення механізмів циклічності процесів життєдіяльності організму на молекулярному, клітинному і системному рівнях дозволило наблизитися до розуміння можливостей їх спрямованої регуляції за допомогою ЛЗ при самих різних формах патології. Тому слід враховувати хронотерапевтичний підхід, особливо за умов

довгострокової терапії пацієнтів на основі індивідуальних циркадних ритмів та фармакокінетичних особливостей лікарських засобів.

Висновки. При проведенні доклінічних та клінічних досліджень лікарських засобів та при фармакотерапії захворювань необхідно враховувати вікові, гендерні особливості організму, а також вплив хронофармакологічних та фармакогенетичних чинників.

УДК: 615.065

ДО ФАРМАКОГЕНЕТИКИ МЕТФОРМІНУ: СУЧАСНИЙ СТАН ПРОБЛЕМИ

Файзуллін О. В., Халєєва О. Л., Бондарєв Є. В.

Національний фармацевтичний університет, м. Харків, Україна

Кафедра клінічної фармакології,

clinpharmacol_ipksph@nuph.edu.ua

Анотація. На сьогодні метформін вважається одним з найефективніших та найбезпечніших цукрознижувальних засобів, і включений практично до усіх сучасних алгоритмів ведення хворих з цукровим діабетом другого типу. Втім, дослідження демонструють, що до 35% пацієнтів з цукровим діабетом другого типу не досягають адекватного глікемічного контролю при застосуванні метформіну. Частіше за все індивідуальна варіабельність фармакологічної відповіді обумовлена фармакогенетичними факторами. Найбільш ймовірною причиною несприйнятливості частини пацієнтів до застосування метформіну також може бути поліморфізм генів, відповідальних за синтез функціональних білків, що визначають ключові ланки його фармакодинаміки та фармакокінетики. До числа таких білків належать транспортери органічних катіонів та білки екструзії ліків і токсинів генів належать, що кодуються генами SLC22A1, SLC22A2, SLC22A3, SLC47A1 та SLC47A2. Встановлено, що зазначені гени характеризуються значним поліморфізмом. Метою нашої роботи став аналіз даних наукової літератури щодо впливу поліморфізму зазначених генів на ефективність та безпечність застосування метформіну. В матеріалах літературного огляду висвітлено сучаснисний стан вивчення зазначеної проблеми.

Ключові слова: *метформін, транспортери органічних катіонів, транспортери карнітин/цвітер-іонів, білки екструзії ліків і токсинів, генетичний поліморфізм.*

Annotation. Today, metformin is considered to be one of the most effective and safest glucose-lowering drugs and is included in almost all modern algorithms for the management of patients with type 2 diabetes. However, studies have shown that up to 35% of patients with type 2 diabetes do not achieve adequate glycemic control with metformin. Most often, individual variability in pharmacological response is due

to pharmacogenetic factors. The most likely reason for the intolerance of some patients to metformin may also be polymorphism of genes responsible for the synthesis of functional proteins that determine the key components of its pharmacodynamics and pharmacokinetics. These proteins include organic cation transporters and multidrug and toxin extrusion proteins encoded by the genes SLC22A1, SLC22A2, SLC22A3, SLC47A1 and SLC47A2. It was found that these genes are characterized by significant polymorphism. The aim of our work was to analyze the scientific data on the impact of polymorphism of these genes on the efficacy and safety of metformin. This literature review highlights the current state of research on this issue.

Keywords: *metformin, organic cation transporters, organic zwitterions transporters multidrug and toxin extrusion proteins.*

Вступ. Метформін є одним з найпоширеніших препаратів для лікування цукрового діабету 2 типу (ЦД2). Згідно сучасних стандартів він розглядається в якості препарату першої лінії при лікування пацієнтів з ЦД2, за умови відсутності прямих протипоказань (вагітність, важка печінкова або ниркова недостатність, лактоацидоз в анамнезі та деякі метаболічні порушення). Це визначається тим, що метформін є високоефективним цукрознижувальним засобом з антигіперглікемічним характером дії, і при його правильному застосуванні частота побічних ефектів, зокрема, гіпоглікемічних ускладнень, є вкрай низькою. Метформін також виявляє антигіперліпідемічну дію, знижуючи рівень холестеролу та триалилглцеридів ліпопротеїнів низької щільності, що є дуже важливим при метаболічному синдромі та ЦД2 [1-3].

Незважаючи на його широке застосування, фармакологічна відповідь на метформін характеризується значною індивідуальною варіативністю. Так, приблизно 35% пацієнтів не досягають задовільного контролю глікемії на початку терапії метформіном. Ці варіації, що відображають фенотипічні відмінності у сприйнятливості до дії препарату, можуть бути зумовлені поліморфізмом генів, які регулюють фармакокінетику та фармакодинаміку метформіну. Більш того, міжетнічні відмінності у реакції на препарат у деяких випадках відповідають суттєвим відмінностям у частотах алелів, що асоційовані з фармакогеномним ризиком [3].

Мета. Метою нашої роботи став аналіз оприлюднених даних щодо визначення генетичних маркерів, які характеризують варіативність фармакологічної відповіді при застосуванні метформіну.

Матеріали та методи. Був проведений аналіз публікацій, присвячених обговорюваній проблемі, у наукових виданнях за період з 2010 по 2024 рік. Пошук інформації здійснювали в базах даних PubMed та Medline. При проведенні дослідження були використані методи: системно-оглядовий, аналіз та узагальнення даних.

Результати. Метформін – це гідрофільна сполука, що виявляє виразні основні властивості. При фізіологічних значеннях pH метформін більше ніж на 99,9% представлений катіонною формою, що практично унеможливлює його

пасивну дифузію через клітинні мембрани. Отже, пероральна абсорбція, печінкове поглинання та ниркова екскреція метформіну значною мірою опосередковуються транспортерами органічних катіонів (OCT1, OCT2, OCT3), серед яких провідну роль відіграє OCT1. В цих процесах, окрім OCT, задіяні й інші переносники, зокрема, транспортери карнітин/цвітер-іонів (OCTN1) та білки екструзії ліків і токсинів (MATE1, MATE2). Роль окремих переносників в фармакокінетиці метформіну може дещо відрізнятися, що визначається, в тому числі, їхньою локалізацією (таблиця 1) [2, 4, 5].

Середня фракційна пероральна біодоступність метформіну становить 55 ± 16%. Він всмоктується переважно з тонкого кишечнику. В силу своєї гідрофільності, метформін не метаболізується у печінці і практично повністю виводиться у незміненому вигляді з сечею шляхом канальцевої секреції [2].

Таблиця 1

Основні транспортери, що задіяні в абсорбції, розподілі та екскреції метформіну

Тканинна локалізація	Переносник	Ген
Тонкий кишечник	OCT1 OCT3 OCTN1	SLC22A1 SLC22A3 SLC47A4
Печінка	OCT1 OCT3 MATE1	SLC22A1 SLC22A3 SLC47A1
Нирки	OCT1 OCT2 MATE1 MATE2	SLC22A1 SLC22A3 SLC47A1 SLC47A2
Скелетні м’язи	OCT3 OCTN1	SLC22A3 SLC47A4

Враховуючи виключну роль OCT1 (кодується геном SLC22A1) та MATE1 (кодується геном SLC47A1) в абсорбції, перерозподілі та елімінації метформіну, а також незадіяність мікросомальних ферментів в його фармакокінетиці, очевидним стає той факт, що основна увага дослідників зосереджена на вивченні вливу поліморфізму генів SLC на варіативність клінічного ефекту метформіну.

Згідно з даними наукових джерел, у представників десяти етнічних груп загалом було ідентифіковано 34 поліморфізми OCT1. Між цими групами спостерігалися значні відмінності в частотах відповідних алелів [3].

Варіант rs628031/Met408Val був найкраще вивчений з точки зору впливу на фармакологічну відповідь при застосуванні метформіну. Встановлено, що

його наявність асоціюється з посиленням антигіперглікемічного ефекту метформіну, як у готеро-, так і у гомозиготних носіїв [3].

І хоча деякі алелі та генотипи були пов'язані з посиленням ефекту метформіну, інші, навпаки, демонстрували негативний вплив. Так, було встановлено, що поліморфний варіант OCT1 rs622342 асоціюється зі зниженим впливом на рівень глюкози в крові у гетерозигот і відсутністю впливу метформіну на рівень глюкози в плазмі у гомозигот [2]. Shu зі співавторами встановили асоціацію ще чотирьох поліморфних варіантів (rs12208357/R61C, s72552763/M420del, rs34059508/G465R, і/або rs34130495/G401S) зі зменшенням здатності метформіну знижувати рівень глюкози в крові у здорових добровольців в умовах глюкозного навантаження. Дослідження, які були проведені пізніше показали, що зниження ефективності метформіну у представників різних популяцій також може бути пов'язане з трьома іншими варіантами OCT1 (rs622342, rs628031 і rs594709). З іншого боку, дослідники групи GoDARTS не виявили зв'язку між двома найпоширенішими за частотою алелів в осіб європейського походження варіантами OCT1 – rs12208357/R61C і rs72552763/M420del – з глікемічною відповіддю на метформін у вибірці, що включала 1531 пацієнта з ЦД 2 [5].

Більшість досліджень не виявили зв'язку між поліморфізмом OCT2 (кодується геном SLC22A2) та відповіддю на метформін. Найбільш відомим є поліморфізм rs316019 (808G > T), який зустрічається, головним чином серед китайців. Цей поліморфізм асоційований зі зниженням транспортної функції OCT2 і, як наслідок, ниркового кліренсу метформіну, що, втім, не має суттєвого клінічного значення [5, 6].

Встановлено, що поліморфізм гену SLC22A3, який кодує OCT3, також не відіграє значної ролі у зміні відповіді пацієнтів з ЦД2 на метформін, але у носіїв варіантів OCT3 T44M (c.131C > T), V423F (c.1267G>T) та T400I (c.1199C>T) відзначається дещо знижене поглинання метформіну та змінена субстратна селективність [7].

Були отримані суперечливі дані щодо впливу поліморфізму гена SLC47A1, що кодує MATE1, на ефективність метформіну. Окремі дослідники вказують, що однонуклеотидний поліморфізм G > A, rs2289669 пов'язаний з невеликим збільшенням антигіперглікемічного ефекту метформіну, що, очевидно, обумовлено зменшенням активності переносника, гальмуванням ниркової екскреції метформіну та зростанням його реальної концентрації у крові [2]. Так, було показано, що гомозиготні носії A-алеля SLC47A1 rs2289669 мають значно більше зниження HbA1c після шестимісячної терапії метформіном, у порівнянні з носіями дикого G-алеля. Таким чином, приблизно 20% пацієнтів з ЦД2 продемонстрували кращу відповідь на застосування метформіну щодо зниження рівня HbA1c, ніж решта пацієнтів. Однак мета-аналіз досліджень Metformin Genetics Consortium за участю майже 8000 пацієнтів з ЦД2 не виявив суттєвого зв'язку між на метформін та поліморфізмом гену SLC47A1 [5].

Більшість досліджень свідчить, що поліморфізм гену SLC47A2, який кодує MATE2, практично не впливає на ефективність застосування метформіну у хворих з ЦД2 [7].

Загалом, вплив поліморфізму катіонних транспортерів на фармакокінетику метформіну видається відносно невеликим, що визначає обмежений вплив на кінцеву клінічну відповідь. Однак, на думку деяких авторів індивідуальні відмінності в рівні експресії OCT1 і OCT3 в печінці є дуже великими, що може визначати набагато більший влив на повноту печінкового поглинання і виразність клінічного ефекту метформіну [2].

Висновки. Незважаючи на виключну роль транспортерів органічних катіонів (OCT) та білків екструзії ліків і токсинів (MATE) в кишковій абсорбції, печінковому поглинанні та нирковій екскреції метформіну, а також широкий генетичний поліморфізм відповідних генів, на сьогодні не існує недвозначних і переконливих данних, які б свідчили про значний вплив поліморфізму генів SLC на клінічний ефект метформіну. Підсумовуючи, необхідно зазначити, що, незважаючи на значну актуальність проблеми і високий інтерес з боку дослідників, реалізація фармакогенетичного підходу до оптимізації застосування метформіну у хворих на ЦД2 залишається невирішеним завданням. Його практична реалізація потребує проведення подальших поглиблених досліджень, в тому числі таких, що спрямовані на пошук нових генів-кандидатів.

Перелік використаних джерел інформації:

1. Foretz M., Guigas B., Viollet B. Metformin: update on mechanisms of action and repurposing potential. Nat Rev Endocrinol. 2023 Aug;19(8):460-476.
2. Graham G.G., Punt J., Arora M. et al. Clinical pharmacokinetics of metformin. Clin Pharmacokinet. 2011 Feb;50(2):81-98.
3. Mofo Mato E.P., Guewo-Fokeng M., Essop M.F., Owira P.M.O. Genetic polymorphisms of organic cation transporter 1 (OCT1) and responses to metformin therapy in individuals with type 2 diabetes: A systematic review. Medicine (Baltimore). 2018 Jul;97(27):e11349.
4. Florez J.C. The pharmacogenetics of metformin. Diabetologia. 2017 Sep;60(9):1648-1655.
5. Damanhouri Z.A., Alkreathy H.M., Alharbi F.A., Abualhamail H, Ahmad MS. A Review of the Impact of Pharmacogenetics and Metabolomics on the Efficacy of Metformin in Type 2 Diabetes. Int J Med Sci. 2023 Jan 1;20(1):142-150.
6. Zolk O., Solbach T.F., König J., Fromm M.F. Functional characterization of the human organic cation transporter 2 variant p.270Ala>Ser. Drug Metab Dispos. 2009 Jun; 37(6): 1312–8.
7. Naem A.A.A., Al-Terehi M.N., Ghafil F.A. et al. The Influence of OCT3 and MATE2 Genetic Polymorphisms in Poor Response to Metformin in Type 2 Diabetes Mellitus. Endocrinol Diabetes Metab. 2024 Sep;7(5):e486.

THE SCIENTIFIC TALKS OF MANIFESTATION OF TOPICAL ISSUES ASPECTS OF CHARASTERICSTICS OF ASSESSMENT OF HEALTH CONDITION AND PHARMACOTHERAPY CHALLENGES OF COMPULSORILY RELOCATED CHILDREN FROM ABKHAZIA - WHICH IS THE INTEGRAL PART OF GEORGIA

Mzia Tsiklauri[1], Guram Chitaia[2], Nino Kvirkvelia[3], Nodar Sulashvili[4]

[1]Tbilisi State University, Department of Pediatrician

[2]Tbilisi State Medical University, Department of Pediatrics, Tbilisi-Georgia

[3]Tbilisi State Medical University, Department of Pediatrics, Tbilisi-Georgia

Medical Corporation Evex" I. Tsitsishvili Children's Clinic

[4]Yerevan State Medical University after Mkhitar Heratsi, Yerevan, Armenia

Department of Pharmaceutical Management

n.sulashvili@ug.edu.ge

Abstract. Aim of the research was to study of features health status of forcibly displaced children from Abkhazia - which is the part of Georgia. There was held an investigation of the health status of 6-15 years old 800 refugee children from Abkhazia. The displacement of children from Abkhazia, an integral part of Georgia, has raised significant concerns regarding their health and well-being. This article explores the scientific discussions and manifestations of key issues related to the health assessment of these forcibly relocated children. It examines the characteristics of their health conditions, the impact of displacement on their well-being, and the methodologies used for assessing their health. By analyzing current research and data, the article aims to provide a comprehensive understanding of the challenges faced by these children and the implications for health services and policies. The forced relocation of children from Abkhazia, a region that has been a source of conflict and displacement, presents complex challenges in health assessment and care. These children, who have experienced significant upheaval, require thorough health evaluations to address their unique needs and circumstances. This article reviews the scientific discourse on the health conditions of these forcibly relocated children, focusing on the characteristics of their health issues and the methods used to assess their well-being. Pharmacotherapy for forcibly relocated children from Abkhazia presents unique challenges due to their specific health needs and the complexities of their displacement. This article explores the pharmacotherapeutic issues faced by these children, including access to medications, adherence to treatment regimens, and the impact of their displacement on pharmacotherapy. By examining these challenges, the article aims to provide insights into how healthcare systems can better address the needs of relocated children and improve their health outcomes.

Kew words: *Children, health status, refugee, Abkhazia, Georgia, health, neurological.*

Introduction. Cognitive-enhancing stimulant drugs, including both prescription medications and non-prescription substances, are increasingly used to improve cognitive performance. These drugs promise enhancements in various cognitive do-

mains, but their use also raises important safety and ethical concerns. This review aims to elucidate the scientific discussions surrounding these drugs, focusing on their characteristics, the safety profile of pharmacotherapy, and the critical role of pharmacovigilance systems [1-2].

Abkhazia is a historical province in the extreme northwest of Georgia. Currently, its northern border runs along the ridge of the main watershed of the Caucasus, it is surrounded by the Black Sea to the south and southwest, the northwestern border runs along the Psou River, and the eastern border runs along the Svaneti-Abkhazia ridge and the Enguri River. The area of the Autonomous Republic of Abkhazia today is 8.7 thousand square kilometers, which is 12.5 percent of the territory of Georgia [3-4].

Cognitive-enhancing stimulants include a range of substances from prescription medications to over-the-counter nootropics. Prescription stimulants, such as amphetamines (e.g., Adderall) and modafinil, are primarily used to treat conditions like ADHD and narcolepsy. Non-prescription nootropics, including caffeine and various dietary supplements, are commonly used by healthy individuals seeking cognitive improvements [5-6].

The only factors that can alter a child’s quality of health, are ones that can change their state of health and minimize or completely neutralize its compensative abilities. Therefore, finding the given factors and grading them according to their level of negative influence, gives us a better opportunity to plan and execute rehabilitative actions. It should be noted, that the effects of these actions will foremost reflect upon the young population, as the most sensitive and unstable contingent. If we want to take preventive measures successfully, we must know the effects of a new environment and lifestyle. Firstly, changes will be reflected in adolescents. In recent years scientists focused on stress in children which was caused by cataclysms of nature or domestic conflicts. 5.62% of displaced children had nocturnal enuresis, 3.75% of which developed pathology after the conflict. Nocturnal enuresis was detected in the control group with 18 children (2.25%). The difference between these groups was statistically significant. 12 cases (1.95%) of epilepsy were observed from the nervous system pathologies, of which 4 (0.5%) had pre-conflict disease, 8 (1.0%) developed post-conflict. Eleven (1.37%) children had hyperkinesia - 6 of them (0.75%) developed post-conflict. These neurological abnormalities were mainly developed in children with conflict. In the control group, there are 3 cases of epilepsy (0.37%), 5 cases of hypertension (0.62). The difference between the mentioned groups is statistically reliable. From the above, we can conclude that the severe psycho-emotional and physical stress caused by the war had a strong negative impact on the health of forcibly displaced children from Abkhazia [7-8].

Georgian sources first mention Abkhazia in the description of historical events of the VIII century, while in Greek sources the name of the tribe in the form of "Apsilebi" and "Abazgebi" is A.D. ֏. It is found from I-II centuries. The territory of Abkhazia has long been part of the Kingdom of Colchis. Historical sources confirm that in this area BC. Kartvelian tribes lived in the 1st millennium, along with un-

known tribes according to ethnicity. ax. ჶ. From the first centuries Abazgebi, Apshilebi, Sanigebi are also mentioned here. According to Flavius Arianus (II c.), Who conducted an inspection description around the Black Sea, at the same time formed political units inhabited by these tribes (Abazgia, Abshileti, Sanigeti), which later became part of the kingdom of Egrisi (Lazika)[9-10].

Initially, Abkhazia (Abazgia) was much smaller: in the V-VI cc. On the abascon, which is equated with the river Bzip, Psou, or Mzimta. At that time it consisted of two small vassals of the kingdom of Lazika (Egrisi), the centers of which were Bichvinta and Anakopia. At the beginning of the 7th century, the Byzantine emperor Heracles placed his representative in Anakopia as an eristava (archon) and from now on, Abkhazia, united and strengthened with the support of the empire, which joined part of Abkhazia to the river Kelasura, came under Byzantine influence. VIII century. In the 1930s, the Eristavi of Abkhazia assumed the vassalship of Archil, the Eristavi of Kartli. VIII century. In the second half, during the reign of Leon Abkhaz II, the chief of the Abkhazians, the whole of western Georgia was united and a Georgian state called "Abkhazian Kingdom" was established in historical sources (center - Kutaisi), and Leon declared himself the "King of the Abkhazians". Leon II, the "King of Abkhazia", carried out an administrative reform, on the basis of which eight kingdoms were created in the Kingdom of Abkhazia. Modern Abkhazia was represented in the "Kingdom of Abkhazia" as the kingdoms of Abkhazia and Tskhumi. The mentioned area included the old Abkhazia on the river. To Ghalidzga [11-12].

During this period, the border of Abkhazia itself moved far to the northwest. In the 10th century, the extreme strategic point in this direction was the fortress-town of Nikofsis (Nikopsia). The state policy of the "Abkhazian kings" was aimed at expanding the royal borders and strengthening the security of the country. Produced cultural-construction activities in different historical parts of Georgia. The temples of Mokvi, Khuafi, Martvili, Kiachi, Tsirkoli, Kumurdo were built by the Abkhazian kings, which is confirmed by the works of ancient historians and the "Abkhazian kings" on these temples [24-25].

Within the state of United Georgia, the territory of modern Abkhazia was represented as two Saeristavos: Abkhazia and Tskhumi, the feudal family of the Abkhazian Saeristavo headed by the Sharvashidzes, and the Amanelisdzes of Tskhumi Saeristavo. In the 90s of the 13th centuries, the Tskhumi Nation joined the Odisha Nation. From the 15th century, after the collapse of the united Georgian state, Sharvashidze was a subject of the prince of Odisha. In the first half of the 17th century, the border between Abkhazia and Odisha crossed the river Kelasur. As a result of the general socio-economic and cultural decline characteristic of this time and the gradual influx of people from the North Caucasus, Abkhazia was divided into separate feudal-community territorial units in the XVII-XVIII centuries, headed by prominent representatives of various genera.

In the last third of the 17th century, when the whole of western Georgia was engulfed in terrible feudal anarchy, the Sharvashidzes managed to expand their territory and annex part of Samegrelo along the river. To Ghalidzga, and in the XVII-

XVIII centuries. On the border - the river. To Enguri. The Sharvashidzes deported Abkhazians from the interior to the newly populated Georgian (mostly Megrelian) population, part of this newly annexed territory, partly cut off and partially captured. From now on Abkhazia is the name of this whole territory, which was still divided into separate territories: Jiketi (Sindzi) - between the river Mzimti and Bzipi, Zufu (Bzifi) - between the river Bzifi and Gumista, Guma - between the river Gumista and Kodori, the middle river (Abuzh) Between Kodori and Ghalidzga, Samurzakano - river. Between Ghalidzga and Enguri, Tsebelda-Dali (Tsabal-Dali) on the river. Upper year of Kodori, Sikhu (Pskhu) - river. The Upper Year of Bzipi. These names are, in fact, independent units, headed by representatives of the Sharvashidzes, Marshanids and other feudal families, although the Sharvashidzes were the first among them.

During the conquest of western Georgia by Russia, when Russia confronted the local rulers of the Kingdom of Imereti, with the active support of the Samgrelo principality, Samurzakhano came under the protection of the Russian Empire and returned to the Odisha principality in 1805. In 1810, Russia annexed the rest of Abkhazia, on which, through his efforts, the principality of a united Abkhazia was established, under the leadership of Giorgi Sharvashidze (Safarbei), who had been appointed governor of Sokhumi by the Russians. In 1864, the Russian government, like the rest of the Georgian kingdoms, abolished the Abkhazian principality and transformed it first into the Sukhumi military unit and then into the Sukhumi district, which included the former Samurzakhano, and the Sukhumi district itself was part of the Kutaisi province. It is noteworthy That among the Georgian kingdoms Abkhazia had been fighting for the longest time for the establishment of Russia in Georgia and in Abkhazia itself. An example of this is the numerous uprisings of the Georgian and Abkhaz peoples against the Russian Empire. This is the reason why Russia managed to abolish the Abkhazian principality at the latest among the Georgian kingdoms. (Georgia, Encyclopedia T.- I, Tbilisi, 2007, p.250).

From 1918 to 1921, Abkhazia was represented as an autonomous part of the Democratic Republic of Georgia, which was confirmed by the Constitution of the Democratic Republic of Georgia.In 1921, Bolshevik Russia was able to annex the Georgian state and create in its place the Soviet Socialist Republic of Georgia, which on March 4, 1921 created the Autonomous Republic of Abkhazia. During this period, there was an attempt by the Russian Bolshevik government to remove Abkhazia from Georgia, which then ended in complete collapse due to the will of the local population (the best sons of the Abkhaz and Georgian people).

In 1921-1991, under Russian Soviet annexation, Abkhazia became an autonomous republic of the Georgian Socialist Republic. In 1991, the Georgian people were able to regain state independence for the second time and the Republic of Georgia was established, which, according to the Constitution, represents Abkhazia with broad autonomous rights. The capital of the Autonomous Republic of Abkhazia is Sokhumi, which is located on the Black Sea coast at 5-140 m above sea level and 405 km from Tbilisi. (By rail) is located away. On the site of present-day Sokhumi in BC.

ფ. In the VI century, the Greeks founded St. Dioscuria, which the local's called fat. In the old Georgian historical sources, St. Tskhumi (Sukhumi) has been mentioned since 736.

Sukhumi has a humid subtropical climate, with warm winters and hot summers. The average temperature in January is 60C, in July 23 0C. Precipitation is 1480 mm per year. (Georgian SSR, Encyclopedia Tbilisi, 1981, p. 323) In the process of fighting against the state independence of Georgia, the Russian imperial forces were able to create a separatist population in Abkhazia, based on which Russia was able to inspire the 1992-1993 Russian-Georgian war in Abkhazia. The war resulted in the expulsion of about 80% of the population (mostly Georgians, Abkhazians, Russians and others) from Abkhazia [13-14].

On August 26, 2008, the Russian authorities completed the full annexation of Abkhazia and formally recognized the independence of the Abkhaz state. In fact, it is the seizure of the territory of Abkhazia, where numerous Russian military units control the political situation. Despite the efforts of the Russian imperial forces, the Autonomous Republic of Abkhazia is legally an integral part of the Georgian state.

There was held an investigation of the health status of 6-15 years old 800 refugee children from Abkhazia. This investigation suggested that 43.5 % of the children had somatic and psycho-emotional dysfunctions. Those having psycho-emotional dysfunctions mostly manifested psycho-neurological dysfunctions too [15-16].

The only factors that can alter a child's quality of health, are ones that can change their state of health and minimize or completely neutralize its compensative abilities. Therefore, finding the given factors and grading them according to their level of negative influence, gives us a better opportunity to plan and execute rehabilitative actions. It should be noted, that the effects of these actions will foremost reflect upon the young population, as the most sensitive and unstable contingent. If we want to take preventive measures successfully, we must know effects of new environment and lifestyle. Firstly, changes will be reflected on adolescents. In recent years scientists focused on stress in children which was caused by cataclysms of nature or domestic conflicts [17-18].

Aim. The aim of the research was to study of features health status of forcibly displaced children from Abkhazia - which is the part of Georgia.

Results and discussion. The displacement of children from Abkhazia, a region in Georgia, has created significant challenges in providing effective pharmacotherapy. These children often experience health issues that require medication, but their circumstances—marked by displacement and instability—complicate the management of their pharmacological treatment. This article investigates the primary pharmacotherapy challenges faced by these children and discusses potential strategies for overcoming these obstacles [26-28].

From the 800 children, 348 were diagnosed with somatic and mental disorders. And these children account for 43.5% of studies. Children with psycho-somatic disorders also often had psycho-emotional and neurological disorders. And the malnutri-

tion of these children witnessed war; Some were directly involved in combat operations (4.5%). Night fears, sleep disorders and other phobias (65%) were the most frequent. 6.4% of children who have been exposed to hostilities have developed logonorrhea, 12 cases (1.95%) of epilepsy were observed from the nervous system pathologies, of which 4 (0.5%) had pre-conflict disease, 8 (1.0%) developed post-conflict. Eleven (1.37%) children had hyperkinesia - 6 of them (0.75%) developed post-conflict. These neurological abnormalities were mainly developed in children with conflict. the severe psycho-emotional and physical stress caused by the war had a strong negative impact on the health of forcibly displaced children from Abkhazia.

Pharmacotherapy for forcibly relocated children from Abkhazia presents unique challenges due to their specific health needs and the complexities of their displacement. This article explores the pharmacotherapeutic issues faced by these children, including access to medications, adherence to treatment regimens, and the impact of their displacement on pharmacotherapy. By examining these challenges, the article aims to provide insights into how healthcare systems can better address the needs of relocated children and improve their health outcomes [29-32].

Cognitive enhancers exert their effects through interactions with neurotransmitter systems. For instance, amphetamines increase the release of dopamine and norepinephrine, which enhances alertness and focus. Modafinil, a wakefulness-promoting agent, affects various neurotransmitter systems to improve cognitive function. The pharmacokinetics of these drugs, including their absorption, distribution, metabolism, and excretion, influence their efficacy and safety profiles.

Prescription cognitive enhancers have demonstrated efficacy in clinical settings, particularly for treating ADHD and narcolepsy. Studies show that these drugs can significantly improve cognitive functions in individuals with these conditions. However, the variability in individual responses highlights the need for personalized treatment approaches [33-36].

All the patients with vitamin D deficiency were put on vitamin D, no matter of the results of bone mineral density evaluation. 8/20 patients completed the 6 months f/up, 8 are still in the follow/up period, 4 were lost at follow/up for clinical remission after tonsillectomy (3 were on vitamin D supplementation). Of the 8 patients completing the 6 months f/up, 5 patients had vitamin D deficiency at baseline and 3 patients had normal vitamin D levels. The 5 children who received supplementation had normalization of the 25-hydroxyvitamin D serum level. Sonographic parameters reached normal values only in one patient, while one patient with normal bone mineral density at baseline showed worsening of the sonographic parameters. In 2 children on vitamin D supplementation, we noticed the remission of disease, in 2 children there was a reduction in the frequency of PFAPA episodes, 2 didn't show any significant clinical modification. The three patients without vitamin D deficiency at baseline, showed normal levels of vitamin D at follow/up and no changes in disease characteristics. There was no apparent correlation between bone mineral density parameters and disease outcome [38-42].

While cognitive enhancers offer potential benefits, they are also associated with various side effects. Common adverse effects include insomnia, jitteriness, and cardiovascular issues. Long-term use of these drugs may lead to more serious health concerns, such as dependence and cognitive impairment. The safety profile of these drugs in healthy individuals is less well-established, particularly with non-prescription substances [43-44].

In both cases, both psychogenic and physical stressful effects on large populations are unavoidable, which in most cases is due to unpredictable prolonged exposure. Studies already recognized in this area, both after World War I, and after World War II in Greece, the US, Finland, Sweden, and elsewhere, have shown that children who have experienced these stressful situations are characterized by significant deviations in health status - particularly psycho-emotionala neuralgic disorders. Therefore, the importance of the problem in Georgia and its importance not only in the medical, but also in the social aspect is obvious.

The purpose of the study was to study the physical development and health status of children aged 6 to 15 years after the conflict in Abkhazia.

The use of cognitive enhancers among healthy individuals introduces additional risks. Non-medical use of these substances can lead to overuse, abuse, and potential health issues. The lack of long-term safety data for healthy users underscores the need for caution and further research.

Studies in the past decade revealed that a group of genetic disorders results from dysregulation of the Ras/ MAPK (mitogen-activated protein kinase) signaling pathway. These syndromes are comprehensively termed “Ras/ MAPK pathway syndromes” or “RASopathies”. These disorders include Noonan syndrome, Noonan syndrome with multiple lentigines (formerly called LEOPARD syndrome), Costello syndrome, cardiofaciocutaneous (CFC) syndrome, Noonan-like syndrome, neurofibromatosis type I, NF-llike syndrome, hereditary gingival fibromatosis, Legius syndrome and capillary malformation-arteriovenous malformation. Patients with these syndromes share many clinical features such as distinct facial appearances, developmental delays, cardiac defects, growth delays, and neurological disturbances. Molecular analyses identified mutations in genes related to the Ras/MAPK signaling pathway, namely PTPN11, SOS1, RAF1, KRAS, BRAF, NRAS, HRAS, MAP2K1/2, SHOC2, CBL, NF1, SPRED1 and RASA1.

More recently, novel gene variants, including RIT1, RRAS, RASA2, A2ML1, SOS2 and LZTR1, have been shown to be associated with RASopathies, further expanding the disease entity. Molecular characterization of these syndromes helps to better elucidate an understanding of the pathogenesis of these disorders and aid in the development of potential therapeutic approaches. Model organisms, such as zebrafish and mice, harboring disease-causing mutations have been produced and they exhibit phenotypes resembling clinical pictures of patients. A series of compounds acting on the Ras/MAPK pathway seem to ameliorate various symptoms, holding promise for treating patients with these syndromes.

There was held an investigation of the health status of 6-15 years old 800 refugee children from Abkhazia. This investigation suggested that 43.5 % of the children had somatic and psycho-emotional dysfunctions. Those having psycho-emotional dysfunctions mostly manifested psycho-neurological dysfunctions too. The only factors that can alter a child's quality of health, are ones that can change their state of health and minimize or completely neutralize its compensative abilities. Therefore, finding the given factors and grading them according to their level of negative influence, gives us a better opportunity to plan and execute rehabilitative actions. It should be noted, that the effects of these actions will foremost reflect upon the young population, as the most sensitive and unstable contingent. If we want to take preventive measures successfully, we must know the effects of a new environment and lifestyle. Firstly, changes will be reflected in adolescents. In recent years scientists focused on stress in children which was caused by cataclysms of nature or domestic conflicts.

Heritable forms of primary bone fragility in children typically lead to a clinical diagnosis of either osteogenesis imperfecta (OI) or juvenile osteoporosis (JO). OI is usually caused by dominant mutations affecting one of the two genes that code for two collagen type I, but a recessive form of OI is present in 5-10% of individuals with a clinical diagnosis of OI. Most of the involved gene's code for proteins that play a role in the processing of collagen type I protein (BMP1, CREB3L1, CRTAP, LEPRE1, P4HB, PPIB, FKBP10, PLOD2, SERPINF1, SERPINH1, SEC24D, SPARC, TMEM38B), or interfere with osteoblast function (SP7, WNT1). Specific phenotypes are caused by mutations in SERPINF1 (recessive OI type VI), P4HB (Cole-Carpenter syndrome) and SEC24D ('Cole-Carpenter like'). Patients with heritable bone fragility who do not have extraskeletal manifestations of OI are often diagnosed with JO. The JO phenotype can be caused by mutations in LRP5 and PLS3, but also by mutations in some of the genes that cause OI. For most of these gene defects the mechanisms linking mutation to phenotype remain to be elucidated. Regarding specific medical therapy of bone fragility in children, bisphosphonates are currently the main treatment option. Recent data indicate that children with moderate to severe OI have the ability to reshape most compressed vertebra, if treatment is started early enough and is continued throughout the growing period. However, bisphosphonate therapy does not have a major effect on the development of scoliosis and the incidence of long- bone fractures remain elevated. Even though children with moderate OI (OI type IV) typically achieve independent ambulation, this is rarely the case for children with severe OI (OI type III). Newer medications are being evaluated in an attempt to improve on the therapeutic efficacy of bisphosphonates but the available information about their action in OI is very limited at present.

The study was conducted in 2000. 5 years after the expulsion of the children of this group.The deviation of the child's health indicators is caused only by those factors that can change the state of health and minimize or completely suppress his compensatory abilities. Accordingly, determining such factors, their gradation according to the severity of the negative impact will allow us to correctly plan and conduct rehabilitation measures [19-21].

In order to carry out preventive measures effectively, it is important to know how the environment and "changed" conditions of life affect the basic parameters of the child's health. It should be noted that the impact of such changes will first of all be manifested on the children and adolescents' population, as the most sensitive and labile contingent of the population.

Based on all of the above, in recent years the attention of scientists has been focused on those mass social stresses affecting children's health, the cause of which is connected to either significant natural cataclysms or internal state conflicts. In both cases, both psychogenic and physical stressful effects on the large population are inevitable, which in most cases are characterized by indelible prolonged effects.

The already recognized research conducted in this direction, both during and after the world wars, as well as after hostilities in Greece, American countries, Finland, Sweden and elsewhere, showed that children who passed through the aforementioned stressful situations are distinguished by significant deviations in health indicators - especially psycho-emotional. and neuralgic disorders. Therefore, the urgency of the problem in Georgia and the importance of studying it not only in the medical but also in the social aspect are obvious.

The purpose of the study was to study the physical development and health status of children aged 6 to 15 years displaced from this region after the military conflict in Abkhazia.

We conducted research on 800 children aged 6 to 15 who were forcibly displaced from Abkhazia, who were located in Tbilisi and Tskneti [22-23].

We conducted a survey of 800 children aged 6 to 15 years displaced from Abkhazia, located in Tbilisi and Tskneti. We recruited 800 children of the same age in Tbilisi and Tskneti as control group[13-15].

In the control group, I included children with similar data (age, sex, place of residence).In both the main and control groups, we studied the constitutional - inherited characteristics of children, socioeconomic status of the family, physical - mental stress factor, severity of evacuation conditions, and more. To this end, we have designed a proper questionnaire that includes all of the factors listed above [16-18].

Findings: 800 children, 348 were diagnosed with somatic and mental disorders. And these children account for 43.5% of studies. Children with psycho-somatic disorders also often had psycho-emotional and neurological disorders. And the malnutrition of these children witnessed war, some were directly involved in combat operations (4.5%). Night fears, sleep disorders and other phobias (65%) were the most frequent. 6.4% of children who have been exposed to hostilities have developed logonorrhea (Fig. 1) [19-22]. 5.62% of displaced children had nocturnal enuresis, 3.75% of which developed pathology after the conflict. Nocturnal enuresis was detected in the control group with 18 children (2.25%). The difference between these groups was statistically significant.

12 cases (1.95%) of epilepsy were observed from the nervous system pathologies, of which 4 (0.5%) had pre-conflict disease, 8 (1.0%) developed post-conflict.

Eleven (1.37%) children had hyperkinesia - 6 of them (0.75%) developed post-conflict. These neurological abnormalities were mainly developed in children with conflict [23-35].




Fig. 1 Children with psycho-somatic disorders also often had psycho-emotional and neurological disorders (main group).

In the control group children, nocturnal fears, phobias, and logonaviruses were ignored. 43.0% of children expressed aggression that developed after the war. Aggressiveness was manifested by unmotivated conflicts, followed by a rash of toys and family items, with 9.0% of animals being burned. The aggressiveness of 4% of control children was expressed only by non-motivated conflicts (Fig. 2).




Fig. 2 Children with psycho-somatic disorders also often had psycho-emotional and neurological disorders (control group).

In the control group, there are 3 cases of epilepsy (0.37%), 5 cases of hypertension (0.62). The difference between the mentioned groups is statistically reliable [22, 31, 45]

From the above, we can conclude that the severe psycho-emotional and physical stress caused by the war had a strong negative impact on the health of forcibly displaced children from Abkhazia [28-30].

Conclusion. The health assessment of forcibly relocated children from Abkhazia presents significant challenges and requires a nuanced understanding of their unique needs. The impact of displacement on physical and psychological health necessitates comprehensive health evaluations and targeted interventions. By addressing the characteristics of health issues and improving methodologies for assessment, policymakers and healthcare providers can better support the well-being of these children. Enhanced health services, supportive interventions, and targeted policies are essential for addressing the complex health needs of forcibly relocated children and improving their overall quality of life. Pharmacotherapy for forcibly relocated children from Abkhazia presents several challenges, including disrupted access to medications, adherence issues, and the impact of displacement on health conditions. Addressing these challenges requires a multifaceted approach, including improving healthcare systems, supporting adherence, and addressing legal, cultural, and logistical barriers. By implementing these strategies, healthcare providers can better meet the needs of relocated children and improve their health outcomes.

References:

1. Stinivasen P. La Kahamanaswany A. socio–economic factor in birth weight of besies. Antiseptic. 2011-V.82. P27-30.
2. Von Schoen-Angerer T., Vagedes J., Schneider R. et al. Acceptance, satisfaction and cost of an integrative anthroposophic program for pediatric respiratory diseases in a Swiss teaching hospital: an implementation report. Complement Ther Med. 2017.
3. Zuzak T.J., Boňková J., Careddu D., Garami M., Hadjipanayis A., Jazbec J. et al. Use of complementary and alternative medicine by children in Europe: published data and expert perspectives. Complement Ther Med. 2013; 21: S34–S47. doi: 10.1016/j.ctim.2012.01.001.
4. Zuzak T.J., Zuzak-Siegrist I., Simões-Wüst A.P., Rist L., Staubli G. Use of complementary and alternative medicine by patients presenting to a paediatric emergency department. Eur J Pediatr. 2009; 168: 431–437. doi: 10.1007/s00431-008-0765-3.
5. Jani Asim A., Trask Jennifer, Ali Ather. Integrative Medicine in Preventive Medicine Education. American Journal of Preventive Medicine. 2015;49(5):S222–S229. doi: 10.1016/j.amepre.2015.08.019.
6. Armishaw J., Grant C.C. Use of complementary treatment by those hospitalised with acute illness. Arch Dis Child. 1999; 81:133–137. doi: 10.1136/adc.81.2.133.
7. Chan E., Rappaport L.A., Kemper K.J. Complementary and alternative therapies in childhood attention and hyperactivity problems. J Dev Behav Pediatr. 2003; 24: 4–8. doi: 10.1097/00004703-200302000-00003.
8. Roth I., Highfield L., Cuccaro P., Wells R., Misra S., Engebretson J. Employing evidence in evaluating complementary therapies: findings from an ethnography of integrative pain Management at a Large Urban Pediatric Hospital. J Altern Complement Med. 2019; 25: S95–S105. doi: 10.1089/acm.2018.0369.

9. Sanders H., Davis M.F., Duncan B., Meaney F.J., Haynes J., Barton L.L. Use of complementary and alternative medical therapies among children with special health care needs in southern Arizona. Pediatrics. 2003; 111: 584–587. doi: 10.1542/peds.111.3.584.
10. Sawni-Sikand A., Schubiner H., Thomas R.L. Use of complementary/alternative therapies among children in primary care pediatrics. Ambul Pediatr. 2002; 2: 99–103. doi: 10.1367/1539-4409(2002)002<0099: UOCATA>2.0.CO;2.
11. So M., Dziuban E.J., Franks J.L. et al. Extending the Reach of Pediatric Emergency Preparedness: A Virtual Tabletop Exercise Targeting Children's Needs. Public Health Rep. 2019;134(4):344-353. doi:10.1177/0033354919849880
12. Jackson B.A. The Problem of Measuring Emergency Preparedness: The Need for Assessing "Response Reliability" as Part of Homeland Security Planning. Santa Monica, CA: RAND Corporation; 2008;
13. Kirkpatrick D.L., Kirkpatrick J.D. Evaluating Training Programs: The Four Levels. 3rd ed San Francisco, CA: Berrett-Koehler; 2006.
14. US Department of Homeland Security. Homeland Security Exercise and Evaluation Program (HSEEP). Washington, DC: US Department of Homeland Security; 2013;
15. Behar S., Upperman J.S., Ramirez M., Dorey F., Nager A. Training medical staff for pediatric disaster victims: a comparison of different teaching methods. Am J Disaster Med. 2008; 3(4):189–199.
16. Bank I., Khalil E. Are pediatric emergency physicians more knowledgeable and confident to respond to a pediatric disaster after an experiential learning experience? Prehosp Disaster Med. 2016; 31(5):551–556. doi:10.1017/S1049023X16000704
17. Subbarao I., Lyznicki J.M., Hsu E.B. et al. A consensus-based educational framework and competency set for the discipline of disaster medicine and public health preparedness. Disaster Med Public Health Prep. 2008; 2(1):57–68. doi:10.1097/DMP.0b013e31816564af
18. Siegel D, Strauss-Riggs K, Needle S. Prioritization of pediatric BRNE disaster preparedness education and training needs. Clin Pediatr Emerg Med. 2014;15(4):309–317.
19. Abramson D.M., Park Y.S., Stehling-Ariza T., Redlener I. Children as bellwethers of recovery: dysfunctional systems and the effects of parents, households, and neighborhoods on serious emotional disturbance in children after Hurricane Katrina. Disaster Med Public Health Prep. 2010;4(suppl 1): S17–S27. doi:10.1001/dmp.2010.7
20. National Advisory Committee on Children and Disasters. Healthcare Preparedness for Children in Disasters: A Report of the NACCD Healthcare Preparedness Working Group. Washington, DC: US Department of Health and Human Services; 2015.
21. Adams J. and Wong J. (2015). Health issues in forcibly displaced populations: A review of current literature. Global Health Action. 8(1), 274-282.
22. Baker K., Robinson M. (2019). The psychological impact of displacement on children: A systematic review. Journal of Child Psychology and Psychiatry. 60(2), 124-135.
23. Jones C., Lee M. (2020). Nutritional challenges and interventions in displaced children: Evidence from recent studies. Nutrition Reviews. 78(3), 189-200.
24. Khan A., Patel V. (2017). Addressing mental health needs in displaced populations: A review of strategies and practices. Lancet Psychiatry. 4(12), 914-922.
25. Miller H., Cooper J. (2018). Evaluating health care access and quality for displaced children: A case study approach. Journal of Refugee Studies. 31(4), 585-601.
26. Smith R., Collins P. (2021). Addressing health disparities in displaced populations: Policy and practice implications. Health Policy. 125(4), 317-326.
27. Fetz K., Ostermann T., Schwermer M. et al. Do patients of integrative anthroposophic pediatric inpatient departments differ? Comparative analysis to all pediatric inpatients in Germany considering demographic and clinical characteristics. BMC Public Health. 2019 Dec 3;19(1):1623. doi: 10.1186/s12889-019-7972-x. PMID: 31795983; PMCID: PMC6889203.

28. Sagara T. Studies on the growth of Japanese schoolpils alter world waar …..j. Naramed ass 2010. 12, 2-3.
29. Sundquist J. Johansson S. The influence of exile and reparation on mental and physical health. Apopulation-basad study. Soc. Psychiatry Epidemiol. 2006 30(2), 20-26.
30. Wolansky N.The secular trend: Mikroevolution, physoilogial adaptation and migration and causative factors.7Intern COngr…oxford.2015.
31. Sundquist J. Johansson S. The influence of exile and reparation on mental and physical health. Apopulation-basad study. Soc. Psychiatry Epidemiol. 2016 30(2), 20-26.
32. McMaster H. Personal child health record and advice booklet program in Tuzlam Bosnia Herzegovina. Yug. soc. med. 2002, 90(5), 2010-2014.
33. Schumacher L. Growth of immigrant children in the newcomer schools of San Francisco. Pediatrics. 2015,80(6), 864-869.
34. Anheyer D., Anheyer M., Finetti C., Dobos G., Cramer H. Implementierung von Phytotherapie in eine Akutklinik für Kinder-und Jugendmedizin. Z Für Phytother. 2018; 39: 251–256. doi: 10.1055/a-0791-8927.
35. Beer André-Michael, Uehleke Bernhard, Wiebelitz Karl Rüdiger. The History of Inpatient Care in German Departments Focussing on Natural Healing. Evidence-Based Complementary and Alternative Medicine. 2013; 2013:1–8.
36. Eckert M., Amarell C., Anheyer D., Cramer H., Dobos G. Integrative pediatrics: successful implementation of integrative medicine in a German hospital setting—concept and realization. Children. 2018; 5:122. doi: 10.3390/children5090122.
37. Huber B.M., Schneider R., Bapst V, Wildhaber J. Integrative Pädiatrie am HFR Freiburg – Kantonsspital. Paediatrica. 2018; Vol. 29 Nr. 1 2018.
38. Längler A., Zuzak T.J. Complementary and alternative medicine in paediatrics in daily practice – a European perspective. Complement Ther Med. 2013;21: S26–S33.
39. World Health Organization. (2022). Health of refugees and migrants: Toolkit for practitioners. World Health Organization.
40. Gordon, M. S., & Anselmo, M. (2018). Managing chronic conditions in displaced populations: Challenges and strategies. Journal of Refugee Health. 9(2), 102-112.
41. Haddad S., Whelan M. (2016). Access to medications for displaced children: A review of barriers and solutions. Global Health Journal. 10(1), 45-56.
42. Johnson A., Patel V. (2020). Addressing medication adherence in vulnerable populations: Insights and interventions. Pharmacy Practice. 17(3), 75-84.
43. Miller R., Johnson K. (2017). Displacement and health: A review of the impact of forced migration on chronic illness management. International Journal of Environmental Research and Public Health. 14(5), 1234-1245.
44. Smith L., Nguyen D. (2019). Cultural competence in healthcare for displaced populations: A critical review. Journal of Health Disparities Research and Practice, 12(2), 34-45.
45. World Health Organization. (2021). Medication access and adherence in displaced populations: Guidelines and recommendations. World Health Organization.

Розділ 4

АДМІНІСТРУВАННЯ АНТИМІКРОБНИХ ПРЕПАРАТІВ ЯК МУЛЬТИДИСЦИПЛІНАРНИЙ ПІДХІД ПРОТИДІЇ АНТИБІОТИКОРЕЗИСТЕНТНОСТІ

ПИТАННЯ ПОДОЛАННЯ АНТИБІОТИКОРЕЗИСТЕНТНОСТІ В УКРАЇНІ

Паламар А. О., Круглик Я. Д.
Буковинський державний медичний університет,
м. Чернівці, Україна
Кафедра фармації
palamar@bsmu.edu.ua

Вступ. Протимікробні препарати вкрай важливі для успішного лікування інфекційних захворювань, особливо важких, проте їх надмірне та нераціональне застосування значно прискорило процес формування резистентності мікроорганізмів до відомих препаратів. Стрімке поширення антибіотикорезистентності підтверджує, що подальше їх безконтрольне застосування призведе до неефективності цих лікарських засобів, збільшенню захворювань та значної кількості летальних випадків. Згідно зі статистикою ВООЗ, понад 1 млн смертей щорічно пов'язані з антибіотикорезистентністю, при цьому прогнози вказують на те, що в найближчі десятиліття ця цифра зростатиме, і до 2050 року може досягти понад 10 млн.

Мета. Провести аналіз раціонального застосування антибактеріальних препаратів з урахуванням наказів та стандартів МОЗ України, спрямованих на подолання антибактеріальної резистентності.

Матеріали та методи. Використано офіційні документи міжнародних організацій, законодавчі акти та нормативно-правові документи, публічні матеріали та результати досліджень з питань проблеми антибіотикорезистентності, що опубліковані у відкритому доступі.

Результати. На сьогодні, більша частина пацієнтів не дотримуються схеми прийому антибактеріальних препаратів, призначеної лікарем. При зникненні симптомів хвороби, ні в якому разі не можна припиняти лікування, оскільки це підвищує ризик виникнення стійких до антибіотиків штамів бактерій, чи самовільно знижувати дозу призначеного препарату або приймати ліки, що залишилися після минулого курсу лікування. Важливо пам'ятати, що антибіотики не тільки не ефективні при вірусних захворюваннях, але, й навпаки, можуть завдати шкоди ослабленому організму. Саме тому, їх не слід призначати собі самостійно, тим паче з метою профілактики.

Безконтрольне вживання антибіотиків сприяє розвитку стійкості бактерій до цих препаратів і, як наслідок, вони перестають діяти, а інфекційні захворювання не піддаються лікуванню – антибіотикорезистентність значно поширюється.

Фармацевтичні працівники, найчастіше, це перші до кого звертаються пацієнти із проханням консультації та відпуску антибактеріального препарату. Фармацевт при відпуску антибіотика повинен надати максимальне пояснення щодо препарату, правил прийому, можливі побічні дії та зазначити, що антибіотикотерапія має тривати стільки, скільки призначив лікар, і не перерива-

тися самостійно. При цьому, відпуск антибактеріальних препаратів має здійснюватися виключно за рецептом лікаря, разом з тим, впровадження електронного рецепта на антибіотики (наказ МОЗ від 21.07.2022 № 1284) дозволяє відстежувати обсяги застосування антибактеріальних засобів і протидіяти розвитку антибіотикорезистентності.

Важливу роль у боротьбі з антибіотикорезистентністю відіграє просвітницька робота серед пацієнтів. Не від'ємним у роботі фармацевта є уточнення доцільності призначення того чи ін. препарату для конкретного пацієнта (призначення згідно з віком, масою тіла та діагнозом).

З березня 2019 МОЗ України розпочав інтенсивну боротьбу з поширенням антибіотикорезистентності в країні, було розроблено «Національний план дій щодо боротьби із стійкістю до протимікробних препаратів», який включає розширення мережі мікробіологічних лабораторій, посилення інфекційного контролю в закладах охорони здоров'я та епідеміологічний нагляд за резистентними збудниками.

Розробка МОЗ України Інструкції з впровадження адміністрування антимікробних препаратів у стаціонарах (наказ МОЗ України від 03.08.2021 № 1614) свідчить, що в закладах охорони здоров'я, які надають медичну допомогу в стаціонарних умовах має бути сформований Відділ інфекційного контролю, де фармацевт клінічний відповідає за моніторинг антибіотиків і консультує лікарів щодо оптимізації фармакотерапії.

Затверджено Стандарт медичної допомоги «Раціональне застосування антибактеріальних і антифунгальних препаратів з лікувальною та профілактичною метою» (наказ МОЗ від 23.08.2023 № 1513), який встановлює загальні вимоги до призначення, коригування і припинення антибактеріальної терапії.

Для запобігання розвитку антибіотикорезистентності, медичні працівники повинні дотримуватися гігієни, призначати антибіотики лише за потребою, передавати інформацію про стійкість у пацієнтів, пояснювати, як правильно приймати ці препарати, і розповідати про профілактику. Важливо, щоб медичні працівники усвідомлювали проблему стійкості до антимікробних препаратів і ділилися цією інформацією з пацієнтами, допомагаючи їм правильно ставитися до лікування та профілактики інфекцій.

Висновки. Враховуючи надмірне та необґрунтоване застосування антибактеріальних препаратів, що призводить до модифікації нових збудників, які стають нечутливими до лікування відомими препаратами, слід продовжувати боротьбу з поширенням антибіотикорезистентності в Україні. Найдієвіший спосіб подолання цієї проблеми – комплексний підхід, що базується на просвітницькій роботі серед медичних, фармацевтичних працівників та населення, дотриманні протоколів призначення антибіотиків і постійне їх оновлення, а також регулярний моніторинг і контроль за частотою їх використання.

ПЕРЕХРЕСТЯ АНТИБІОТИКОРЕЗИСТЕНТНОСТІ ТА БІОСИНТЕЗУ АНТИБІОТИКІВ

Кравченко В. М., Філімонова Н. І., Сенюк І. В.
Національний фармацевтичний університет, м. Харків, Україна
Кафедра клінічної лабораторної діагностики, мікробіології та біологічної хімії
citochrom@gmail.com

Вступ. Антибіотики залишаються найважливішим ресурсом у глобальному лікуванні інфекційних захворювань. Зростання резистентності патогенів людини до антибіотиків викликало глобальне занепокоєння, оскільки антибіотики невпинно втрачають ефективність у клінічних умовах. Незважаючи на зростання резистентності до антибіотиків і зміни програм розробки ліків у приватному секторі, ринок антибіотиків залишається потужним. У 2018 році FDA схвалило чотири нові антибіотики з традиційних класів антибіотиків, включаючи аміноглікозид плазоміцин і тетрацикліни еравациклін, омадациклін і сарециклін. Ці нові антибіотики отримали схвалення, зокрема, завдяки подоланню встановлених клінічних механізмів резистентності для задоволення клінічних потреб. Згідно досвіду клінічної практики, нові механізми резистентності з'являться після широкомасштабного застосування цих нових препаратів

Мета. Проведення аналізу походження антибіотикорезистентності задля актуалізації питань перехрестя біосинтезу антибіотиків і самозахисту продуцента, що призводить до клінічно значущих механізмів резистентності.

Матеріали та методи. Використані пошукові системи Google, Google Scholar, Scopus, PubMed і ResearchGate з використанням ключових слів за обраною тематикою.

Результати. Самозахист є необхідною умовою для будь-якого мікроорганізма, що продукує антибіотик. Кластери біосинтетичних генів антибіотиків (БГА) містять гени, що кодують ензими, необхідні як для біосинтетичної збірки каркасу антибіотиків, так і для механізмів самозахисту. Ця спільна кластеризація призводить до спільної експресії генів і забезпечує самозахист під час синтезу токсичного антибіотика. Самозахист продуцентів антибіотиків часто називають вродженою стійкістю, тоді як поява генів самозахисту у синтезі антибіотиків відома як набута стійкість.

Потужна форма резистентності до антибіотиків виникає у формі інактивації через активність ензимів. На відміну від інших механізмів резистентності, ковалентна модифікація антибіотиків знижує загальну концентрацію антибіотика нижче інгібуючого рівня. Потік ферментативного конструювання антибіотиків (біосинтез) і руйнування (самозахист/резистентність) впливає на динаміку мікробної популяції та сприяє появі ензимів, що інактивують антибіотики. Ензими, що нейтралізують антибіотики, забезпечують клітинний імунітет, модифікуючи частину каркаса антибіотиків, необхідну для зв'язування мішені.

Відмінності виникають у типах хімічних зав'язків при порівнянні самозахисту та резистентності у клінічних патогенів.

У той час як макролідне фосфорилювання є достатньо незворотною модифікацією, щоб надати резистентність, більш ефективною стратегією резистентності є деструктивний гідроліз макролактонового кільця, спільного для всього класу антибіотиків. Кодовані плазмідами макролідні естерази, включаючи еритроміцинестеразу EreA, були виявлені у клінічних патогенах.

Висококонсервативні функціональні групи та реактивні «м'які місця» в каркасі є основними мішенями для інактивації ензимів резистентності. Стійкість може бути досягнута, лише якщо хімічна модифікація є незворотною у біологічному масштабі за часом і якщо модифікація блокує доступ та зв'язування з мішенню. Найнебезпечніші ензими, що інактивують антибіотики, повністю «знищують» каркас антибіотиків (наприклад, бета-лактамази та тетрацикліндеструктази), мають широке покриття субстрату в усьому класі антибіотиків і експресуються у відповідь на антибіотичний виклик. Жоден антибіотик, навіть відомий тейксобактин, який, як стверджується, не має «виявленої стійкості» не може повністю уникнути появи інактивуючих ензимів.

Зв'язок між біосинтезом антибіотиків і резистентністю відіграє все більш важливу роль у пошуках антибіотикорезистентності та антибіотиків наступного покоління. Біосинтетичний генний кластер (БГК) мікробних антибіотиків є довгостроковим сховищем для генів стійкості до антибіотиків у різноманітних середовищах. Формування БГК, керований резистентністю, за останні роки виявив кілька нових антибіотиків і допоміг розширити структурний пул відомих класів антибіотиків. Шляхом скринінгу невеликої бібліотеки з 1000 актиноміцетів проти ванкомцину було досягнуто 4% показника резистентності, що корелювало з експресією оперону vanHAX для стійкості до глікопептидів. Секвенування генома за допомогою спільного кластеризування оперону vanHAX з глікопептидними БГК привело до відкриття пекіскоміцину з новим глікопептидим каркасом. Скринінг, орієнтований на резистентність, виявився ще більш корисним в останні роки за рахунок зростання кількості секвенованих геномів штамів продуцентів антибіотиків.

Стійкість до антибіотиків – це те, що слід враховувати на самих ранніх стадіях розробки антибіотиків. Резистентністю слід керувати, а не уникати її шляхом фармацевтичної оптимізації.

Висновки. Стійкість до антибіотиків врівноважується біосинтезом антибіотиків у природному середовищі. Стійкість патогенів людини виникає через механізми самозахисту у продуцентів антибіотиків.

Ензими, що інактивують антибіотики, розвиваються, щоб модифікувати збережені дизайни каркасу, необхідні для зв'язування мішені.

Гібридні антибіотики, ад'юванти та синергетичні комбінації є природною платформою для подолання резистентності.

Проактивне вивчення резистентності може бути використано для розробки антибіотиків наступного покоління.

ПАНДЕМІЯ COVID-19: РОКИ ВПЛИВУ НА АНТИМІКРОБНУ РЕЗИСТЕНТНІСТЬ

Шаповалова О. В. [1], Кучма І. Ю. [2]

[1]Національний фармацевтичний університет, м. Харків, Україна
Кафедра клінічної фармакології
[2]Навчально-науковий медичний інститут НТУ «Харківський політехнічний інститут», кафедра мікробіології, бактеріології, вірусології та мікології
shapolga2002@gmail.com, irina_kuchma@ukr.net

Вступ. Пандемічний вірус SARS-CoV-2 та хвороба COVID-19, яку він спричинює, відомі людству вже майже п'ять років, а саме з 31 грудня 2019 року. Натепер відомі біологія збудника, особливості епідеміології захворювання, досягнуті значні успіхи в діагностиці, лікуванні та профілактиці COVID-19. Однак все гостріше постає проблема взаємозв'язку тривалої боротьби з пандемією та невдач у протистоянні розвитку антимікробної резистентності мікроорганізмів (АМР).

Мета. Провести аналіз впливу періоду пандемії COVID-19 2020-2023 р.р. на стан проблеми антимікробної резистентності мікроорганізмів.

Матеріали та методи. Проводили аналіз доступних джерел наукової літератури з використанням бази данних MEDLINE, а також матеріалів сайтів ВООЗ та CDC.

Результати. Невтішна статистика свідчить, що на тлі критичного стану системи охорони здоров'я, яка виявилась майже безсильною проти SARS-CoV-2 на початку 2020 р., почалось значне збільшення використання антимікробних препаратів та недотримання рекомендацій щодо профілактики АМР. В США протягом першого року пандемії спостерігали ріст інфекцій, пов'язаних з наданням медичної допомоги, на 15%. При цьому антибіотики отримували приблизно 80% пацієнтів, госпіталізованих із COVID-19. В цей період 29400 людей померли від інфекцій, що були викликані стійкими до антимікробних препаратів бактеріями.

Перенавантаження лікарень, тривале перебування хворих у відділеннях інтенсивної терапії, брак персоналу, лікарських препаратів, засобів індивідуального захисту та ізольованих приміщень мали негативний вплив на реалізацію стратегії боротьби з АМР. Цьому також сприяли організаційні прогалини через перенесення пріоритетів на COVID, в тому числі й матеріальних витрат.

Матеріали звіту CDC щодо аналізу числа резистентних до антимікробних препаратів госпітальних штамів, проведеного в 2020 р. на базі 600 лікарень в США, свідчать про зростання кількості стійких до карбапенемів бактерій роду Acinetobacter на 78% та родини Enterobacteriacea на 35%; ентеробактерій, що продукують бета-лактамази розширеного спектру дії (ESBL) на 32%; мультирезистентних Pseudomonas aeruginosa на 32%; метицилінрезистентних Staphylococcus aureus (MRSA) на 13%; ентерококів, стійких до ванкоміцину на 14%. В той же час спостерігали розвиток стійкості пріорітетних збудників гри-

бкових інфекцій до антимікотичних препаратів: у 60% ізолятів виду Candida auris та 26% штамів інших видів роду Candida.

Як відомо, вторинні бактеріальні та грибкові інфекції суттєво впливають на епідеміологічні показники захворюваності та смертності під час спалахів вірусних інфекцій і пандемій, прикладом цього слугує грип. Однак, результатом неконтрольованого застосування антибіотиків є підвищення АМР, виникнення нової загрози – бактеріальної пандемії або вірусної пандемії, що супроводжується вторинними бактеріальними або грибковими інфекціями, стійкими до наявних протимікробних засобів.

Щодо COVID-19, за даними Американського товариства інфекційних захворювань (IDSA) при коронавірусній хворобі лише 8% пацієнтів з вторинною бактеріальною/грибковою інфекцію потребують призначення антибіотиків. Однак згідно клінічних рекомендацій на початку пандемії COVID-19 пацієнти отримували антибіотики широкого спектру дії навіть за відсутності бактеріальних ускладнень. Так, в Італії антибіотики отримували 66,67% пацієнтів, в Іспанії – 80%, в Китаї – 95% пацієнтів пульмонологічних лікарень. Загалом від 70% до 95% госпіталізованих пацієнтів отримували або доксициклін, або амоксицилін. Стандартні рекомендації також включали бензилпеніцилін, кларитроміцин, метронідзол, моксифлоксацин, ципрофлоксацин як препарати першої лінії лікування COVID-19.

Окреме занепокоєння викликає лікування інфекції SARS-CoV-2 у носіїв резистентних штамів, оскільки за даними CDC приблизно 2% населення є носіями MRSA та штамів, що продукують ESBL/AmpC. Як наслідок, такі пацієнти з COVID-19 мають підвищений ризик розвитку вторинних інфекцій резистентними бактеріями. Одним з ранніх проявів побічних ефектів невдалих призначень для лікування COVID-19 також є розвиток мукоромікозу, найбільш вірогідною причиною якого вважається поєднання терапії антибіотиками та стероїдами, зниження імунного захисту та дисглікемії (епідемія мукоромікозу була оголошена в Індії в 2021 р.).

Додатковим фактором поширення АМР є масштабне потрапляння біоцидів у довкілля в результаті збільшення споживання дезінфікуючих засобів та фармацевтичних продуктів для стримування поширення вірусу.

Висновки. Пандемія COVID-19 висвітлила багато викликів, з якими стикається людство під час реагування на сучасні надзвичайні ситуації у сфері охорони здоров’я. Одним з таких викликів є ріст АМР патогенів. Шляхами стримування АМР є раціональне застосування антимікробних препаратів, широке впровадження швидких тестів для виявлення збудників та тестування їх чутливості до антимікробних препаратів, заборона призначення антибіотиків при вірусних інфекціях, створення нових лікарських засобів, надійний контроль за їх призначенням, моніторинг розвитку АМР.

АНАЛІЗ РИНКУ ДИТЯЧИХ АНТИМІКРОБНИХ ЛІКАРСЬКИХ ЗАСОБІВ ДЛЯ ПЕРОРАЛЬНОГО ПРИЙОМУ: РИЗИКИ АНТИБІОТИКОРЕЗИСТЕНТНОСТІ

Воронкіна А. С., Дідківська А. О., Хорунжий Д. С.
Вінницький національний медичний університет ім. М.І. Пирогова, м. Вінниця, Україна
Кафедра фармації
voronkina@vnmu.edu.ua

Вступ. Антибіотикорезистентність – одна з найсерйозніших загроз для глобального здоров`я, особливо в умовах, коли неконтрольоване використання антибіотиків стає частим явищем. В Україні додатковим фактором ризику є зростання кількості пацієнтів, що потребують тривалої антибіотикотерапії, зокрема серед військових з гнійно-опіковими ранами та післяопераційними ускладненнями. Згідно з даними ВООЗ, найбільш частими збудниками внутрішньолікарняних антибіотикорезистентних інфекцій є патогени групи ESKAPE: *Enterococcus faecium, Staphylococcus aureus, Klebsiella pneumonia, Acinetobacter baumannii, Pseudomonas aeruginosa, Enterobacter spp.* Частково вони можуть перетинатися з типовими збудниками респіраторних інфекцій у дітей, зокрема *S. aureus* та *K. pneumoniae* є збудниками пневмонії. Натомість ринок дитячих антимікробних засобів для перорального прийому демонструє тривожні тенденції, що в подальшому можуть призвести до погіршення ситуації з антибіотикорезистентністю.

Мета роботи. Дослідити ринок дитячих антимікробних препаратів для перорального прийому в Україні, співставивши його асортимент з класифікацією антибіотиків за системою AWaRe від ВООЗ 2021 та 2023 років, а також проаналізувати динаміку роздрібних продажів (в упаковках) антибіотиків груп Access та Watch у першому півріччі 2024 року.

Матеріали та методи. Використано дані з відкритих джерел (Державний реєстр лікарських засобів, Компендіум, Таблетки.юа), списки ВООЗ AWaRe 2021, 2023 років, базу ProximaResearch від компанії Моріон для отримання даних щодо продажів препаратів. Проведено кількісний аналіз обраного сегменту ринку (продажі в упаковках) за перше півріччя 2024 року, а також порівняння даних за попередніх чотири роки.

Результати. ВООЗ згідно класифікації AWaRe поділяє антибіотики на три групи: **Access** (Доступу): антибіотики першого вибору, що мають бути широко доступні і підходять для емпіричного призначення без визначення чутливості збудника з мінімальним ризиком розвитку резистентності; Watch (Спостереження): препарати, що мають підвищений ризик розвитку резистентності та потребують контрольованого використання, зокрема попереднього визначення чутливості збудника до антибіотикотерапії; та Reserve (Резерв): антибіотики останньої лінії, що застосовуються лише у крайніх випадках, в разі підтвердженої резистентності мікроорганізму до дії препаратів перших двох груп,

індивідуальної непереносимості, тощо. Щодвароки переліки антибіотиків у списках AWaRe змінюються, що свідчить про визнання глобальної потреби в обмеженні використання критично важливих препаратів через зростання резистентності. Серед дитячих лікарських засобів для пероральної антибіотикотерапії (у формі порошків або гранул для приготування суспензії) в Україні зареєстровані лише антибіотики груп Access (2 міжнародних непатентовані назви (МНН), 9 торгові назви (ТН)) та Watch (7 МНН, 19 ТН). Загальне співвідношення за кількістю МНН між препаратами груп Access, Watch та Reserve серед дитячих пероральних лікарських засобів становить 2:7:0. Натомість, у переліку ВООЗ "Model list of essential medicines for children 2023" це співвідношення складає 25:14:6 відповідно. Очевидно, що асортимент антибіотиків групи спостереження в Україні значно ширший за такий для групи доступу, що може бути спричинене більшим попитом. Але антибіотики спостереження превалюють не лише за асортиментними показниками. За даними ProximaResearch, продажі дитячих антибіотиків для перорального прийому в групі Access за перше півріччя 2024 року в упаковках складають лише близько 32% загальних продажів, а у групі Watch – 68% від загальної кількості відповідно). Домінують за кількістю ТН у групі Watch препарати групи цефалоспоринів (4 МНН, 12 ТН), зокрема, переважно третього покоління (3 МНН, 11 ТН), відомі своєю здатністю в тому числі до формування перехресної резистентності з іншими бета-лактамними антибіотиками, які, власне, і формують групу Access. За об'ємами продажів в упаковках серед препаратів спостереження переважають макроліди (зокрема, азитроміцин). Таке співвідношення зберігається протягом останніх чотирьох років, з незначними коливаннями: препарати з вищим ризиком розвитку резистентності (група Watch), використовуються частіше за антибіотики першого ряду, а це може призвести до тяжких наслідків.

Висновки. Значне домінування продажів препаратів з групи Watch свідчить про недотримання рекомендацій щодо використання препаратів першого вибору з групи Access, що створює ризики для розвитку антибіотикорезистентності. Динаміка змін списків AWaRe вказує на значне занепокоєння світової медичної спільноти та посилення заходів щодо обмеження використання критично важливих антибіотиків. Важливо забезпечити і в Україні контрольоване використання антибіотиків та підвищити рівень обізнаності серед медичних працівників та пацієнтів щодо небезпеки антибіотикорезистентності, особливо в умовах військового стану.

THERAPY FOR MULTIDRUG-RESISTANT TUBERCULOSIS

Kolomiets Y., Kharchenko Y.
Dnipro State Medical University, Dnipro, Ukraine
Department of pharmacology, general and clinical pharmacy
209_14@dmu.edu.ua

Introduction. Tuberculosis is an airborne disease caused by the pathogen, Mycobacterium tuberculosis, which predominantly affects the lungs. This type of bacteria latently infecting approximately a quarter of the worldwide population.

Aim. New strategy of multidrug-resistant tuberculosis (MDR TB) treatment.

Materials and methods. An information search and analysis of the scientific literature on the treatment of MDR TB was conducted on the PubMed platform.

The results. World Health Organization has reported that about 85% of tuberculosis patients are cured with the existing 6-month antibiotic regimen. However, the lengthy oral administration of high-dose anti-tuberculosis drugs is associated with significant side effects and leads to drug resistance cases. Alternatively, reformulating existing anti-tubercular drugs into inhalable nanoparticulate systems is a promising strategy to overcome the challenges associated with oral treatment as they could enhance drug retention in the pulmonary region to achieve an optimal drug concentration in the infected lungs. Many preclinical studies have shown that inhalable nanoparticles increased therapeutic efficacy and minimized adverse drug reactions when delivered through the pulmonary route, but none of them has progressed into clinical trials to date. Poverty, difficulty to access proper treatment, and lack of patient compliance contributed to the emergence of more sinister drug-resistant strains, which demand a longer duration of treatment with more toxic and more expensive drugs. Only three new drugs, bedaquiline, delamanid and pretomanid were approved in the last decade for treatment of tuberculosis. Bedaquiline is a first of its class drug that inhibits mycobacterial adenosine triphosphate synthase's proton pump. There are many clinical trials and cohort studies that reported a higher culture conversion rate with bedaquiline treatment as compared to the control group. Many meta-analyses and systematic reviews have reported higher cure rate, and lower mortality rate in patients with MDR TB treated with a bedaquiline-containing regimen. Delamanid and pretomanid share many similarities, buy several differences could be identified. Is that certain Mycobacterium tuberculosis isolates have been described that are resistant to either delamanid or pretomanid, but with preserved susceptibility to the other compound. This might imply that delamanid and pretomanid could replace one another in certain regimens. Regarding bactericidal activity, based on in vitro and preclinical in vivo activity, delamanid has lower MICs and higher mycobacterial load reductions at lower drug concentrations and doses compared with pretomanid.

Conclusions. Bedaquiline and the two nitroimidazole derivatives delamanid and pretomanid the first anti-tuberculosis drugs with novel mode of actions to be introduced to the market in more than 50 years. Nanoparticles, which the patient inhales, are also effective for the treatment of MDR TB. In this way, the therapeutic effect increases and the risks of side effects of drugs during delivery through the lungs decrease.

ВИВЧЕННЯ РОЛІ ФАРМАЦЕВТА В ЗАСТОСУВАННІ ПРОБІОТИКІВ ПРИ ПРОВЕДЕННІ АНТИБАКТЕРІАЛЬНОЇ ТЕРАПІЇ

Паскевич М. П., Жаботинська Н. В.
Національний фармацевтичний університет, м. Харків, Україна
Кафедра фармакології та клінічної фармації
bronkevih@gmail.com

Вступ. Антибактеріальна терапія є ефективним способом лікування інфекцій, проте вона часто супроводжується побічними ефектами, зокрема розвитком дисбіозу або антибіотико-асоційованої діареї. Пробіотики можуть допомогти відновити баланс мікрофлори та запобігти цим ускладненням. Однак, застосування пробіотиків серед населення не завжди є раціональним, часто використовується безконтрольно та без належної консультації з фахівцями. Фармацевт відіграє важливу роль у забезпеченні грамотного використання пробіотиків під час антибактеріальної терапії, надаючи рекомендації пацієнтам щодо їхнього застосування.

Мета. Метою цього дослідження є статистичний аналіз поширеності застосування пробіотиків серед населення України під час і після антибактеріальної терапії та вивчення ролі фармацевта у покращенні якості фармацевтичної опіки при відпуску пробіотиків.

Матеріали та методи. Планується проведення двох онлайн-анкетувань: анкетування серед населення України для з'ясування рівня обізнаності та практики використання пробіотиків під час та після антибактеріальної терапії та анкетування фармацевтів для оцінки їхньої ролі в наданні рекомендацій щодо пробіотиків та виявлення прогалин у знаннях і практиках консультування. По завершенню анкетування планується статистичний аналіз зібраних даних для виявлення ключових тенденцій та впливових факторів на використання пробіотиків населенням.

Результати. Очікується, що в рамках дослідження буде оцінено ступінь поширеності практики застосування пробіотиків серед населення України під час антибактеріальної терапії. Планується виявити основні джерела інформації, якими користуються пацієнти при виборі пробіотиків, а також їхні мотиви щодо застосування цих препаратів. Очікується також визначити ключові аспекти фармацевтичної опіки, які потребують вдосконалення з метою підвищення ефективності антибактеріальної терапії та зменшення ризиків побічних ефектів. На основі результатів дослідження будуть сформовані рекомендації щодо підвищення кваліфікації фармацевтів у сфері застосування пробіотиків, що сприятиме покращенню фармацевтичної опіки та ефективності лікування.

Висновок. Дослідження допоможе виявити основні проблеми та тенденції у використанні пробіотиків населенням, а також визначити, як можна підвищити роль фармацевтів у забезпеченні раціонального застосування цих препаратів.

УДК 614.2

АНАЛІЗ ПРОБЛЕМНИХ ПИТАНЬ ПРОФЕСІЙНОЇ ДІЯЛЬНОСТІ КЛІНІЧНИХ ФАРМАЦЕВТІВ У ВІДДІЛАХ ІНФЕКЦІЙНОГО КОНТРОЛЮ

Міщенко О. Я., Андріяненков О. В., Шаповалова О. В.
Національний фармацевтичний університет, м. Харків, Україна
Інститут підвищення кваліфікації спеціалістів фармації
Кафедра клінічної фармакології
clinpharmacol_ipksph@nuph.edu.ua

Анотація. Метою дослідження було проведення аналізу проблемних питань професійної діяльності клінічних фармацевтів у відділах інфекційного контролю.

Методи дослідження: проведено анкетування 59 клінічних фармацевтів ВІК ЗОЗ України щодо їх трудових функцій та успішності їх виконання на посаді клінічного фармацевта.

Результати: Отримані результати дослідження висвітлюють певні проблемні питання в роботі клінічних фармацевтів ВІК, зокрема щодо наявності медичної інформаційної системи із можливістю відстежування призначення АМП (54,2 % опитуваних клінічних фармацевтів стверджують про відсутність такої у ЗОЗ), також щодо наявності затверджених нових клінічних протоколів емпіричної АМП-терапії щодо негоспітальних інфекцій (інфекції дихальної системи, шкіри та підшкірної клітковини, кишкові інфекції) та щодо інфекцій, пов'язаних з наданням медичної допомоги (ІПНМД). Найбільш налагодженим є створення СОП з ААМП, процес преавторизації АМП та моніторингу призначення АМП (96,6 % клінічних фармацевтів розробили та затвердили керівником ЗОЗ СОП з ААМП; 89,8% – впровадили процедуру преавторизації для АМП групи резерву та відповідно 98,3% опитуваних проводять моніторинг призначення АМП (щонайменше групи резерву).

Висновки. Найбільш проблемними питаннями в роботі клінічних фармацевтів ВІК є питання наявності медичної інформаційної системи із можливістю відстежування призначення АМП, розробки нових клінічних протоколів емпіричної АМП-терапії щодо негоспітальних інфекцій (інфекції дихальної системи, шкіри та підшкірної клітковини, кишкові інфекції) та щодо ІПНМД. Найбільш налагодженими процесами є створення СОП з ААМП, преавторизації АМП, моніторингу призначення АМП, зокрема групи резерву.

Ключові слова: *клінічний фармацевт, адміністрування антимікробних препаратів, відділ інфекційного контролю*.

UDC 614.2

ANALYSIS OF PROBLEM ISSUES OF PROFESSIONAL ACTIVITY OF CLINICAL PHARMACISTS IN INFECTION CONTROL DEPARTMENTS

Mishchenko O.Ya., Andriyanenkov O.V., Shapovalova O.V.
National University of Pharmacy, Kharkiv, Ukraine
Institute for Advanced Training of Pharmacy Specialists
Department of Clinical Pharmacology
clinpharmacol_ipksph@nuph.edu.ua

Abstract. The purpose of the study was to analyze problematic issues of professional activity of clinical pharmacists in infection control departments (ICD).

Research methods: a survey was conducted of 59 clinical pharmacists of ICD of health care institutions of Ukraine regarding their work functions and the success of their performance as a clinical pharmacist.

Results: The obtained results of the study highlight certain problematic issues in the work of clinical pharmacists of ICD, in particular, regarding the availability of a medical information system with the possibility of tracking the prescription of antimicrobial drugs (AMD) (54.2% of the surveyed clinical pharmacists claim that there is no such system in the health care center), as well as the presence of approved of new clinical protocols of empiric antimicrobial therapy therapy for nosocomial infections (infections of the respiratory system, skin and subcutaneous tissue, intestinal infections) and for healthcare-associated infections (HCAI). The most well-established is the creation of standard operating procedures (SOPs) for the administration of antimicrobial drugs (AAMDs), the process of preauthorization of AMPs and monitoring of the appointment of AMPs (96.6% of clinical pharmacists developed and approved the SOP for AAMDs by the head of the health care center; 89.8% implemented the preauthorization procedure for AMDs of the reserve group and, accordingly, 98.3% of respondents monitor the appointment of AMPs (at least of the reserve group).

Conclusions. The most problematic issues in the work of clinical pharmacists of ICD are the availability of a medical information system with the possibility of tracking the appointment of AMDs, the development of new clinical protocols of empirical antimicrobial therapy therapy for non-hospital infections (respiratory system infections, skin and subcutaneous tissue infections, intestinal infections) and for HCAI. The most well-established processes are the creation of SOPs with AAMDs, pre-authorization of AMDs, monitoring of the assignment of AMDs, in particular the reserve group.

Key words: *clinical pharmacist, administration of antimicrobial drugs, infection control department.*

Вступ. Основним завданням професійної діяльності клінічного фармацевта у клінічному закладі охорони здоров’я є підвищення ефективності та безпеки лікарської терапії конкретного хворого, зокрема при призначенні антимікробних препаратів (АМП) з метою протидії резистентності до АМП

(АМР) [1, 2]. З цієї точки зору, діяльність клінічного фармацевта в клінічних закладах охорони здоров'я (ЗОЗ) на посаді координатора з адміністрування антимікробних препаратів у відділі інфекційного контролю є дуже важливою і актуальною.

Адміністрування АМП (ААМП, англійською Antimicrobial stewardship) - це послідовний набір дій, які сприяють відповідальному їх застосуванню. Це визначення можна використовувати щодо призначень АМП як на індивідуальному, так і на національному та глобальному рівнях, а також це стосується здоров'я як людей, так і тварин і навколишнього середовища. Вперше організаційну діяльність з ААМП описано на початку 1970-х років. Термін «адміністрування антимікробних препаратів» застосований у 1997 р. як опис набору стратегій, правил, рекомендацій або інструментів, які могли б покращити призначення АМП з метою зменшення їх необгрунтованого застосування та АМР [1].

Завданнями впровадження заходів з ААМП є: профілактика розповсюдження АМР в ЗОЗ і поза ним; раціональне застосування АМП з профілактичною і лікувальною метою; підвищення ефективності емпіричної АМП-терапії; оптимізація / зниження витрат ЗОЗ на АМП; зменшення терміну перебування пацієнтів на стаціонарному лікуванні; оптимізація підходів до лікування інфекцій, пов'язаних знаданням медичної допомоги (ІПНМД) [2, 3].

Адміністрування АМП, що здійснює клінічний фармацевт ВІК, включає: проспективне фармацевтичне консультування; аналіз причин безпідставної видачі антимікробного препарату широкого спектру дії з аптеки у клінічні відділення; перевірка наявності та інформативності записів у медичній документації щодо обґрунтування призначення АМП, заміни та відміни антимікробного препарату; ретроспективний аудит відповідності призначення антимікробної терапії затвердженим медико-технологічним документам; оцінка рівня споживання антимікробних препаратів в ЗОЗ; забезпечення зворотного зв'язку з лікарями (обов'язково донесення результатів до керівництва та обговорення їх з лікарями і визначення подальших дій (рекомендацій) щодо вдосконалення застосування ЛЗ) [2-4].

Етапи впровадження адміністрування антимікробних препаратів визначені Наказом МОЗ України від 03.08.2021 № 1614 «Про організацію профілактики інфекцій та інфекційного контролю в закладах охорони здоров'я та установах / закладах надання соціальних послуг / соціального захисту населення».

Формування ВІК передбачено наказом Міністерства охорони здоров'я України «Про організацію профілактики інфекцій та інфекційного контролю в закладах охорони здоров'я та установах/закладах надання соціальних послуг/соціального захисту населення» від 3 серпня 2021 р. № 1614. Положення про відділ з інфекційного контролю закладу охорони здоров'я є частиною цього наказу. Склад відділу з інфекційного контролю залежить від рівня надання медичної допомоги. У закладах, які надають амбулаторно-поліклінічну допомогу, ВІК потенційно може складатися з одного спеціаліста. Цей спеціаліст – і

керівник відділу, і водночас координатор з гігієни рук. Для допомоги йому потрібен тренер з гігієни рук та спеціаліст зі збору даних, які можуть працювати за сумісництвом. У ЗОЗ, що надають медичну допомогу в стаціонарних умовах, склад ВІК більш чисельний, а саме: керівник відділу, лікар-епідеміолог, координатор з гігієни рук, координатор з навчання і підготовки, клінічний фармацевт та статистик.

Основне завдання ВІК – організація профілактики інфекцій та інфекційного контролю в ЗОЗ/ЗСЗ з метою запобігання поширенню інфекційних хвороб, у тому числі пов’язаних із наданням медичної допомоги, мікроорганізмів із антимікробною резистентністю та формування культури безпеки в ЗОЗ.

У ВІК працює мультидисциплінарна команда. Фахівці кожної спеціальності досліджують ситуацію зі своєї точки зору, а потім обмінюються результатами, висновками, що створює підґрунтя для розроблення узгодженого плану менеджменту протидії АМР. Першочерговою умовою ефективної роботи ВІК є злагодженість роботи членів мультидисциплінарної команди, чітке розуміння своїх особистих обов'язків і загального плану дій, а також взаєморозуміння і взаємопідтримка. Велике значення має чітке розуміння керівництвом ЗОЗ та лікарями нових вимог і підходів щодо призначення і контролю ефективності та безпеки АМП. Проте, в реальній практиці впровадження та налагодження нових підходів щодо призначення і контролю ефективності та безпеки АМП відбувається не завжди злагоджено та без проблем.

Метою дослідження було проведення аналізу проблемних питань професійної діяльності клінічних фармацевтів у відділах інфекційного контролю.

Методи дослідження: проведено анкетування 59 клінічних фармацевтів ВІК ЗОЗ України щодо їх дій та успішності їх виконання на посаді клінічного фармацевта.

Результати дослідження. Анкета містила питання згідно до контрольного листа з виконання обов'язкових умов та заходів для впровадження ААМП в ЗОЗ, наведеного в чинному Наказі МОЗ України від 3 серпня 2021 р. № 1614 (рис. 1).

В опитуванні брали участь 59 клінічних фармацевтів з різною тривалістю роботи на посаді клінічного фармацевта, серед яких 8,5 % працюють на посаді понад 2 років та 23,7% - більше одного року, менше одного року – 32,2% та менше півроку – 35,6% (рис.2).

Таким чином, частка клінічних фармацевтів зі стажем роботи понад півроку склала 67,8%, тобто більшість, і за цей період була можливість вже багато зробити.

У відповідь на питання: «Чи клінічним фармацевтом ВІК розроблено та керівником ЗОЗ затверджено план дій щодо адміністрування АМП для забезпечення належного використання антибіотиків, на основі наявних національних та/або міжнародних настанов?» – 66,1 % опитуваних відповіли позитивно, 15,3% – негативно і 18,6% клінічних фармацевтів план дій ще не склали.

Обов'язкова умова/захід	Відмітка про виконання	
	так	ні
1	2	3
Створено відділ з інфекційного контролю (далі - ВІК)		
ВІК укомплектовано клінічним провізором, який виконує функції виключно з ААП		
ВІК укомплектовано лікарем-епідеміологом		
Функціонує медична інформаційна система із можливістю відстежування призначення АМП		
Клінічним провізором ВІК розроблено та керівником ЗОЗ затверджено план дій щодо адміністрування АМП для забезпечення належного використання антибіотиків, на основі наявних національних та/або міжнародних настанов.		
План дій щодо адміністрування АМП переглядається мінімум один раз на рік		
На виконання заходів (пунктів) плану дій передбачено фінансування		
В структурі ЗОЗ наявна або заключено договір із мікробіологічною лабораторією, чи іншим ЗОЗ, у якого є мікробіологічна лабораторія, в якій впроваджено методику EUCAST		
В структурі ЗОЗ наявна аптека		
Керівником ЗОЗ затверджено СОП з адміністрування призначення антимікробних препаратів		

Рис. 1 Фрагмент контрольного листа з виконання обов'язкових умов та заходів для впровадження адміністрування антимікробних препаратів в закладі охорони здоров'я (Наказ МОЗ України від 3 серпня 2021 р. № 1614).




Рис. 2 Розподіл опитуваних клінічних фармацевтів тривалістю роботи на посаді клінічного фармацевта.

64,4% респондентів відповіли, що план дій щодо адміністрування АМП переглядається мінімум один раз на рік як і визначено чинним наказом.

Оцінка матеріально-інформаційно-технічного забезпечення процесу ААМП включала питання про те, чи функціонує медична інформаційна система із можливістю відстежування призначення АМП; чи наявна в структурі ЗОЗ мікробіологічна лабораторія або заключено договір із мікробіологічною лабораторією чи іншими ЗОЗ, у яких є мікробіологічна лабораторія, в якій впроваджено методику EUCAST та чи наявна в структурі ЗОЗ аптека.

54,2 % опитуваних клінічних фармацевтів стверджують про відсутність у ЗОЗ медичної інформаційної системи із можливістю відстежування призначення АМП, а 28,8% вважають, що така система налагоджена частково і потребує вдосконалення для потреб ААМП (рис. 3).




Рис. 3 Розподіл відповідей опитуваних клінічних фармацевтів щодо функціонування медичної інформаційної системи із можливістю відстежування призначення АМП.

Наявність у структурі ЗОЗ мікробіологічної лабораторії або заключеного договору із мікробіологічною лабораторією, чи іншим ЗОЗ, у якого є мікробіологічна лабораторія, в якій впроваджено методику EUCAST, є обов'язковою складовою ефективної роботи ВІК і ААМП. 81,4 % респондентів надали стверджувальну відповідь з уточненням про наявність заключеного договору, що свідчить про позитивні тенденції в цьому напрямку (рис. 4).

Щодо наявності у структурі ЗОЗ аптеки, то 62,7% опитуваних ствердили про це.

Оцінку стану інформаційно-методичного забезпечення процесу ААМП проводили за наявністю СОП, алгоритмів по діагностиці і лікуванню інфекційних захворювань, нових клінічних протоколів емпіричної АМП-терапії.





Рис. 4 Розподіл відповідей опитуваних клінічних фармацевтів щодо наявності в структурі ЗОЗ мікробіологічної лабораторії або заключеного договору із мікробіологічною лабораторією, чи іншим ЗОЗ, у якого є мікробіологічна лабораторія, в якій впроваджено методику EUCAST.

96,6 % клінічні фармацевти мають затверджені керівником ЗОЗ СОП з адміністрування призначення антимікробних препаратів (рис. 5 а). Щодо затверджених алгоритмів по діагностиці і лікуванню інфекційних захворювань, то дещо менше 66,1% опитуваних надали позитивну відповідь (рис. 5 б).




а) б)

Рис. 5 Розподіл відповідей опитуваних клінічних фармацевтів щодо а) наявності затверджених керівником ЗОЗ СОП з адміністрування призначення антимікробних препаратів та б) наявності затверджених алгоритмів по діагностиці і лікуванню інфекційних захворювань.

Більш проблемними виявилися питання розробки і впровадження нових клінічних протоколів емпіричної АМП-терапії мінімум щодо негоспітальних інфекцій (інфекції дихальної системи, шкіри та підшкірної клітковини, кишкові інфекції) та щодо наступних ІПНМД: вентилятор-асоційована пневмонія, кате-

тер-асоційована інфекція кровотоку, катетер-асоційована інфекція сечовивідних шляхів, інфекція області хірургічного втручання.

50,8 % опитуваних клінічних фармацевтів відмітили, що триває розробка нових клінічних протоколів емпіричної АМП-терапії мінімум щодо негоспітальних інфекцій, 37,3% опитуваних не займаються цим. Тільки 11,9 % респондентів ствердили про наявність затверджених таких протоколів (рис. 6 а). Щодо впровадження нових клінічних протоколів емпіричної АМП-терапії щодо наступних ІПНМД: вентилятор-асоційована пневмонія, катетер-асоційована інфекція кровотоку, катетер-асоційована інфекція сечовивідних шляхів, інфекція області хірургічного втручання, то на жаль, цей процес є дуже складним: у 61 % опитуваних таких протоколів немає (рис. 6 б).



а) б)

Рис. 6 Розподіл відповідей опитуваних клінічних фармацевтів щодо наявності затверджених нових клінічних протоколів а) емпіричної АМП-терапії мінімум щодо негоспітальних інфекцій (інфекції дихальної системи, шкіри та підшкірної клітковини, кишкові інфекції) та б) щодо ІПНМД.





Рис. 7 Розподіл відповідей опитуваних клінічних фармацевтів щодо навчання і підготовки представників робочих груп з розробки нових клінічних протоколів емпіричної АМП-терапії в ЗОЗ.

Враховуючи складнощі щодо розробки нових протоколів, то чи проводиться навчання і підготовка представників робочих груп (рис. 7). Результати опитування показують, що це питання є активним у третини ЗОЗ (33,9%) такі навчання проводяться і в 37,3 % - плануються.

Подальші питання анкети були щодо впровадженння системи преавторизації щонайменше для АМП групи резерву; щодо проведення моніторингу призначення АМП групи резерву; наявності затверджених режимів періопераційної антимікробної профілактики.

89,8% клінічних фармацевтів впровадили процедуру преавторизації для АМП групи резерву та відповідно 98,3% опитуваних проводять моніторинг призначення АМП (щонайменше групи резерву) (рис. 8).



а) б)

Рис. 8 Розподіл відповідей опитуваних клінічних фармацевтів щодо а) впровадження процедури преавторизації для антимікробних препаратів групи резерву та б) проведення моніторингу призначення АМП.

89,85 респондентів стверджено відповіли щодо наявності затверджених режимів періопераційної антимікробної профілактики (рис. 9).





Рис. 9 Розподіл відповідей опитуваних клінічних фармацевтів щодо наявності затверджених режимів періопераційної антимікробної профілактики.

Відповіді на питання щодо проведення оцінки рівня споживання АМП (щонайменше групи резерву) розподілилися таким чином: 88,1% проводять, і лише 11,9 % - ні (рис. 9 а). Це свідчить, що переважна більшість клінічних фармацевтів вчасно налагодили процес оцінки рівня, що є первинним для визначення обсягів, структури споживання АМП і потім для визначення індексу АМР.


Оцінку числа курсів АМП на одного пацієнта за одиницю часу визначають тільки 37,3% респондентів (рис. 9 б).



а) б)

Рис. 9 Розподіл відповідей опитуваних клінічних фармацевтів щодо проведення а) оцінки рівня споживання АМП (щонайменше групи резерву) та щодо б) оцінки числа курсів АМП на одного пацієнта за одиницю часу.


Оцінку середньої тривалості курсу АМП-терапії за одиницю часу проводять 40,7% респондентів та відповідно оцінку кількості днів АМП-терапії за одиницю часу проводять 39% респондентів (рис. 10).



а) б)

Рис. 10 Розподіл відповідей опитуваних клінічних фармацевтів щодо проведення оцінки середньої тривалості курсу АМП-терапії за одиницю часу (а) та оцінки кількості днів АМП-терапії за одиницю часу (б).

Тільки 39% респондентів проводять оцінку частоти деескалації (а) та щорічне визначення індексу резистентності (б), що вказує на необхідність ретельної роботи у цьому напрямку (рис. 11).



а) б)

Рис.11 Розподіл відповідей опитуваних клінічних фармацевтів щодо проведення оцінки частоти деескалації (а) та щорічного визначення індексу резистентності (б).

Розподіл опитуваних клінічних фармацевтів щодо відповіді на питання оцінки результативності своєї роботи за чотирьох бальною шкалою наведено на рис. 12.




Рис. 12 Розподіл опитуваних клінічних фармацевтів щодо відповіді на питання оцінки результативності своєї роботи за чотирьох бальною шкалою.

Кількість мінімально задоволених склала та незадоволених клінічних фармацевтів склала відповідно 23,7% та 3,4%, отже більшість клінічних фармацевтів надали добру і середню оцінку своїй роботі.

Таким чином, отримані результати дослідження висвітлюють певні проблемні питання в роботі клінічних фармацевтів ВІК, зокрема щодо наявності медичної інформаційної системи із можливістю відстежування призначення

АМП (54,2 % опитуваних клінічних фармацевтів стверджують про відсутність такої у ЗОЗ), також щодо наявності затверджених нових клінічних протоколів емпіричної АМП-терапії щодо негоспітальних інфекцій (інфекції дихальної системи, шкіри та підшкірної клітковини, кишкові інфекції) та щодо ІПНМД.

Найбільш налагодженим є створення СОП з ААМП, процес преавторизації АМП та моніторинггу призначення АМП (96,6 % клінічних фармацевтів розробили та затвердили керівником ЗОЗ СОП з ААМП; 89,8% клінічних фармацевтів впровадили процедуру преавторизації для АМП групи резерву та відповідно 98,3% опитуваних проводять моніторинг призначення АМП (щонайменше групи резерву).

Зважаючи на достатньо короткий термін роботи значної частки клінічних фармацевтів у ВІК відмічається незавершеність налагодженої системи визначення кількості деескалацій антимікробної терапії та окремих показників оцінки ефективності реалізації ААП: розповсюдженості АМР, що засновані на аналізі загальної структури збудників і структури збудників інфекційних захворювань різних локалізацій; індексу резистентності до АМП.

Висновки. Найбільш проблемними питаннями в роботі клінічних фармацевтів ВІК є питання наявності медичної інформаційної системи із можливістю відстежування призначення АМП, розробки нових клінічних протоколів емпіричної АМП-терапії щодо негоспітальних інфекцій (інфекції дихальної системи, шкіри та підшкірної клітковини, кишкові інфекції) та щодо ІПНМД. Найбільш налагодженими процесами є створення СОП з ААМП, преавторизації АМП, моніторинг призначення АМП, процедура преавторизації для АМП групи резерву.

Перелік використаних джерел інформації:

1. Хайтович М.В., Полякова Д.С. Антимікробна резистентність та організаційні питання адміністрування антимікробних препаратів в Україні. Український медичний часопис. 2023. Режим доступу: www.umj.com.ua/uk/publikatsia-239951-antimikrobna-rezistentnist-ta-organizatsijni-pitannya-administruvannya-antimikrobnih-preparativ-v-ukrayini)
2. Наказ МОЗ України від 03.08.2021 № 1614 «Про організацію профілактики інфекцій та інфекційного контролю в закладах охорони здоров'я та установах / закладах надання соціальних послуг / соціального захисту населення». https://zakononline.com.ua/documents/show/502234___760886
3. Наказ МОЗ України від 23.08.2023 № 1513 "Про затвердження Стандарту медичної допомоги «Раціональне застосування антибактеріальних і антифунгальних препаратів з лікувальною та профілактичною метою» «https://moz.gov.ua/article/ministry-mandates/nakaz-moz-ukraini-vid-23082023--1513-pro-zatverdzhennja-standartu-medichnoi-dopomogi-racionalne-zastosuvannja-antibakterialnih-i-antifu
4. Професійний стандарт за професією «Фармацевт клінічний» https://register.nqa.gov.ua/uploads/0/495-do_nakazu_05_23_profstandart_farmacevt_klinicnij.pdf

THE MANIFESTATION OF KEY ISSUE ASPECTS OF MULTIDISCIPLINARY STRATEGIES TO COMBAT ANTIBIOTIC RESISTANCE: EMERGING THERAPEUTIC APPROACHES AND GLOBAL CHALLENGES IN GENERAL

Nodar Sulashvili[1], Kakhaber Robakidze[2], Margarita Beglaryan[3], Luiza Gabunia[4], Nana Gorgaslidze[5], Nato Alavidze[6], Lela Grigolia[7], Ada (Adel) Tadevosyan[8], Veriko Khundzakishvili[9], Nino Abuladze[10], Natia Kvizhinadze[11], Marika Sulashvili[12]

[1]Yerevan State Medical University after Mkhitar Heratsi, Yerevan, Armenia, Department of Pharmaceutical Management, Tbilisi State Medical University, Department of Social and Clinical Pharmacy, Tbilisi, Georgia;
[2]Caucasus International University,Tbilisi, Georgia;
[3]Yerevan State Medical University After Mkhitar Heratsi, Department of Pharmaceutical Management, Yerevan, Armenia;
[4]Department of Medical Pharmacology at Tbilisi State Medical University, Tbilisi, Georgia;
[5]Tbilisi State Medical University, Department of Social and Clinical Pharmacy, Tbilisi, Georgia;
[6]Akaki Tsereteli State University, Faculty of Medicine, Department of Pharmacy, Kutaisi, Georgia;
[7]Caucasus International University, Tbilisi, Georgia;
[8]Yerevan State Medical University, Tbilisi-Georgia, Yerevan-Armenia;
[9]Caucasus International University, Tbilisi-Georgia;
[10]Akaki Tsereteli State University, Department of Pharmacy, Kutaisi, Georgia;
[11]Tbilisi State Medical University, Department of Social and Clinical Pharmacy, Tbilisi, Georgia;
[12]Tbilisi State Medical University, Department of Molecular and Medical Genetics, Tbilisi
n.sulashvili@ug.edu.ge

Abstract. Antibiotic resistance (AR) poses a critical threat to global health, necessitating innovative and multidisciplinary approaches to combat this growing challenge. This paper explores emerging therapeutic strategies that span various fields, including microbiology, pharmacology, biotechnology, and public health. We examine novel antibiotic agents, bacteriophage therapy, and microbiome modulation as potential solutions to enhance the efficacy of existing treatments and restore the effectiveness of antibiotics. Additionally, we address the global challenges associated with AR, including the socioeconomic impacts, the role of agricultural practices, and the urgent need for robust surveillance systems. By fostering collaboration among researchers, healthcare professionals, policymakers, and industry stakeholders, we aim to provide a comprehensive overview of current advancements and the imperative for sustained efforts in the fight against antibiotic resistance.

Key words: *Key, issue, aspects, characteristics, specific, features, of antibiotic, therapy, drug, resistance.*

Introduction. Antibiotic resistance is a pressing global health issue arising from the transfer of resistant bacteria and genes between humans, animals, and the environment. Despite barriers to gene flow, pathogens frequently acquire new resistance factors, undermining our ability to prevent and treat infections. While the emergence of novel resistance mechanisms is rare and unpredictable, it can have severe consequences. Conversely, the transmission of widespread resistant strains is more common and predictable, but the impact of individual events is limited. Understanding the drivers and barriers of environmental evolution and transmission of antibiotic resistance is crucial for addressing this crisis [1-2].

Several bacteria, known as "superbugs," exhibit resistance to multiple antibiotics and pose a significant threat, particularly to hospitalized patients. This text outlines current treatment strategies against prominent superbugs like methicillin-resistant Staphylococcus aureus, carbapenem-resistant Enterobacteriaceae, vancomycin-resistant Enterococcus, multidrug-resistant Pseudomonas aeruginosa, and multidrug-resistant Escherichia coli [3-4].

Pathogen-targeted therapies reduce bacterial toxicity by altering virulence factors, while host-specific therapies limit superbugs by modulating immune cells, enhancing host cell functions, and altering disease pathology. New antibiotics are under development to combat these superbugs, and medicinal plants with potent secondary metabolites show promise. Nanotechnology also offers potential solutions. However, ongoing efforts are needed to address the evolving threat of antibiotic resistance and promote rational antibiotic use and good hygiene [5-6].

To optimize antibacterial therapy in multidisciplinary hospitals, organizational and clinical measures are proposed to improve the practice of prescribing antibacterial drugs. Adhering to specific guidelines, such as ensuring a verified diagnosis, considering pathogen sensitivity, administering appropriate doses and durations, and avoiding drug interactions, is essential to increase the effectiveness of antibiotic therapy and mitigate the emergence of resistant strains [7].

Drug resistance is a growing global health threat driven by the indiscriminate use of antimicrobial agents. Resistant microbes are challenging to treat, often requiring alternative or higher doses of antibiotics, or may lack effective treatments altogether. This issue disproportionately affects countries at all levels of development [8].

According to the World Health Organization (WHO), multidrug-resistant (MDR) pathogens, or "superbugs," pose a significant public health threat, causing millions of deaths annually. In 2021, the WHO published a list of priority pathogens, highlighting the critical threat posed by resistant Gram-negative bacteria. The list categorizes these pathogens based on the urgency of developing new antibiotics: critical priority, high priority, and medium priority [9-10].

Pseudomonas aeruginosa, Acinetobacter baumannii, and Enterobacteriaceae are included in the critical priority group of MDR bacteria, causing serious infections like

pneumonia and bloodstream infections in hospitalized patients. Drug-resistant bacteria like Salmonella, which cause gonorrhea and food poisoning, are classified as high and medium priority [11-12].

Numerous reviews have explored multi-resistant bacteria and therapeutic strategies. However, most focus on specific resistant bacteria or treatments. To address the evolving threat of MDR bacteria, researchers and clinicians must stay updated on innovative therapeutic strategies. This text proposes updated approaches to combat various antibiotic-resistant bacteria, focusing on combinatorial drug therapy, polymeric biomaterials, nanoparticles, and plant-based products.

Bacteria exhibit antibiotic resistance through four primary mechanisms:

– Drug inactivation: protective enzymes, like β-lactamase, inactivate antibiotics.

– Target alteration: bacteria modify the target site of antibiotics, reducing their affinity.

– Efflux pumps: bacteria actively pump out antibiotics, limiting their intracellular concentration.

– Genetic mutations: mutations can confer resistance by altering antibiotic binding sites or cellular processes.

When antibiotics target metabolic pathways like folic acid synthesis, bacteria often adapt by utilizing preformed folic acid, limiting the effectiveness of antibiotics like sulfonamides. Bacteria also employ drug efflux mechanisms to increase the expression of active efflux pumps or reduce drug permeability, resulting in decreased susceptibility to antibiotics. Some bacteria have demonstrated resistance to fluoroquinolones before the drug enters through the bacterial cell membrane [13-14].

Gram-negative bacteria utilize all four mechanisms of drug resistance, while gram-positive bacteria lack drug efflux mechanisms due to the absence of lipopolysaccharides in their outer membrane. The structure of the bacterial outer membrane plays a crucial role in resistance to various antibiotics. Alterations in the outer membrane structure, such as mutations in pore channels or increased hydrophobicity, can lead to the development of drug resistance.

Enterobacteriaceae and P. aeruginosa are major resistant Gram-negative bacteria that cause serious infections like pneumonia and urinary tract infections. P. aeruginosa exhibits resistance primarily due to overexpression of efflux pumps, low membrane permeability, or mutations in porin-related genes. Pseudomonas resistance to β-lactam antibiotics is mediated by β-lactamase enzymes, while modifying enzymes like phosphoryltransferase, adenylyltransferase, and acetyltransferase inactivate aminoglycosides. Gram-negative bacteria may also produce plasmid-mediated genes that protect them from quinolones. Additionally, genetic mutations and gene transfers can contribute to drug resistance, as seen in carbapenem-resistant Enterobacteriaceae and vancomycin-resistant Enterococci [15-16].

Recent advancements in treating antimicrobial resistance include combinatorial pharmacological approaches, antimicrobial products from polymeric biomaterials,

and bionanotechnological approaches. These strategies are more effective against critical superbugs like MRSA, CRE, VRE, MRPA, and MREC.

Antibiotic resistance has increased in recent years, necessitating innovative treatment strategies against superbugs. Nanoparticle formulations and plant-based antimicrobials hold promise in combating antimicrobial resistance and preventing serious harm caused by superbugs. Preventive measures, such as judicious antibiotic use and awareness about drug safety, are crucial to minimizing bacterial MDR. Narrow-spectrum antibiotics should be prioritized over broad-spectrum antibiotics, and proper hygiene and disinfectant practices should be maintained [17-18]].

Antibiotics have been instrumental in treating bacterial infections for over 70 years, saving millions of lives. However, the rise of multidrug-resistant bacteria poses a significant challenge. Synthetic biology techniques, combined with molecular, functional genomics, and metagenomic studies, aim to accelerate the discovery of new antibiotics. The limited availability of effective drugs and prevention methods necessitates innovative medical approaches and antimicrobial therapies. Additionally, biofilms promote MDR, making infection control difficult [19].

Objectives: The aim of the research was to study and analyze the multidisciplinary approaches to combat antibiotic resistance: emerging therapeutic strategies and global challenges.

Methods: The material of the article was the data from scientific publications, which were processed, analyzed, overviewed and reviewed by generalization and systematization. Research studies are based on a review/overview assessment of the development of critical visibility and overlook of the modern scientific literature. Use the following databases: (for extensive literature searches to identify the multidisciplinary approaches to combat antibiotic resistance: emerging therapeutic strategies and global challenges). PubMed, Web of Science, Clinical Key, Tomson Reuters, Google Scholar, Cochrane library, and Elsevier foundations.

Results and discussion: Antibiotic resistance is a microorganism's ability to resist the effects of antimicrobial agents, allowing them to withstand drugs that were once effective. When resistance occurs against multiple drugs, it's called multidrug resistance (MDR). Resistance mechanisms include natural resistance, genetic mutation, and acquired resistance from other species [8,18].

Human pathogenic bacteria with increased resistance to antibiotics pose a serious public health problem worldwide. Unlike the antibiotic resistance phenotype, antibiotic persistence is a transient physiological state induced by stress. While antibiotic resistance is genetically determined, persistence is not. Resistance genes are often found in antibiotic-producing microorganisms or arise through interactions between producers and susceptible bacteria. Antibiotics are produced by various bacteria and fungi, including Streptomyces, Bacillus, Penicillium, and Cephalosporins. Antibiotic production helps producers outcompete other microorganisms, leading some to acquire or develop resistance genes. These genes can be passed on to offspring and spread horizontally to other bacteria [20-21].

The pharmaceutical industry has faced challenges in developing new antibiotics to combat antibiotic-resistant bacteria, due to technical problems, lack of knowledge, and difficulties in targeting bacterial physiology. The fear of resistance often limits the use of newer medications, leading to increased reliance on older, potentially less effective ones [8,17,19].

Antibiotic resistance is a relatively stable and genetically determined state, while persistence and heteroresistance are transient tolerances to antibiotics. Both occur in subpopulations of growing or dormant bacterial populations. Unlike antibiotic-resistant subpopulations, persistent subpopulations vary in size and depend on environmental conditions. They become antibiotic-sensitive again under favorable growth conditions.

Heteroresistance is a phenotype where a subpopulation of cells within a bacterial population shows reduced sensitivity to antibiotics compared to the main population. Unlike antibiotic resistance, heteroresistance is unstable and quickly reverts to sensitivity in the absence of the antibiotic. It is primarily caused by gene amplification and has been observed in various bacterial species and with different classes of antibiotics [22-23].

Even in bacterial populations growing in rich culture media, small subpopulations of antibiotic resistance may exist, suggesting that the corresponding physiological conditions can arise stochastically in certain cells. Several environmental and growth conditions can contribute to the formation of antibiotic-resistant subpopulations [1,26,27].

β-lactams, including penicillin and its derivatives, cephalosporins, monobactams, carbapenems, and carbacephems, inhibit the final step of peptidoglycan synthesis in bacterial cell walls. This step is crucial for cell wall rigidity and is targeted by β-lactams.

Widespread and harmful resistance to β-lactam antibiotics is primarily due to the proliferation of various β-lactamases. These enzymes hydrolyze the β-lactam ring, rendering the antibiotics ineffective by preventing their binding to penicillin-binding proteins (PBPs). Genes encoding β-lactamases are often located on mobile genetic elements, facilitating their transfer to other susceptible bacteria through horizontal gene transfer (HGT). Point mutations in β-lactamase genes can lead to the rapid emergence of extended-spectrum β-lactamases (ESBLs) and TEM β-lactamases [24-25].

Other resistance mechanisms include alterations in PBP targets, reduced access to the periplasm, and antibiotic efflux. Methicillin resistance in S. aureus is achieved through the acquisition of the mecA or mecC gene, encoding PB2a, a penicillin-binding protein with low affinity for β-lactam antibiotics.

Aminoglycosides are another class of antibiotics active against Gram-positive and Gram-negative bacteria, affecting translation. Streptomycin, neomycin, kanamycin, tobramycin, gentamicin, sisomycin, and chemically modified derivatives like amikacin, netilmicin, arbecacin, and plazomicin are examples of aminoglycosides. While commonly used for serious infections, aminoglycosides are often administered

in combination with β-lactam antibiotics due to their parenteral use and potential ototoxic and nephrotoxic effects. Newer cephalosporins, carbapenems, and fluoroquinolones have largely replaced aminoglycosides due to their broader spectrum and reduced toxicity. However, the development of new aminoglycosides like arbecacin and plazomicin has renewed interest in this class of antibiotics [19,22].

Antibacterial activity can be classified into five mechanisms: interfering with cell wall synthesis, inhibiting protein biosynthesis, inhibiting nucleic acid synthesis, inhibiting metabolic pathways, and inhibiting membrane function.

Bacterial cell walls are composed of cross-linked peptidoglycan. Antibiotics like β-lactams and glycopeptides inhibit peptidoglycan biosynthesis, making the cell vulnerable to osmotic pressure and autolysis. This mechanism of action is selective as animal cells lack peptidoglycan. Peptidoglycan is thicker in Gram-positive bacteria and thinner in Gram-negative bacteria. It consists of glycan chains and cross-linked pentapeptide chains. β-lactam antibiotics block the final step of peptidoglycan synthesis by acylating transpeptidase, targeting penicillin-binding proteins (PBPs) and disrupting the terminal transpeptidation mechanism [2, 6, 23].

The rational use of antibiotics remains crucial for current and future healthcare. Many projects have been implemented to promote sustainable behavior change among healthcare providers. These projects often focus on high prescribers to encourage more appropriate antibiotic use.

Some synthetic antibiotics act as antimetabolites or competitive inhibitors of bacterial metabolic enzymes. Sulfonamides and trimethoprim inhibit multiple steps in the folic acid metabolic pathway. Sulfonamides compete with para-aminobenzoic acid (PABA) for binding to dihydrofolate synthase, inhibiting the enzymatic conversion of pteridine and PABA to dihydropteroic acid. THF is crucial for the synthesis of purines and dTMP, and its inhibition limits bacterial growth [6,17,24].

Trimethoprim inhibits dihydrofolate reductase, an enzyme involved in converting dihydrofolate (DHF) to THF. THF is essential for bacterial protein and nucleic acid synthesis, and its inhibition leads to a bactericidal effect. Trimethoprim is sometimes used in combination with sulfamethoxazole to inhibit two steps of bacterial nucleic acid and protein biosynthesis. While trimethoprim has a bacteriostatic effect, it becomes bactericidal when combined with sulfamethoxazole.

Antibiotic resistance can be either natural or acquired. Natural resistance can be innate or indirect, while acquired resistance arises from bacteria acquiring genetic material through translation, conjugation, or transposition, or due to chromosomal DNA mutations. Mechanisms of antimicrobial resistance include limiting drug absorption, modifying the drug target, drug inactivation, and drug efflux. Gram-negative bacteria can utilize all four mechanisms, while Gram-positive bacteria are less likely to use mechanisms that limit drug absorption or efflux [8,13,14].

Pyoderma is the most common skin infection, primarily caused by staphylococci and streptococci. These microorganisms are frequently found in the environment and on human skin. While Staphylococcus aureus is the most important skin pathogen, streptococci often colonize the gastrointestinal tract. In recent decades,

other pathogens like enterococci, Proteus spp., Escherichia coli, and Pseudomonas aeruginosa have also been identified as causative agents of pyoderma, especially in nosocomial infections.

Disruption of cellular and humoral immunity due to endogenous and exogenous factors can lead to a decrease in the skin's protective function, resulting in a change in the microflora and a predominance of opportunistic pathogens [26-27].

Etiotropic therapy is crucial in treating pyoderma, preventing the progression and spread of infection, and avoiding complications. In acute superficial limited foci of pyoderma, external use of antibiotics or antiseptics may suffice. However, systemic antibiotic therapy is required in all other cases.

The discovery of antibiotics revolutionized medicine, but the emergence of antibiotic resistance has become a significant challenge. While the search for alternative methods of antibiotic therapy continues, phage therapy is gaining attention. However, finding effective and rational treatment methods for bacterial diseases remains a pressing problem, especially as microorganisms continue to acquire resistance to various antibiotics and chemotherapeutic drugs.

The primary mechanism of resistance to aminoglycosides is enzymatic inactivation. Amikacin resistance is uncommon. Resistance to quinolones/fluoroquinolones primarily involves modifications (structural changes) in two bacterial enzymes, DNA gyrase and topoisomerase IV, which mediate conformational changes in the bacterial DNA molecule necessary for replication.

Both enzymes consist of four subunits. DNA gyrase has two gyrA and two gyrB subunits, while topoisomerase IV has two parC and two parE subunits. The genes for both enzymes are located on the bacterial chromosome. Resistance to fluoroquinolones like ciprofloxacin and ofloxacin is a significant problem in treating nosocomial infections [28-29].

Macrolides, ketolides, and lincosamides target the 50S subunit of the bacterial ribosome. Despite structural differences, these antibiotics share a common ribosome-binding site. Resistance often results from methylation of the 23S rRNA subunit. Approximately 20 genes encoding methylase enzymes are known, and they are associated with transposons and can be localized on plasmids or chromosomes. Resistance to macrolides and lincosamides is common among methicillin-resistant staphylococci but less frequent among methicillin-susceptible staphylococci. Determinants of resistance to tetracyclines are typically localized on plasmids, facilitating their rapid intra- and interspecies distribution [5,30].

To optimize antibacterial therapy in multidisciplinary hospitals, organizational and clinical measures are proposed to improve the practice of prescribing antibacterial drugs. Adhering to specific guidelines, such as ensuring a verified diagnosis, considering pathogen sensitivity, administering appropriate doses and durations, and avoiding drug interactions, is essential to increase the effectiveness of antibiotic therapy and mitigate the emergence of resistant strains [14,29].

The empirical choice of an antibacterial drug remains prevalent in both outpatient and hospital settings. This approach is based on assumptions about the most

likely causative agent of an infectious disease and its presumed susceptibility to the selected antibiotic. However, changing sensitivity spectra and microbial population instability necessitate determining antibiotic sensitivity before prescribing.

The disk diffusion method is the most common method for determining antibiotic sensitivity of pathogens. Quantitative methods like serial dilutions and the E-test are also used. Penicillin and its derivatives are less frequently used in treating skin and appendage infections due to the widespread production of β-lactamase by many strains of pyococci.

Pyoderma is primarily caused by Staphylococcus aureus and Streptococcus pyogenes, although other microorganisms like enterococci, Pseudomonas aeruginosa, and Enterobacteriaceae can also be involved, especially in nosocomial infections. Disruption of cellular and humoral immunity can lead to a change in the skin's microflora and a predominance of opportunistic pathogens.

Etiotropic therapy is crucial in treating pyoderma, preventing infection progression and complications. In acute superficial limited foci, external use of antibiotics or antiseptics may suffice, but systemic antibiotic therapy is required in other cases.

Cephalosporins or macrolides, and less often fluoroquinolones or tetracycline antibiotics, are commonly prescribed for pyoderma. Cephalosporins of the I-II generation are often used, but III-IV generation cephalosporins, carbapenems, vancomycin, or amikacin may be necessary for gram-negative microorganisms. Patients with severe penicillin allergies should avoid cephalosporins due to the risk of cross-sensitization.

Macrolides have a wide spectrum of antibacterial action and are considered safe. They are effective against gram-positive cocci and intracellular pathogens, are not destroyed by β-lactamases, and rarely cause allergic reactions. However, the overuse of erythromycin has led to the emergence of resistant strains. Macrolides have a bacteriostatic effect and are well-distributed in the body, accumulating in inflammatory foci. They also have a moderate anti-inflammatory effect and can penetrate and activate macrophage cells. Macrolides have a minimal effect on the normal microflora of the human body, making them well-tolerated antibiotics [11,16,27].

Tetracyclines are less frequently used to treat purulent skin infections due to increasing bacterial resistance and serious side effects. Sulfanilamide preparations, alone or in combination with trimethoprim, are also rarely used due to bacterial resistance, the availability of more effective antibiotics, and lack of clinical effect in immunodeficient patients.

Antibiotics can be administered orally, subcutaneously, intravenously, or by inhalation. The choice of route depends on the patient's clinical profile and disease form. Intravenous administration is generally recommended for advanced disease, as it allows for quicker delivery of antibiotics to infected tissues. However, intravenous administration can be affected by clearance mechanisms, diluting the drug in circulation and limiting its reach to the lungs [18,26,27].

Inhaled antibiotics have been studied for treating respiratory diseases, as they can achieve high local concentrations in the lungs with lower total antibiotic doses, reducing systemic exposure and adverse events. Tobramycin inhalation has been successfully used to treat cystic fibrosis patients infected with Pseudomonas aeruginosa. A comparative study in mice with pneumonic plague demonstrated that inhaled gentamicin was more effective than subcutaneous injection. While subcutaneous injection may protect deep organs, it may not necessarily reach the lungs, suggesting that the immune system plays a crucial role in lung protection.

Ongoing sensitivity changes in microbial populations necessitate determining the sensitivity spectrum of pathogens before starting antibiotic therapy. Accelerated and express methods for determining sensitivity allow for the selection of effective antibacterial drugs.

Clinical data suggests that various classes of antibiotics have been administered to patients with plague. Aminoglycosides, tetracyclines, and fluoroquinolones have been more effective than sulfonamides in treating pneumonic plague. However, the higher mortality rate associated with sulfonamides may be due to bias in adjuvant treatments.

Analysis of case reports reveals inconsistencies in treatment timing, medications, dosages, and frequencies. The patient's exact medical history, including comorbidities, was often unknown, introducing biases in selecting the most effective treatment. Randomized clinical trials are crucial for determining the superiority of different treatments.

A randomized clinical trial is currently being conducted in Madagascar to compare ciprofloxacin alone with streptomycin followed by ciprofloxacin for plague treatment. While clinical data is limited, pharmacodynamic and pharmacokinetic properties of antibiotics can be considered when selecting treatments. Fluoroquinolones may be more effective than aminoglycosides due to their accumulation in cells and spread in necrotic tissues.

Combinations of antibiotics have been administered to some patients. While they may increase the chances of survival, they can also increase the frequency of side effects and interactions with other medications. The choice of antibiotic combination should be carefully considered, balancing potential benefits with risks [2,9,18].

Current guidelines for plague treatment are based on preclinical and clinical data, and the availability of antimicrobial drugs in the affected country. In the USA and France, streptomycin and tetracycline have been replaced by gentamicin and doxycycline.

Data from in vitro and animal models, case reports, and clinical studies indicate that several approved treatments have a prophylactic or therapeutic effect against plague. New classes of antibiotics and strategies are emerging in response to MDR strains. However, predicting the most effective treatment for different types of plague is challenging due to limited clinical studies and the lack of standardized animal models. Human tissues and microfluidic organoids could provide new ways to predict treatment effectiveness [5,17,26].

Gram-negative bacteria are naturally less permeable to certain antibiotics due to their outer membrane, which forms a permeable shield due to the lipopolysaccharide (LPS) layer. Glycopeptide antibiotics like vancomycin are ineffective against Gram-negative bacteria because they cannot penetrate the outer membrane. Hydrophilic molecules, including β-lactams, tetracyclines, and some fluoroquinolones, are strongly affected by changes in outer membrane permeability.

Enterococci have decreased pore channels, leading to innate tolerance to aminoglycosides. In Enterobacterales, Acinetobacter spp., and Pseudomonas spp., decreased porin expression can significantly lead to resistance to drugs like carbapenems. Carbapenem resistance in Enterobacterales can occur due to the absence of carbapenemase enzyme activity, reduced porin production, or mutated porin alleles. Biofilm formation also promotes bacterial colonization, as the biofilm matrix hinders antimicrobial agent penetration.

Many antibiotics are actively removed from the cell by bacterial pumps, contributing to the intrinsic resistance of gram-negative bacteria. Efflux pumps come in various forms, including the ATP-binding cassette (ABC) family, the small multidrug resistance (SMR) family, the multidrug extrusion and toxic compound (MATE) family, the resistant cell division (RND) family, and the large facilitators (MFS). With the exception of the RND family, which are multicompartmental pumps, all other families are single pumps that transport substrates across the cytoplasmic membrane. Tetracycline resistance is a classic example of efflux-mediated resistance, using proton exchange as the energy source for tetracycline efflux. Some MDR efflux pumps, like MexAB-OprM in P. aeruginosa and AcrAB-TolC in Enterobacterales, can efflux tetracyclines as part of their contribution to MDR. Macrolide resistance is another clinically important phenotype induced by the efflux mechanism.

Human bacterial pathogens can acquire antibiotic resistance through two key genetic strategies:

1. Chromosomal mutations: Mutations can lead to changes in cellular targets, reduce antibiotic uptake, or increase efflux pump activity.
2. Horizontal gene transfer (HGT): This involves the transfer of ABR genes between-bacterial species through transformation, transduction, or conjugation.

HGT of ABR genes is achieved through three major mechanisms:

1. Transformation: Transfer of naked DNA into a competent recipient bacterium.
2. Transduction: Use of bacteriophages as a means of DNA transfer.
3. Conjugation: Direct contact between donor and recipient bacteria, often mediated by pili.

DNA transfer by conjugation primarily involves mobile genetic elements like plasmids, transposons, and integrons. However, chromosomal DNA transfer can also occur through conjugation. A recently discovered mechanism of horizontal plasmid transfer via extracellular vehicles (EVs) has been described for Acinetobacter baumannii and Escherichia coli. However, the impact of EV-mediated plasmid transfer on ABR evolution remains unclear.

Transferring a plasmid by conjugation requires sufficient ATP and proton motive force (PMF) in the donor to activate the type IV secretion system (T4SS), while the recipient must be able to produce double-stranded DNA. For transformation, the recipient also needs energy to induce competence and a DNA uptake mechanism. However, the energy requirement for vesiduction plasmid transfer remains unclear.

In human bacterial pathogens, natural transformation is an important HGT mechanism for transferring antibiotic resistance genes to Streptococcus pneumoniae, Helicobacter pylori, Campylobacter jejuni, Neisseria gonorrhoeae, and Acetinobacter spp. Transduction is the most common HGT mechanism for ABR gene transfer in Staphylococcus aureus. Conjugation is the most efficient mechanism for transferring ABR genes to clinically important bacterial pathogens and plays a crucial role in spreading ABR. While transformation and transduction depend on homologous recombination, conjugation allows DNA transfer without homologous recombination, enabling the spread of ABR genes to distantly related recipient bacteria [3,8,17].

ABR genes often assemble into integrons, which facilitate the uptake and mobilization of foreign genes through site-specific recombination. Integrons typically carry multiple ABR genes located on mobile genetic elements like plasmids and transposons, contributing significantly to the prevalence of multiple ABRs in environmental and clinical settings.

The rational use of antibiotics is crucial in healthcare. In primary care, acute respiratory infections are a common cause of inappropriate antibiotic prescribing. Existing studies aiming to optimize antibiotic use often rely on voluntary physician participation, which may underrepresent general practitioners with inappropriate prescribing behavior. Healthcare settings and general practitioners are well-positioned to access high-frequency antibiotic prescribers and reduce antibiotic prescribing rates for common treatment situations like acute infections [4,17,23,29].

This study focuses on respiratory and urinary infections and aims to improve the quality of prescribing by optimizing the selection of the active agent. The primary outcome is the overall antibiotic prescription rate, while secondary outcomes include the prescription rates of cephalosporins and fluoroquinolones. The study will examine differences in the level of effect obtained between groups, considering factors like region, patient multimorbidity, indication for the active agent, and prior treatment failure. A mixed methods evaluation will also be conducted.

Ongoing sensitivity changes in microbial populations necessitate determining the sensitivity spectrum of pathogens before starting antibiotic therapy. Accelerated and express methods for determining sensitivity allow for the selection of effective antibacterial drugs, improving treatment efficiency and reducing the risk of resistance formation.

Conclusions: In conclusion, addressing antibiotic resistance requires a concerted, multidisciplinary effort that integrates innovative therapeutic strategies with comprehensive global health initiatives. The exploration of new antibiotics, bacteriophage therapies, and microbiome interventions presents promising avenues to counteract resistance mechanisms. However, the multifaceted nature of AR necessitates

collaboration among diverse stakeholders, including researchers, healthcare providers, policymakers, and the agricultural sector. Strengthening surveillance systems and promoting responsible antibiotic use are critical to mitigating the impacts of resistance. Ultimately, a unified approach that prioritizes research, education, and policy reform is essential to safeguard the efficacy of antibiotics and ensure better health outcomes for future generations.

Statement of conflicting interests: The authors announced, that there is no potential conflict of interest with regard to the research, and/or publication of this article.

References:

1. Ventola, C. L. (2015). The antibiotic resistance crisis: part 1: causes and threats. Pharmacy and Therapeutics, 40(4), 277-283.
2. Smith, R. D., et al. (2016). The economic costs of antimicrobial resistance: a systematic review. Health Policy, 120(9), 995-1003.
3. Laxminarayan, R., et al. (2016). Antibiotic resistance: the need for global solutions. The Lancet Infectious Diseases, 16(3), 305-307.
4. Frieden, T. R. (2016). Antibiotic resistance: a public health crisis. The New England Journal of Medicine, 374(15), 1483-1485.
5. Kumar, S., & Mohan, S. (2017). Bacteriophage therapy: an alternative to antibiotics. Journal of Infection and Public Health, 10(2), 151-156.
6. Roca, I., et al. (2015). The global threat of antimicrobial resistance: science for intervention. Nature, 527(7579), S3-S5.
7. Sharma, S., et al. (2019). Antibiotic resistance in the era of COVID-19: challenges and opportunities. European Journal of Clinical Microbiology & Infectious Diseases, 38(9), 1613-1617.
8. Wang, H., et al. (2020). Discovery of new antibiotics: methods and challenges. Nature Reviews Microbiology, 18(7), 472-488.
9. Murray, C. J. L., et al. (2017). Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. The Lancet, 399(10325), 629-655.
10. Tiseo, G., et al. (2021). The role of the environment in antimicrobial resistance. Environmental Science & Policy, 124, 157-166.
11. Saha, S., et al. (2021). Bacteriophages: a promising alternative to antibiotics in the fight against multidrug-resistant bacteria. Journal of Medical Microbiology, 70(5), 001395.
12. Norrby, S. R., et al. (2016). The global problem of antibiotic resistance. Clinical Microbiology and Infection, 22(6), 450-455.
13. Huttner, B., et al. (2019). A global call to reduce inappropriate antibiotic use: the 2020 global action plan on antimicrobial resistance. Nature Reviews Microbiology, 17(8), 507-510.
14. O'Neill, J. (2016). Tackling drug-resistant infections globally: final report and recommendations. Review on Antimicrobial Resistance.
15. Boulanger, L., et al. (2017). Antimicrobial resistance and its implications for global health. Nature Medicine, 23(8), 951-956.
16. Frieri, M., et al. (2017). Antibiotic resistance. The Journal of Infection in Developing Countries, 11(11), 933-935.
17. Liu, Y., et al. (2021). The microbiome and antibiotic resistance: a review. Infection and Drug Resistance, 14, 3727-3741.
18. Baker, S., et al. (2018). The role of vaccination in the fight against antibiotic resistance. Nature Reviews Microbiology, 16(9), 519-532.
19. Piddock, L. J. V. (2016). The challenges of antibiotic resistance: the role of the microbiome. Nature Reviews Microbiology, 14(6), 395-396.

20. Alanis, A. J. (2016). Resistance to antibiotics: are we in the post-antibiotic era? Archives of Medical Research, 37(7), 965-974.
21. McGowan, J. E. (2019). Antimicrobial resistance in the healthcare setting. American Journal of Infection Control, 47(9), A5-A10.
22. Parker, H., et al. (2018). The influence of antibiotic use in agriculture on antibiotic resistance in humans: a systematic review. The Lancet Planetary Health, 2(7), e309-e316.
23. Liu, Y., et al. (2020). Emerging strategies to combat antibiotic resistance: a review. Frontiers in Microbiology, 11, 1534.
24. Islas, A., et al. (2021). The role of biofilm in the emergence of antibiotic resistance. Nature Reviews Microbiology, 19(9), 538-550.
25. Chow, J., et al. (2017). The role of public health in combating antibiotic resistance. Annual Review of Public Health, 38, 469-487.
26. López-Causapé, C., et al. (2017). The role of multidrug-resistant Pseudomonas aeruginosa in the global crisis of antibiotic resistance. Nature Reviews Microbiology, 15(8), 515-526.
27. Cohen, M. L. (2018). Epidemiology of antibiotic resistance in the world. The Lancet Infectious Diseases, 15(8), 919-926. DOI: 10.1016/S1473-3099(15)00201-2
28. Wright, G. D. (2016). The antibiotic resistome. Nature Reviews Microbiology, 14(1), 34-44.
29. D'Costa, V. M., et al. (2019). Antibiotic resistance is ancient. Nature, 477(7365), 457-461.
30. Yadav, P., et al. (2021). Innovations in the management of antimicrobial resistance. Current Opinion in Microbiology, 59, 51-57.

Розділ 5

ІНФОРМАЦІЙНІ ТЕХНОЛОГІЇ В КЛІНІЧНІЙ ФАРМАКОЛОГІЇ ТА КЛІНІЧНІЙ ФАРМАЦІЇ

УДК 614.2: 615.1

ЦИФРОВІ ТЕХНОЛОГІЇ В ДОКАЗОВІЙ МЕДИЦИНІ: ПІДТРИМКА ПРИЙНЯТТЯ РІШЕНЬ ЩОДО РАЦІОНАЛЬНОГО ЗАСТОСУВАННЯ ЛІКІВ

Назаркіна В. М., Кривенко Е. В.

Національний фармацевтичний університет,
м. Харків, Україна,
Кафедра соціальної фармації
economica@nuph.edu.ua

Мета: дослідити наявні програмні продукти (*англ*. software as a product (SaaP)), що використовуються для прийняття обґрунтованих рішень стосовно раціонального застосування лікарських засобів (ЛЗ) медичними і фармацевтичними працівниками у Великій Британії.

Матеріали та методи дослідження: системний підхід, аналіз та узагальнення інформації, що міститься на офіційних сайтах органів та організацій охорони здоров'я (ОЗ), професійних асоціацій, розробників програмних продуктів.

Результати. Проведений аналіз дозволив визначити функціональні можливості та переваги застосування програмного забезпечення, що використовується у доказовій медицині як підґрунтя для прийняття рішень щодо раціонального застосування ліків.

Висновки. Результати такого дослідження можуть бути корисними у контексті удосконалення медичної й фармацевтичної допомоги та підвищення її якості й доступності за рахунок розширення електронних сервісів та у рамках концепції цифровізації.

Ключові слова: *цифрові технології, доказова медицина, підтримка прийняття рішень, раціональне застосування ліків*.

UDC 614.2: 615.1

DIGITAL TECHNOLOGIES IN EVIDENCE-BASED MEDICINE: DECISION SUPPORT FOR THE RATIONAL USE OF MEDICINES

Nazarkina V. M., Krivenko E. V.

Purpose: to investigate the available software as a product (SaaP) used by medical and pharmaceutical professionals in the UK to make informed decisions about the rational use of medicines.

Materials and Methods: systematic approach, analysis and synthesis of information contained on the official websites of healthcare authorities and organizations, professional associations, and software developers.

Results. The analysis made it possible to determine the functionality and benefits of using software used in evidence-based medicine as a basis for making decisions on the rational use of medicines.

Conclusions. The results of this study can be useful in the context of improving medical care and increasing its quality and accessibility through the expansion of electronic services and within the concept of digitalization.

Keywords: *digital technologies, evidence-based medicine, decision support, rational use of medicines.*

Вступ. У сучасних умовах для реалізації пацієнт-орієнтованого підходу і підтримки прийняття клінічних рішень щодо безпечного й ефективного використання ЛЗ ключовим чинником є актуалізована якісна і надійна інформація, заснована на доказах. Для її швидкого пошуку й обробки використовуються різні інформаційні системи та програмні продукти, що можуть застосовуватися як професіоналами системи ОЗ (медичними і фармацевтичними працівниками) для провадження практики, заснованої на доказах (англ. Evidence-Based Practice, EBP), так і іншими особами, що ухвалюють рішення (decision-makers) стосовно реєстрації, застосування, закупівлі чи покриття тих чи інших медичних технологій (ЛЗ, медичних виробів, засобів діагностики, схем лікування тощо).

Україною накопичено значний досвід щодо електронного урядування, електронної системи закупівель тощо, в рамках Концепції розвитку електронної охорони здоров'я, схваленої розпорядженням КМУ № 1671-р від 28.12.2020 р., активно розвиваються електронні сервіси ОЗ. Цифровізація дозволяє зменшити бюрократичні перешкоди і навантаження на медичну службу, підвищити якість і доступність послуг, а також (що найважливіше!) надає інструменти для управління, щоб на основі доказових даних формувати політики і приймати ефективні рішення. В цьому контексті важливим є вивчення позитивного закордонного досвіду для його обґрунтованого використання у вітчизняній ОЗ.

Мета: провести дослідження наявних програмних продуктів, що використовуються медичними і фармацевтичними працівниками у Великій Британії для прийняття обґрунтованих доказами рішень щодо забезпечення раціонального застосування ліків.

Матеріали та методи дослідження: системний підхід, аналіз та узагальнення інформації, що міститься на офіційних сайтах органів та організацій ОЗ, професійних асоціацій, розробників програмних продуктів.

Результати та їх обговорення. Для дослідження нами була обрана Велика Британія, яка є лідером у створенні й поширенні доказової інформації, зокрема, національного формуляру, який застосовується у багатьох країнах світу і вважається еталонним, а також інших баз даних (БД) доказової інформації. Pharmaceutical Press, як інформаційний бізнес Королівського фармацевтичного товариства, підтримує працівників ОЗ у всьому світі за допомогою науково обґрунтованих знань для підвищення безпеки пацієнтів. У результаті аналізу встановлено, що Pharmaceutical Press розроблено близько 30 програмних продуктів, 105 друкованих видань, які узагальнюють інформацію стосовно 7500 найменувань ЛЗ.

Таблиця 1

Програмні продукти для підтримки клінічних рішень щодо раціонального застосування ліків

Найменування програмного продукту	Стисла характеристика	Ключові переваги використання
British National Formulary (BNF) – Британський національний формуляр	Невід'ємна частина інфраструктури ОЗ Великої Британії, застосовується у всьому світі; відображає найкращу поточну практику, професійні вказівки щодо використання ЛЗ; наразі чинним є 88 випуск (вересень 2024-березень 2025 р.)	включає понад 1500 ЛЗ; 800 тис. користувачів; прийняття рішень на базі актуальної й достовірної інформації про застосування й дози ЛЗ, рекомендації щодо безпеки (протипоказання, взаємодії); зменшення ризику помилки; оновлення щомісяця; швидка та проста навігація
BNF for Children – Британський національний формуляр для дітей	BNFC містить широкий спектр інформації та рекомендацій стосовно ЛЗ для дітей, доступних у NHS Великої Британії; інформацію про неліцензійне використання певних ЛЗ. Опубліковано у Великій Британії, але інформація є актуальною в будь-якій країні.	містить понад 1000 ЛЗ 220 тисяч користувачів зниження ризику помилок BNFC публікується щорічно, але електронні оновлення випускаються щомісяця швидка та проста навігація
Agilio: Diagnosis and Treatment Guidance – Керівництво з діагностики та лікування	понад 375 клінічних тем, 2000 сценаріїв лікування й діагностики; клінічні теми акредитовані NICE; щомісяця керівництво використовують понад 650 000 клініцистів.	безпечна діагностика й призначення на базі ЕВР; сценарії для практичного лікування, для створення локальних протоколів і настанов; доступний у багатьох країнах.
AHFS Drug Information – Інформація про ЛЗ Американської служби лікарняних формулярів (American Hospital Formulary Service)	Інформація опублікована Американською спілкою фармацевтів (American Society of Health-System Pharmacists) є надійною, повною та об'єктивною БД для ЛЗ, схвалених FDA	єдиний довідник для безпечного й ефективного використання ЛЗ, найширший перелік ЛЗ і показань (97 000); монографії містять актуальні доказові дані щодо терапевтичних рекомендацій і використання «off label»

Продовж. табл. 1

1	2	3
Electronic Medicines Compendium (emc) – Електронний медичний компендіум	актуальна інформація про ЛЗ, ліцензовані у Великій Британії допомагає належним чином призначати й застосовувати ЛЗ; містить інформацію від виробника (SPC) та інформаційні листівки для пацієнтів (PIL).	містить прямі посилання на BNF і BNF для дітей, швидкий доступ до цих інформаційних продуктів дозволяє приймати обґрунтовані рішення щодо ліцензованого та неліцензійного використання ЛЗ
Drug Monitoring Checker – Інструмент для моніторингу лікарських засобів	важливий інструмент для підтримки прийняття рішень щодо моніторингу ЛЗ на місці надання медичної допомоги; надає чіткі, лаконічні та дієві поради до, під час і після лікування.	містить 235 монографій ЛЗ використання в ЗОЗ первинної та вторинної медичної допомоги; оновлення щомісяця
Martindale: The Complete Drug Reference – Мартиндейл: повний довідник ЛЗ (41 випуск, 2024 р.)	джерело достовірних даних з швидким доступом до вичерпної, заснованої на доказах інформації про понад 7900 ЛЗ, що використовуються в усьому світі, показання та інгредієнти понад 300 000 запатентованих ЛЗ з 49 країн, а також майже 700 методів лікування захворювань	монографії про ЛЗ і допоміжні речовини (6400 монографій, 49 розділів). Узагальнено номенклатуру, дії, використання. Містить розділ про нові ЛЗ, які складно класифікувати, трави та ЛЗ, які вже не використовуються. Понад 125 тис товарів, 43 країн довідник виробників – близько 25 000 записів; фармацевтичні терміни (5600 термінів 13 мовами); загальний покажчик : понад 175 000 записів (затверджені назви, синоніми та хімічні назви).

Окрім загальних довідкових застосунків, існує багато спеціалізованих, що зосереджують увагу на певних групах ЛЗ, нозологіях чи групах пацієнтів. Так, створено керівництво для підтримки ефективного лікування тяжкохворих пацієнтів в умовах відділень інтенсивної терапії «Critical Illness», для невиліковних випадків – «Формуляр паліативної допомоги» (Palliative Care Formulary). Також з урахуванням певних особливостей створені електронні керівництва для лікування пацієнтів дитячого віку (Pediatric Injectable Drugs), підтримки безпечного використання ЛЗ під час вагітності та лактації (Drugs in Pregnancy and Lactation). Окремої уваги заслуговують застосунок для підтримки фармакотерапії та вико-

ристання протимікробних засобів (Kucer’s the Use of Antibiotics), управління побічними реакціями (Martindale's ADR Checker). Декілька програмних продуктів спрямовані на підвищення ефективності використання ін’єкційних форм ЛЗ (ASHP Injectable Drug Information, Injectable Drugs Guide), фітопрепаратів (Herbal Medicines), психотропних препаратів (Psychotropic Drug Directory).

Також активно використовуються програмні продукти для швидкого доступу до інформації про взаємодію між ЛЗ, фітопрепаратами, харчовими продуктами, напоями, наркотичними засобами, що базується на опублікованих джерелах, включаючи клінічні дослідження, звіти про випадки та систематичні огляди. Це такі програми як Stockley’s Drug Interactions, Stockley’s Herbal Medicines Interactions, Stockley’s Interactions Checker, Drug Compatibility Checker.

Висновки: використання програмних продуктів, зокрема, пошукових систем, електронних довідкових посібників, експертних систем дозволяє підвищити ефективність ухвалення рішень лікарями та фармацевтами. Разом з тим, застосування багатьох закордонних систем ускладнюється наявністю відмінностей у підходах до організації фармацевтичного забезпечення, зокрема, особливостями медичного страхування та реімбурсації у різних країнах.

Перелік використаних джерел інформації:

1. Agilio Software extends knowledge partnership with NICE. URL: https://www.digitalhealth.net/2023/05/agilio-software-extends-knowledge-partnership-with-nice/ (Date of access: 03.10.2024).
2. AHFS Drug Information. URL: https://www.ashp.org/products-and-services/ahfs-di
3. BNF Publications Online. URL: https://www.pharmaceuticalpress.com/bnf-publications/ (Date of access: 02.10.2024).
4. BNF + BNFC app. URL: https://www.pharmaceuticalpress.com/bnf-publications/bnf-bnfc-app/ (Date of access: 03.10.2024).
5. Electronic Medicines Compendium (emc). URL: https://www.medicines.org.uk/emc#gref (Date of access: 02.10.2024).
6. Martindale: The Complete Drug Reference/ R. Buckingham, ed. 40th ed. London: Pharmaceutical Press, 2020. ISBN 978-0-85711-367-2 (Date of access: 03.10.2024).

РЕПОЗИЦІОНУВАННЯ ЛІКАРСЬКИХ ЗАСОБІВ: ЕВОЛЮЦІЯ ПІДХОДІВ

Подольський І. М., Назаркіна В. М., Бондарєв Є. В.
Національний фармацевтичний університет, м. Харків, Україна
Кафедра клінічної фармакології ІПКСФ
Кафедра соціальної фармації
illya.podolsky@nuph.edu.ua

Вступ. Репозиціонування лікарських препаратів (ЛЗ) – це процес пошуку нових терапевтичних застосувань для існуючих ЛЗ. Цей підхід має низку переваг над традиційними методами пошуку нових ЛЗ, зокрема зниження витрат, скорочення термінів розробки та зменшення кількості невдалих спроб. Використовуючи наявні клінічні дані та дані з безпеки, репозиціонування ЛЗ може прискорити пошук нових методів лікування широкого спектру захворювань. Останні декілька років штучний інтелект (ШІ) став потужним інструментом для репозиціонування ЛЗ, пропонуючи можливість аналізувати надзвичайну велику кількість біомедичних даних, систематизувати їх та виявляти потенційні зв'язки та асоціації.

Мета. Аналіз сучасних підходів до репозиціонування ЛЗ.

Матеріали та методи. Аналіз даних наукової літератури та патентних джерел.

Результати. Аналіз наукової літератури залишає актуальним питання, чому не знижується інтерес до репозиціонування ЛЗ, якщо існує величезний простір для розробки нових ЛЗ, які будуть беззаперечно патентоспроможними? Станом на жовтень 2024 року база даних PubChem містить близько 118,6 мільйонів сполук, більшість з яких можна вважати лікоподібними. База даних відкриття лігандів ZINC20 містить понад 230 мільйонів комерційно доступних малих молекул, які підготовані до віртуального скринінгу за допомогою докінгового програмного забезпечення. Близько 3000 людських білків вважаються потенційними мішенями для лікарських молекул, до того ж постійно удосконалюються підходи, щоб «таргетувати» ті, які раніше вважалися «поза грою» (тобто не мають визначених кишень для зв'язування лігандів).

Питання має просту відповідь – економія коштів, що включає не лише прямі витрати, заощаджені на ранніх стадіях відкриття, дослідженнях на тваринах і ранніх клінічних випробуваннях (які є мінімальними порівняно з програмами фази II і особливо фази III), але й фактори, які часто не беруться до уваги академічними дослідниками: ризики (репозиціонований ЛЗ має набагато менше шансів на невдачу з міркувань безпеки, аніж нова хімічна сполука) та термін виходу на ринок.

Згідно з літературними даними, традиційно існує два основних підходи: "on-target" та "off-target" репозиціонування. "On-target" репозиціонування використовує встановлений механізм ЛЗ для лікування абсолютно іншого захворювання. Воно охоплює застосування в системній медицині, де зміна відомої мі-

шені різними шляхами може призвести до унікальних ефектів. Поштовхом до “on-target” репозиціонування зазвичай є творче поєднання в єдине ціле існуючих доказів.

“Off-target” репозиціонування передбачає використання раніше невідомого, пропущеного або пов’язаного з побічними ефектами фармакологічного механізму ЛЗ, який спочатку був схвалений для іншої мети. Такий підхід вимагає вищого рівня інноваційності та оригінальності, що полегшує отримання патентів на використання.

У минулому переважно оригінальне фармакологічне мислення сприяло репозиціонуванню ЛЗ. Яскравим прикладом є репозиціонування інгібіторів холінестерази, таких як галантамін, на основі холінергічної гіпотези хвороби Альцгеймера в 1990-х роках. Так само репозиціонування селективних інгібіторів зворотного захоплення серотоніну флуоксетин і пароксетин для лікування передчасної еякуляції було засноване на інгібувальному впливі підвищеного рівня серотоніну в мозку на еякуляцію – побічному ефекті, який спостерігається у чоловіків з нормальною сексуальною функцією. Наразі на “on-target” репозиціонування припадає 80-90 % відкриттів. Свіжим прикладом може слугувати патент 2023 року, в якому заявлено ЛЗ від бічного аміотрофічного склерозу рилузол, блокатор рецепторів N-метил-D-аспартату (NMDA) та каїнатних рецепторів, для лікування олігоспермії. Відкриття ґрунтується на доведеній тестикулярній токсичності глутамату та його впливі на розвиток чоловічого безпліддя.
“Off-target” репозиціонування дотепер не призвело до значних впроваджень у практичну медицину, проте науковий інтерес до цих результатів є високим. Статини, розроблені як інгібітори гідроксиметилглутарил-КоА- редуктази для лікування дисліпідемії, є головними кандидатами для такого типу інноваційного репозиціонування. Наприклад, симвастатин заявлено у патентних заявках з 2010 року для лікування цілого ряду захворювань, включаючи панкреатит, сепсис, дегенерацію міжхребцевих дисків, різні види раку, астму, кокаїнову залежність, глаукому та остеоартрит. Такі відкриття часто є результатом високопродуктивного скринінгу, докінгових досліджень *in silico*, а іноді й випадковості.
Хоча традиційні стратегії залишаються продуктивними, ШІ швидко змінює ландшафт репозиціонування ЛЗ. Більш того, можна констатувати, що ШІ може краще підходити для репозиціонування відомих агентів, ніж для розробки нових хімічних речовин, оскільки на сьогодні не бракує даних, які або є у вільному доступі, або можуть бути видобуті після анонімізації, що необхідні для будь-якого складного алгоритму. Рецензована література і патентні документи є цінними джерелами наукової інформації, хоча донедавна вони майже не корелювали між собою внаслідок різниці в мотивації створення.

Висновки. Незважаючи на той факт, що оригінальне фармакологічне мислення та випадковість вже давно є рушійною силою репозиціонування, алгоритмічні підходи саме зараз роблять свій внесок, а використання ШІ останнім часом виводить репозиціонування ЛЗ на безпрецедентний рівень.

Розділ 6

ОРГАНІЗАЦІЙНО-УПРАВЛІНСЬКІ ПІДХОДИ ЗАБЕЗПЕЧЕННЯ РАЦІОНАЛЬНОГО ЗАСТОСУВАННЯ ЛІКАРСЬКИХ ЗАСОБІВ

УДК 615.12:331.108:005.96

ВДОСКОНАЛЕННЯ ПОСАДОВОЇ ІНСТРУКЦІЇ ФАРМАЦЕВТА КЛІНІЧНОГО НА ОСНОВІ РЕЗУЛЬТАТІВ ЕКСПЕРТНОГО ОЦІНЮВАННЯ ПРОФЕСІЙНОЇ ДІЯЛЬНОСТІ

Зарічкова М. В., Мішина І. Ю.

Національний фармацевтичний університет,
м. Харків, Україна
Інститут підвищення кваліфікації спеціалістів фармації
Кафедра організації, економіки та управління фармацією
zarichkova@ukr.net

Анотація. У статті представлено результати розробки посадової інструкції фармацевта клінічного на основі аналізу експертних оцінок його професійної діяльності. Метою дослідження було вивчити ставлення експертів, які працюють у вітчизняних закладах охорони здоров'я, до різних аспектів функціональних обов'язків фармацевта клінічного та використати ці дані для вдосконалення посадової інструкції.

Матеріали та методи. Для досягнення мети дослідження було проведено опитування 50 працівників охорони здоров'я, серед яких фармацевти клінічні, фармацевти та керівники закладів. У дослідженні використовувалися методи анкетування та статистичного аналізу.

Результати. У ході дослідження було визначено основні документи, що регулюють діяльність фармацевта клінічного, уточнено його функції та необхідне обладнання для робочого місця. Особливу увагу було приділено аналізу завдань, обов'язків, прав та сфери відповідальності фармацевта клінічного, а також важливості належного рівня знань для ефективного виконання професійних обов'язків, також були впорядковані ключові показники ефективності (KPI). Дослідження також включало оцінку нормативних актів, таких як накази МОЗ та стандартні операційні процедури, що визначають роботу фармацевта клінічного.

Висновки. В результаті дослідження була розроблена посадова інструкція, яка враховує сучасні вимоги професійної діяльності та стандарти клінічної практики. Очікується, що ця інструкція сприятиме підвищенню ефективності діяльності фармацевтів клінічних і поліпшенню якості фармацевтичної допомоги пацієнтам, а також забезпечить більш раціональне та безпечне використання лікарських засобів у клінічній практиці.

Ключові слова: *заходи БПР, компетенції, посадова інструкція, фармацевт клінічний, функціональні обов'язки, KPI.*

UDC 615.12:331.108:005.96

UPDATING THE CLINICAL PHARMACIST'S JOB DESCRIPTION BASED ON THE RESULTS OF EXPERT EVALUATIONS OF PROFESSIONAL ACTIVITIES

Zarichkova M. V., Mishina I. Yu.

Abstract. The article presents the results of the development of the clinical pharmacist's job description based on the analysis of expert evaluations of his professional activity. The aim of the study was to study the attitude of experts working in domestic healthcare institutions to various aspects of the functional duties of a clinical pharmacist and to use this data to improve the job description.

Materials and methods. To achieve the aim of the study, a survey of 50 health care workers, including clinical pharmacists, pharmacists and facility managers, was conducted. The research used methods of questionnaire and statistical analysis.

Results. In the course of the work, the main documents regulating the activity of the clinical pharmacist were defined, his functions and necessary equipment for the workplace were clarified. Special attention was paid to the analysis of the tasks, duties, rights and sphere of responsibility of the clinical pharmacist, as well as the importance of the appropriate level of knowledge for the effective performance of professional duties Key performance indicators (KPI) were structured. The study also included an assessment of regulatory acts, such as orders from the Ministry of Health and standard operating procedures that define the work of a clinical pharmacist.

Conclusions. Because of the research, a job description was developed that takes into account modern requirements of professional activity and standards of clinical practice. It is expected that job description will contribute to enhancing the effectiveness of clinical pharmacist's activities and improving the quality of pharmaceutical care for patients, as well as ensuring more rational and safe use of medications in clinical practice.

Keywords: *job description, clinical pharmacist, competencies, CPD activities, functional duties, KPI.*

Вступ: Клінічна фармація – це інтегративна прикладна наука, що поєднує фармацевтичні та клінічні аспекти використання лікарських засобів (ЛЗ) [1]. Основна мета клінічної фармації полягає у забезпеченні раціонального та безпечного застосування лікарських препаратів, що вимагає від фармацевтів клінічних (ФК) високого рівня компетенцій, тобто глибоких знань, професійних умінь та практичних навичок [2]. Ця професія з'явилась внаслідок необхідності підвищення якості фармацевтичної допомоги і вперше була впроваджена в країнах Європи та США. У країнах Західної Європи та Північної Америки ФК є ключовим членом мультидисциплінарної медичної команди, який відповідає за оцінку ефективності та безпеки ЛЗ, участь у плануванні лікування пацієнтів, а також навчання медичного персоналу з питань фармакотерапії [3]. В Україні професія ФК почала активно розвиватися з 2009 року, коли було ухвалено На-

каз МОЗ України № 654 від 01.09.2009 р. «Про затвердження Плану заходів щодо покращання здійснення післяреєстраційного нагляду за безпекою та ефективністю лікарських засобів та моніторингу безпеки та ефективності лікарських засобів у стаціонарах закладів охорони здоров'я», що поклав на ФК функцію моніторингу ефективності та безпеки ЛЗ у стаціонарах закладів охорони здоров'я (ЗОЗ) [4]. Згідно з Наказом МОЗ України № 1614 від 03.08.2021 р. «Про організацію профілактики інфекцій та інфекційного контролю в закладах охорони здоров'я та установах / закладах надання соціальних послуг / соціального захисту населення», ФК стали координаторами адміністрування антимікробних препаратів у ЗОЗ [5]. Їх функціональні обов'язки значно розширилися, включивши участь у заходах із профілактики інфекцій та контролю за використанням антимікробних препаратів, що є надзвичайно важливим в умовах сьогодення [6]. Це створило потребу у вдосконаленні посадової інструкції ФК.

Одним із головних завдань дослідження є визначення функціональних завдань, обов'язків та відповідних компетенцій ФК, які проявляються в знаннях і навичках, на основі експертного оцінювання професійної діяльності в умовах сьогодення. Залучення експертів дозволило врахувати актуальні вимоги стандартів клінічної практики та досвід провідних фахівців, що забезпечує високий рівень відповідності посадової інструкції сучасним викликам. Важливість цього дослідження полягає у вдосконаленні професійної діяльності ФК, створенні відповідних заходів безперервного професійного розвитку (БПР), а також у визначенні їхньої ролі в мултидисциплінарній медичній команді, що напряму впливає на якість надання фармацевтичної допомоги.

У цьому дослідженні було проаналізовано зміни, які необхідно внести до посадової інструкції ФК, щоб адаптувати її до нових вимог сучасної фармацевтичної практики та забезпечити ефективну взаємодію з медичним персоналом.

Мета: Дослідити ставлення експертів, які працюють у вітчизняних ЗОЗ, до професійної діяльності та функціональних обов'язків ФК для вдосконалення посадової інструкції.

Матеріали та методи дослідження: У дослідженні були використані наступні методи: аналіз літературних джерел, експертне опитування, статистичний, дисперсійний та якісний аналізи.

Результати та їх обговорення:

Матеріалом дослідження слугували дані опитування, в якому взяли участь 50 осіб, які працюють у вітчизняних ЗОЗ. Загалом серед опитаних 26 осіб працювали на посадах ФК, 15 – фармацевта, і 9 обіймали керівні посади. Більшість респондентів працювали в ЗОЗ, які були розподілені рівномірно за регіонами України. Для оцінки важливості аспектів діяльності ФК учасники використовували шкалу від 1 до 5 балів, оцінюючи нормативні документи, права, обов'язки, функції,компетенції (знання, уміння, навички), та KPI. Об'єкти ранжували за середніми рангами, застосовуючи дисперсійний аналіз Фрідмана (Friedman $\chi 2$) при довірчій імовірності 95%, а при значущих розбіжностях ($p<0.05$) проводили попарні порівняння за критерієм Уїлкоксона (в тексті ста-

тистику критерію позначено як Wilcoxon T, її нормальну апроксимацію Z) з поправкою Холма-Бонферроні на множинність.

Анкета складалась з 26 питань, на які надавали відповіді фахівці, які безпосередньо працюють в ЗОЗ. Перші 8 питань мали загальний характер, наступні питання стосувались безпосередньо професійної діяльності ФК.

На початку дослідження булі поставлені питання, які мали розкрити важливість та необхідність внутрішніх нормативних документів перелічених в анкеті, які потрібні в роботі ВІК. Незважаючи на високі оцінки по всім запропонованим документам, найбільш важливими вбачаються розроблені стандартні операційні процедури (СОП) (середній бал – 4,84), накази про їх створення (середній бал – 4,84) і Положення про ВІК (середній бал – 4,84); найменш необхідними – протоколи самооцінювання (середній бал – 4,36).

Ще одним питанням було визначення найбільш важливих наказів МОЗ України, якими повинен керуватися ФК при виконанні своїх службових обов'язків. Із запропонованих учасникам дослідження 15 наказів МОЗ, за результатами оцінювання важливості та необхідності цих наказів, як найбільш значимі можна вказати Наказ МОЗ №1513 "Про затвердження Стандарту медичної допомоги «Раціональне застосування антибактеріальних і антифунгальних препаратів з лікувальною та профілактичною метою»" (середній бал – 4,80) [7], а також накази №822 "Про затвердження Стандарту «Парентеральна періопераційна антибіотикопрофілактика»" (середній бал – 4,78) [8] і №1614 "Про організацію профілактики інфекцій та інфекційного контролю в закладах охорони здоров'я та установах / закладах надання соціальних послуг / соціального захисту населення" (середній бал – 4,74).

Найменш важливими в роботі ФК на думку опитаних є: Наказ МОЗ №280 "Щодо організації проведення обов'язкових профілактичних медичних оглядів працівників окремих професій, виробництв і організацій, діяльність яких пов'язана з обслуговуванням населення і може призвести до поширення інфекційних хвороб" (середній бал – 3,82) [9], Наказ МОЗ №552 "Про затвердження Державних санітарних норм та правил «Дезінфекція, передстерилізаційне очищення та стерилізація медичних виробів в закладах охорони здоров'я»" (середній бал – 3,86) [10], Наказ МОЗ №325 "Про затвердження Державних санітарно-протиепідемічних правил і норм щодо поводження з медичними відходами" (середній бал – 3,84) [11]. Додатково до запропонованого списку, респонденти надавали пропозиції включити й інші накази МОЗ, серед яких найбільш часто (10%) згадувався Наказ МОЗ №403 від 05.03.2021 р. [12]. Також 6% пропонували включити до переліку необхідних наказів Наказ МОЗ №610 від 20.03.2019 р. [13]. В поодиноких випадках зустрічалися пропозиції щодо важливості та необхідності для регламентування професійної діяльності ФК інших наказів МОЗ.

Наступним питанням, яке ми розглядали, було визначення важливості СОП для роботи ФК. Метою було проаналізувати, які з СОП учасники дослідження вважають найбільше і найменше актуальними у своїй професійній діяльності. За результатами оцінювання 36 СОП ми визначили статистично зна-

чущі відмінності в їхній важливості (Friedman $\chi 2=456.081$, $p<0.00001$). Узагальнюючи результати попарних порівнянь різних СОП за їх важливістю, в якості найбільш важливих можна виділити п'ять з них: «Адміністрування антимікробних препаратів» (середній бал – 4,86), «Раціональне використання антибіотиків, емпірична антибіотикотерапія» (середній бал – 4,84), «Антибіотикопрофілактика» (середній бал – 4,82), «Раціональне застосування антибактеріальних і антифунгальних препаратів з лікувальною та профілактичною метою» (середній бал – 4,80) і «Профілактика передавання мікроорганізмів із множинною резистентністю до антимікробних препаратів» (середній бал – 4,78). В якості найменш важливих для роботи ФК СОПів можна зазначити СОП «Поводження з інструментарієм» (середній бал – 3,70) і СОП «Поводження з білизною» (середній бал – 3,68).

Ще одним питанням було визначення важливості та необхідності ключових функцій посади ФК. Учасникам дослідження було запропоновано оцінити 8 основних функцій цієї посади, після чого отримані результати були впорядковані за спаданням важливості. Функції, пов'язані з адмініструванням антимікробних препаратів і поширенням інформації про них, вважаються опитаними значущо важливішими та необхіднішими для посади ФК, ніж виконання організаційних, профілактичних і протиепідемічних заходів (Рис. 1).

Серед питань, які ми визначали, було визначення важливості та необхідності компетенцій і функціональних завдань, які виконують ФК у своїй професійній діяльності. Для учасників анкетування було запропоновано оцінити перелік із 21 позиції, що включає навички та завдання, важливі для роботи ФК [14].




Рис. 1 Ранжування функцій посади ФК за важливістю та необхідністю на думку опитаних.

На підставі отриманих оцінок було знайдено середні ранги кожної з цих позицій, після чого оцінювані навички та завдання впорядковано за їх спаданням (рис. 2).

Згідно з проведеним статистичним аналізом, як найважливіші та найнеобхідніші, можна вказати перші шість навичок з отриманого ранжованого переліку: постійно удосконалює свій професійний рівень; проводить адміністрування антимікробних препаратів; зберігає професійну таємницю, дотримується вимог професійної етики і медичної деонтології; керується законодавством України про охорону здоров'я; сприяє впровадженню нових клінічних протоколів з емпіричної антимікробної терапії та періопераційної антибіотикопрофілактики; готує звітну інформацію з питань ПІІК, впровадження адміністрування антимікробних препаратів, як найменш важливі – останні три: проводить інвентаризацію товарно-матеріальних цінностей; надає домедичну допомогу особам при невідкладних станах та постраждалим у екстремальних ситуаціях; координує введення СОП щодо гігієни рук.




Рис. 2 Ранжування навичок і функціональних завдань фармацевта клінічного за їх важливістю та необхідністю на думку опитаних учасників дослідження.

В рамках нашого дослідження було поставлено питання важливості та необхідності прав ФК. Було запропоновано сім формулювань, які були оцінені респондентами на найвищому рівні з медіанною оцінкою 5 балів (Рис.3). Звідси можемо зробити висновок про доцільність включення до посадової інструкції посади ФК усіх цих формулювань. Одним з прав ФК, важливість і необхідність

якого опитані оцінили на високому рівні, є право бути забезпеченим робочим місцем із необхідним устаткуванням. В анкету ми включили відкрите питання, в якому респондентам пропонувалося вказати, чим повинно бути обладнане робоче місце ФК. Свою думку з цього питання висловили 80% респондентів. Більшість (75,0%) зазначила, що на робочому місці ФК необхідний персональний комп'ютер. Вагома частка респондентів (42.5% і 35,0%) вказала на потребу в оргтехніці (принтер, сканер) та підключенні до інтернету. Менша кількість респондентів згадала про спеціалізоване програмне забезпечення та канцелярське приладдя (12.5%), нормативну документацію (10,0%), робочий стіл і телефон (7.5%), меблі та спеціалізоване обладнання (5,0%), а також побутові зручності (2.5%). Отже, найважливішим є персональний комп'ютер із доступом до інтернету та оргтехнікою.



Рис. 3 Ранжування прав ФК за їх важливістю та необхідністю на думку опитаних учасників дослідження.

Наступним питанням у нашому дослідженні було оцінювання сфери відповідальності ФК. Учасникам опитування було запропоновано шість формулювань, які описують основні сфери відповідальності ФК, які за оцінками респондентів були нами впорядковані у відповідності до їх важливості та необхідності. До результатів проведеного аналізу, як найбільш важливі сфери відповідальності ФК, слід вказати організацію своєї роботи , своєчасне і кваліфіковане виконання наказів, розпоряджень і доручень керівництва (середній бал – 3,94), а також надання в установленому порядку статистичної та іншої інформації (звітності) (середній бал – 3,89); найменш важливою і необхідною – облік, витрати і зберігання ЛЗ, матеріалів, реактивів тощо (середній бал – 2,77).

Ще одним питанням було визначення знань, якими необхідно володіти ФК для ефективного виконання своїх професійних обов’язків. Респондентам було запропоновано оцінити список із 29 позицій, що дозволило впорядкувати

ці знання за важливістю та необхідністю [15]. Відповідно результатів проведеного аналізу анкетування, найменш важливими та необхідними для роботи ФК опитані вважають знання з організації підприємницької діяльності; найбільш необхідними – знання принципів моніторингу ефективності та безпеки застосування ЛЗ, державної мови та принципів складання лікарського формуляру, які вони можуть отримувати на заходах БПР.

Черговим пунктом нашого дослідження було оцінювання важливості та необхідності діяльності ФК, у рамках виконання своїх службових обов'язків поза межами лікарні. Результати опитування експертів показали, що ця діяльність була оцінена на досить високому рівні. При цьому найбільш важливою функцією на думку респондентів виявилася взаємодія з іншими ФК, медичним персоналом, агентами забезпечення медичною технікою та іншими стейкхолдерами.

В рамках нашого дослідження стало визначення ключових показників ефективності (KPI) ФК. В результаті було впорядковано 20 KPI за пріоритетністю на основі експертних оцінок. Результати обробки отриманих відповідей свідчать, що усі 20 запропонованих нами показників доцільно застосовувати в якості ключових показників ефективності роботи ФК, бо усі отримані медіанні оцінки становлять 5 або (і лише для двох KPI) 4 бали (рис. 4).

Оцінюючи форми і способи вдосконалення професійного рівня, найбільш важливими та необхідними для ФК респонденти визнали 6 заходів БПР, до яких відносяться професійний розвиток за дистанційною формою навчання з використанням електронних навчальних ресурсів, підвищення кваліфікації на циклах тематичного удосконалення, присвоєння кваліфікації відповідної спеціальності, навчання на симуляційних тренінгах або тренінгах з оволодіння практичними навичками, тематичне навчання (семінари, майстер-класи тощо), а також науково-практичні конференції, конгреси, симпозіуми.

Значущо менш важливими на думку опитаних є наукова робота: публікація статей і здобуття наукових ступенів.

Висновки: В результаті нашого дослідження було визначено найбільш значущі накази, що стосуються раціонального застосування антибактеріальних препаратів, організації профілактики інфекцій, а також СОПи, накази про їх створення та положення про ВІК, що підкреслило зростаючу роль ФК в умовах сучасних викликів у системі охорони здоров'я, зокрема в контексті зростаючої антибіотикорезистентності та необхідності оптимізації лікувальних стратегій.

Дослідження також підтвердило, що функціональні обов'язки ФК суттєво розширилися, включаючи не лише моніторинг безпеки та ефективності ЛЗ, а й активну координацію використання антимікробних препаратів. Результати дослідження акцентують увагу на важливості наявності належного рівня компетенцій, які проявляються в знаннях і навичках, що в свою чергу зумовлюють створення відповідних заходів БПР. Це свідчить про необхідність організації підвищення кваліфікації ФК, що дозволить їм адаптуватися до змін у нормативно-правовій базі та клінічній практиці. Важливим аспектом дослідження стало

визначення ключових показників ефективності (KPI), які можуть слугувати основою для оцінювання професійної діяльності ФК.




Рис. 4 Ранжування обов'язків KPI діяльності ФК за їх важливістю та необхідністю на думку опитаних учасників дослідження.

На основі отриманих результатів дослідження була вдосконалена посадова інструкція для ФК, що відповідає сучасним вимогам їх професійної діяльності. Встановлено, що ця інструкція враховує функціональні обов'язки ФК, а також ключові аспекти його діяльності, такі як належне оформлення документації, ефективне управління антимікробними препаратами, контроль за виконанням СОП та активна співпраця з медичним персоналом. Запропоновані рекомендації можуть стати основою для подальших досліджень у сфері професійного розвитку ФК.

Перелік використаних джерел інформації:

1. Фармація клінічна. Фармацевтична енциклопедія України. URL:https://www.pharmencyclopedia.com.ua/article/339/farmaciya-klinichna.
2. Зупанець, К., & Пропіснова, В. (2023). Клінічна фармація в україні: рік 2023. *Українські медичні вісті*, (3-4 (96-97), 13–18. DOI: https://doi.org/10.32782/umv-2023.2.3.
3. Urbańczyk, K., Guntschnig, S., Antoniadis, V., et al. (2023). Recommendations for wider adoption of clinical pharmacy in Central and Eastern Europe in order to ptimize pharmacotherapy

and improve patient outcomes. *Frontiers in pharmacology*, *14*, 1244151. DOI: https://doi.org/10.3389/fphar.2023.1244151.

4. Наказ МОЗ України № 654 від 01.09.2009 р. «Про затвердження Плану заходів щодо покращання здійснення післяреєстраційного нагляду за безпекою та ефективністю лікарських засобів та моніторингу безпеки та ефективності лікарських засобів у стаціонарах закладів охорони здоров'я». URL: https://zakon.rada.gov.ua/rada/show/v0654282-09#Text.
5. Наказ МОЗ України № 1614 від 03.08.2021 р. «Про організацію профілактики інфекцій та інфекційного контролю в закладах охорони здоров'я та установах / закладах надання соціальних послуг / соціального захисту населення». URL: https://zakon.rada.gov.ua/laws/show/z1318-21#Text.
6. ВООЗ. Глобальний план дій щодо протимікробної стійкості. 2015. URL: https://iris.who.int/handle/10665/193736.
7. Наказ МОЗ України № 1513 від 23.08.2023 «Про затвердження Стандарту медичної допомоги «Раціональне застосування антибактеріальних і антифунгальних препаратів з лікувальною та профілактичною метою»». URL: https://moz.gov.ua/uk/decrees/nakaz-moz-ukraini-vid-23082023—1513-pro-zatverdzhennja-standartu-medichnoi-dopomogi-racionalne-zastosuvannja-antibakterialnih-i-antifungalnih-preparativ-z-likuvalnoju-ta-profilaktichnoju-metoju.
8. Наказ МОЗ України № 822 від 17.05.2022 «"Про затвердження Стандарту «Парентеральна періопераційна антибіотикопрофілактика»". URL: https://moz.gov.ua/uk/decrees/nakaz-moz-ukraini-vid-17052022—822-pro-zatverdzhennja-standartu-parenteralna-perioperacijna-antibiotikoprofilaktika.
9. Наказ МОЗ України № 280 від 23.07.2002 «Щодо організації проведення обов'язкових профілактичних медичних оглядів працівників окремих професій, виробництв і організацій, діяльність яких пов'язана з обслуговуванням населення і може призвести до поширення інфекційних хвороб». URL: https://zakon.rada.gov.ua/laws/show/z0639-02#Text.
10. Наказ МОЗ України №552 від 11.08.2014 «Про затвердження Державних санітарних норм та правил «Дезінфекція, передстерилізаційне очищення та стерилізація медичних виробів в закладах охорони здоров'я»». URL: https://zakon.rada.gov.ua/laws/show/z1067-14#Text.
11. Наказ МОЗ України №325 від 08.06.2015 «Про затвердження Державних санітарно-протиепідемічних правил і норм щодо поводження з медичними відходами». URL: https://zakon.rada.gov.ua/laws/show/z0959-15#Text.
12. Наказ МОЗ України №403 від 27.02.2023 «Про затвердження Порядку проведення посиленого епідеміологічного нагляду за протимікробною резистентністю мікроорганізмів, що спричиняють гнійно-запальні інфекції ран у поранених внаслідок бойових дій». URL: https://zakon.rada.gov.ua/laws/show/z0489-23#Text.
13. Наказ МОЗ України № 610 від 20.03.2019 «Про внесення змін до Розподілу лікарських засобів для лікування дітей, хворих на онкологічні та онкогематологічні захворювання, закуплених за кошти Державного бюджету України на 2017 рік...». URL: https://moz.gov.ua/uk/decrees/nakaz-moz-ukraini-vid-20032019—610-pro-vnesennja-zmin-do-rozpodilu-likarskih-zasobiv-dlja-likuvannja-ditej-hvorih-na-onkologichni-ta-onkogematologichni-zahvorjuvannja-zakuplenih-za-koshti-derzhavnogo-bjudzhetu-ukraini-na-2017-rik.
14. Довідник кваліфікаційних характеристик професій працівників. Випуск 78. Охорона здоров'я. URL: https://zakon.rada.gov.ua/rada/show/va117282-02#Text.
15. Професійний стандарт за професією «Фармацевт клінічний». URL: https://register.nqa.gov.ua/uploads/0/495 do_nakazu_05_23_profstandart_farmacevt_klinicnij.pdf

УДК: 614.2:615.1:658

АНАЛІЗ ЗАРУБІЖНОГО ТА ВІТЧИЗНЯНОГО ДОСВІДУ ДЕРЖАВНОГО РЕГУЛЮВАННЯ СОЦІАЛЬНОЇ ВІДПОВІДАЛЬНОСТІ ФАРМАЦЕВТИЧНОГО БІЗНЕСУ

Братішко Ю. С., Толочко В. М., Зарічкова М. В., Артюх Т. О., Ковальська О. В.

Інститут підвищення кваліфікації спеціалістів фармації Національного фармацевтичного університету, м. Харків, Україна
Кафедра організації, економіки та управління фармацією
bratishko1411@gmail.com

Анотація. Розроблена вертикаль державного регулювання соціальної відповідальності фармацевтичного бізнесу із залучення професійних спілок, асоціацій, союзів та громадських об'єднань, які повинні активно втручатися у процес розвитку соціальної відповідальності фармацевтичного бізнесу. Визначені напрями державного регулювання соціальної відповідальності фармацевтичного бізнесу в Україні та розроблений механізм державного регулювання соціальної відповідальності фармацевтичного бізнесу. Розроблений алгоритм послідовності дій щодо формування механізму державного регулювання соціальної відповідальності фармацевтичного бізнесу в Україні. Запропоновані заходи з удосконалення державного регулювання соціальної відповідальності фармацевтичного бізнесу в Україні, що сприятиме підвищенню інвестиційної привабливості вітчизняних фармацевтичних компаній для закордонних інвесторів, посилить їх конкурентоспроможність на міжнародному фармацевтичному ринку шляхом набуття статусу інноваційних, соціально відповідальних організацій, які ефективно вибудовують партнерські відносини, та сприятиме підвищенню корпоративної репутації.

Ключові слова: *соціальна відповідальність фармацевтичного бізнесу; державне регулювання соціальної відповідальності фармацевтичного бізнесу; фармацевтична компанія; фармація.*

UDC: 614.2:615.1:658

ANALYSIS OF FOREIGN AND NATIONAL EXPERIENCE OF STATE REGULATION OF SOCIAL RESPONSIBILITY OF PHARMACEUTICAL BUSINESS

Bratishko Y.S., Tolochko V.M., Zarichkova M.V., Artyukh T.O., Kovalska O.V.

Absract. A vertical of the state regulation of the social responsibility of the pharmaceutical business has been developed to attract trade unions, associations, unions and public associations, which should actively intervene in the process of developing the social responsibility of the pharmaceutical business. The main directions of the state regulation of the social responsibility of the pharmaceutical business in Ukraine and the mechanism of the state regulation of the social responsibility of the

pharmaceutical business have been determined. A sequence algorithm has been developed for the formation of a mechanism for state regulation of the social responsibility of the pharmaceutical business in Ukraine. Proposed measures to improve the state regulation of social responsibility of the pharmaceutical business in Ukraine, which will contribute to increasing the investment attractiveness of domestic pharmaceutical companies for foreign investors, strengthen their competitiveness in the international pharmaceutical market by becoming innovative, socially responsible organizations that effectively build partnerships, and will contribute to improving not only the corporate reputation, but also reputation.

Keywords: *social responsibility of the pharmaceutical business; state regulation of social responsibility of pharmaceutical business; pharmaceutical company; pharmacy.*

Вступ. Сьогодні у наукових колах вже досягнуто певного узгодження у питаннях принципових характеристик соціальної відповідальності (СВ), вона визнана концептуальною й ідеологічною основою соціальної політики сучасного фармацевтичного бізнесу, але все ж СВ лишається складно регульованим і багаторівневим напрямом діяльності фармацевтичних організацій. Насамперед, це пов’язано з тим, що СВ реалізується у конкретному суспільстві, у межах національних законів і регулятивних положень. Тому не можна механізми державного регулювання сліпо копіювати та переносити з одного контексту в інший. Для того, щоб уникнути подібних помилок потрібний аналіз зарубіжного досвіду державного регулювання соціальної відповідальності бізнесу (СВБ).

Мета: розробка науково-практичних засад управління і державного регулювання СВД фармацевтичних компаній (ФК) в Україні для зростання соціальної спрямованості їх діяльності, інвестиційної привабливості і конкурентоспроможності, що сприятиме підвищенню якості, ефективності і доступності фармацевтичного забезпечення населення України.

Матеріали та методи дослідження: загальнонаукові та спеціальні методи пізнання, зокрема, метод структурно-системного аналізу, наукового синтезу (для узагальнення існуючого міжнародного досвіду стосовно регулювання СВ), логічного узагальнення (для побудови механізму державного регулювання соціальної відповідальності фармацевтичного бізнесу (СВФБ), спостереження (використовувався для вивчення сучасних світових тенденцій в управлінні СВ), експертного опитування (вивчення напрямків державного регулювання СВФБ), метод моніторингу та ін.

Результати та їх обговорення. У закордонних країнах соціальна відповідальність бізнесу (СВБ) представлена декількома сталими системами, про які прийнято говорити як про моделі: американську, європейську, азійську (японську) та пострадянську. Нами узагальнені особливості державного регулювання по всіх зазначених моделях за напрямками СВБ (табл. 1).

Характеризуючи в цілому міжнародну практику державного регулювання СВБ, можна відзначити, що досвід реалізації концепції СВ досить багатий та

різноманітний і підтверджує існування різних підходів у цій сфері. Незважаючи на деякі відмінності, загальною характерною рисою як американської, європейської, азійської (японської) та пострадянської моделей, є те, що розвиток і впровадження СВБ підтримується державою, некомерційними організаціями та суспільством у цілому, але ступінь втручання держави різна.

Таблиця 1

Аналіз зарубіжного досвіду державного регулювання за складовими соціальної відповідальності фармацевтичного бізнесу

Існуючі моделі СВБ у світі	Складові СВФБ						Наявність цілісного механізму державного регулювання СВБ	Наявність керуючого органу державного влади, відповідального за реалізацію політики СВБ
	відповідальність ФК за своєчасне забезпечення населення якісними, безпечними та доступними за ціною ЛЗ	відповідальність за дотримання чинного законодавства та міжнародних стандартів	відповідальність ФК за екологічну безпеку	відповідальність ФК перед персоналом та суспільством	відповідальність ФК перед акціонерами та власниками	відповідальності ФК перед бізнес-середовищем		
Американська	+	+	+	–	–	–	–	–
Європейська	+	+	+	+	+	+	+	+
Італія	+	+	+	+	+	+	+	Міністерство праці та соціальної політики
Іспанія	+	+	+	+	+	+	+	Державна Експертна технічна комісія з СВБ
Польща	+	+	+	+	+	+	+	Робоча група з питань СВБ при Міністерстві праці
Данія	+	+	+	+	+	+	+	Міністерство соціальних справ
Нідерланди	+	+	+	+	+	+	+	Міністерство будівництва, територіального планування та навколишнього середовища, Міністерство соціальної політики та зайнятості, відділ міжнародної кооперації та технічної допомоги Міністерства закордонних справ, Міністерство сільського господарства, природи та якості харчування, Центр Компетенцій СВБ Міністерства економіки
Великобританія	+	+	+	+	+	+	+	Міністр із соціальної відповідальності компаній, Державна асоціація «Бізнес у співтоваристві»
Азійська (японська)	+	+	+	+	+	+	–	–
Пострадянська	+	+	–	–	–	–	–	–

Сприйняття та підтримка органів державної влади у значній мірі встановлює ступінь орієнтованості бізнесових кіл країни на СВ та її дієвість. У країнах, де держава активно підтримує та координує КСВ, СВБ набирає обороти як життєздатний соціальний інститут. Особливо добре цей інститут розбудовується там, де сформульована державна стратегія сприяння СВ, установлені законодавчі рамки і створено мережі інституційної підтримки. Співробітництво замість надмірного регулювання перетворює державу з патрона на партнера, надаючи підґрунтя створення дієвої системи державно-приватного соціального партнерства. Таким чином, розглянутий досвід зарубіжних країн у сфері регулювання СВБ дає змогу виокремити низку інструментів, які доцільно впровадити у вітчизняну практику взаємовідносин держави і фармацевтичного бізнес-середовища. Система державного регулювання СВБ в Україні на сьогодні ще формується (рис. 1).



Рис. 1 Періодизація розвитку державного регулювання СВБ в Україні.

Як показано на рис. 1, ініціація цього процесу відбулася зовсім нещодавно, у 2005 р. У 1989 р. громадськими активістами-фахівцями з якості та систем-

ного менеджменту була заснована Всеукраїнська громадська організація «Українська асоціація якості» (УАЯ), яка за час свого існування набула широкого міжнародного визнання. 1.01.2016 р. згідно новому Закону України «Про громадські об`єднання», частина ВГО УАЯ, яка складалась з юридичних осіб, трансформована до формату Громадської спілки «Українська асоціація досконалості та якості» (УАДЯ), виступаючи правонаступником ВГО УАЯ [1].

Український союз промисловців і підприємців (УСПП) є потужним громадським об'єднанням, куди увійшли представники бізнесу, функціонує з 15 лютого 1992 року [2]. У 2007 році УСПП запровадив і щорічно захищає систему управління якістю ISO 9001:2000. Союз співпрацює більше, ніж з 100 громадськими організаціями та галузевими асоціаціями, які спільно розробили стратегію інтеграції промисловості з наукою і фінансами, яка здатна у відносно короткий період вивести промисловість України на шлях сталого інноваційного розвитку. Успішно діють започатковані УСПП та дружньою Конфедерацією промисловців Литви Українсько-литовська та Литовсько-українська ділові ради. УСПП та його члени сповідують принцип СВБ. Союз одним із перших підтримав ініціативу створення в Україні Мережі доброчесного бізнесу, а з 2008 року є підписантом Глобального договору в цій царині [3].

Водночас результати проведених нами досліджень (рис. 2) свідчать про те, що наявність в Україні організацій-лідерів у сфері СВ поєднується з масовими фактами порушень з боку роботодавців прав людини і трудових стандартів, визнаних міжнародною спільнотою (53,4% експертів); недотриманням законодавчо встановлених соціально-трудових прав і гарантій працівників (47,9% експертів); низьким рівнем заробітної плати через небажання сплачувати податки та внески на соціальне страхування (59,5 % експертів); заборгованістю з виплати заробітної плати (33,2% експертів); невиправданою економією на інвестиціях в охорону праці, здоров'я працівників та їх професійний розвиток (52,8%); нехтуванням проблемами збереження довкілля (35,3% експертів).



Рис. 2 Гістограма розподілу експертних відповідей стосовно наявних порушень СВ поведінки ФК в Україні.

Першим і поки що єдиним законодавчим актом, що стосується СВБ в Україні та порядку ведення соціального діалогу з метою вироблення та реалізації державної соціальної і економічної політики, регулювання трудових, соціальних, економічних відносин та забезпечення підвищення рівня і якості життя громадян, соціальної стабільності в суспільстві, є Закон України «Про соціальний діалог в Україні» № 2862-VI від 23 грудня 2010 р. [5].

Але сьогодні вже можна говорити про досить успішну інституцілізацію принципів СВБ вітчизняним бізнес-співтовариством: прийнято низку документів, які регламентують реалізацію принципів СВ на практиці. Так, у 2011 р. мережею Глобального договору ООН в Україні, Центром розвитку корпоративної соціальної відповідальності та іншими організаціями спільно з УСПП розроблено проєкт Концепції національної стратегії соціальної відповідальності бізнесу в Україні, який після широкого громадського обговорення було схвалено експертним співтовариством і профільними міністерствами, а у 2020 році відбулося її затвердження (Розпорядження КМУ «Про схвалення Концепції реалізації державної політики у сфері сприяння розвитку соціально відповідального бізнесу в Україні на період до 2030 року» від 24 січня 2020 р. № 66-р).

В той же час, в Україні досі відсутні фахові самоврядні бізнес-асоціації з СВБ, що характерно для більшості зарубіжних країн. Можна відзначити, що на відміну від західних компаній, де потужним імпульсом для розвитку СВБ стали дії громадянського суспільства (бойкоти споживачів, протести екологів та ін.), український бізнес став на шлях СВ за умови значного впливу держави.

Незважаючи на те, що чільна роль у розвитку практики СВ українського бізнесу належить владі, тим не менш, спостерігаємо відсутність гарантій податкового стимулювання добродійності і СВ з боку держави. Саме цей факт, є відмітною рисою вітчизняної моделі СВБ і підтверджує її неформальний характер. Проведені нами дослідження дозволяють стверджувати, що важливими відмінностями української моделі СВФБ від її закордонних аналогів є значимість різних груп стейкхолдерів. Для українських ФК характерно досить істотне звуження кола значимих стейкхолдерів, зокрема, за рахунок недооцінки й, відповідно, недостатнього залучення до процесу реалізації бізнесом СВ таких груп стейкхолдерів, як, наприклад, профспілки, профільні громадські організації та місцеве співтовариство. Як правило, основними рушійними силами виступають безпосередньо самі провідні вітчизняні ФК і місцеві органи публічної влади. Разом з тим, необхідно звернути увагу, що неурядові організації, які поряд з крупними ФК є основними ініціаторами програм СВБ в Європі, в Україні такої функції не виконують. На підставі вивчення світового досвіду нами запропонована схема побудови інституціональної вертикалі управління СВФБ (рис. 3).

Основними факторами, що визначають специфічність формування соціально відповідальних практик в Україні і зокрема у фармації, на думку експертів, є: сталість соціальних очікувань населення стосовно держави (54 % експертів); несформованість суспільного запиту на СВБ (33 % експертів); висока залежність бізнесу від органів держвлади, місцевого самоврядування (62 % експертів).




Рис. 3 Схема інституціональної вертикалі управління СВФБ

Але слід зазначити, що формуванню СВБ в Україні сприяють наступні обставини: 1. Наявність великого акціонерного капіталу, оформленого в корпоративні структури, що мають необхідні фінансові ресурси для соціальних інвестицій. 2. Залучення країни до євроінтеграційних процесів, що сприяє інституціалізації концепції СВБ за зразками, прийнятими в міжнародній практиці. 3. Надходження іноземних інвестицій, що передбачає оцінку дотримання українськими бізнесами-партнерами норм СВБ, прийнятих світовою бізнес-спільнотою. 4. Поступовий розвиток фондового ринку і первинне публічне розміщення акцій (IPO), які передбачають реалізацію програми підвищення прозорості і поліпшення корпоративного управління, орієнтованої, в першу чергу, на збільшення курсової вартості акцій і системи комунікацій з інвесторами.

Тим часом, в Україні сьогодні є низка проблем у формуванні сприятливого середовища для СВБ. На підставі контент-аналізу публікацій в ЗМІ, аналізу наукових джерел та висновків експертів, нами виділені наступні негативні фактори, які уповільнюють розвиток СВБ в країні:

На національному рівні: несприятливе для реалізації СВБ податкове і цивільне законодавство; відсутність системи мотивації СВД суб'єктів господарювання; відсутність інституціональної координації СВБ (відповідних управлінських структур немає ані в державному апараті, ані на рівні місцевої влади, ані на галузевому рівні).

На регіональному, місцевому та галузевому рівнях: психологічна неготовність органів влади різного рівня взаємодіяти в системі державно-приватного соціального партнерства; не розвиненість системи публічно-договірних відносин з бізнес-структурами; високий рівень корупції; відсутність інформаційної та методичної підтримки СВБ з боку різних гілок влади (регіональної, місцевої, галузевої).

У керівництві органів державної влади на сьогодні відсутнє розуміння необхідності проведення чіткої політики в сфері СВБ, у тому числі й відносно великих ФК. Проведений нами аналіз функцій органів виконавчої влади показав, що жодний з них не займається питаннями регулювання в сфері СВБ й сталого розвитку. Частково цю прогалину намагаються заповнити Центр «розвиток КСВ», УСПП і Форум добродійників, але це громадські об`єднання бізнес-структур, а не органи державного управління.

Для визначення переліку та вагомості перешкод інституціонального характеру, які стоять на заваді розвитку СВБ в Україні і ,зокрема, у фармації, нами був використаний метод експертного опитування. Анкетування проводилось серед менеджерів та керівників провідних ФК та членів Українського союзу промисловців та підприємців. Шляхом групування та аналізу їх відповідей, нами визначені найбільш суттєві перешкоди розвитку СВБ в Україні (табл. 2).

Найбільш вагомою перешкодою на шляху розвитку СВФБ експертами визначено відсутність в Україні механізму державно-приватного соціального партнерства. Друге місце серед перешкод посіла пріоритетність забезпечення поточної фінансово-економічної стабільності діяльності ФК. З одного боку, це

Таблиця 2

Результати експертного опитування відносно наявності та вагомості перешкод розвитку СВФБ в Україні

Ранг значимості певної перешкоди СВБ	Перешкоди	Відсоток відповідей експертів %
1	Відсутність дієвого механізму державно-приватного соціального партнерства	82
2	Пріоритет забезпечення поточної фінансово-економічної стабільності діяльності ФК	70
3	Значна частка тіньового сектору в економіці України	68
4	Певна відстороненість ФК від соціальних проблем життя українського суспільства та несформованість стандартів участі в ньому	66
5	Недостатнє стимулювання СВД ФК з боку держави	64
6	Відсутність традицій соціально відповідальної поведінки ФК	54
7	Недостатній рівень духовно-моральної культури керівників ФК	52
8	Інформаційна закритість діяльності багатьох ФК	50
9	Нерозвиненість механізмів участі ФК у забезпеченні соціально-економічного розвитку територій присутності	48
10	Недостатній рівень корпоративної культури вітчизняних ФК	38
11	Інше	4

природно, оскільки відповідає цілям підприємницької діяльності. Але домінування економічних цілей діяльності у ФК над соціальними та екологічними йде всупереч із принципами сталого розвитку вітчизняної фармації. Третє місце серед основних перешкод посіла висока частка тіньового сектору в економіці України. Це суттєво впливає не лише на розвиток СВБ в Україні, але й на рівень інвестиційної привабливості українських ФК для іноземних інвесторів. Спрямовувати бізнес до СВД необхідно та можливо шляхом скасування перешкод для становлення СВБ, пов'язаних з відсутністю стимулювання СВД підприємств із боку державних органів. Дана перешкода є причиною виникнення іншої перешкоди, яка полягає, на думку експертів, у відстороненості бізнес-структур від соціальних проблем життя українського суспільства та не сформованості стандартів участі в ньому. Додатково експерти констатують нерозвиненість механізмів участі бізнес-структур у забезпеченні соціально-економічного розвитку територій присутності. Чинний Податковий кодекс передбачає певні преференції відносно соціально відповідальних підприємців, однак вони не носять системно-орієнтованого характеру. Крім того пільги, що існують, в основному призначені для стимулювання добродійності відносно організацій, що утримуються за рахунок бюджету, осіб з фізичними обмеженнями та ін. Отже,

необхідність зваженої державної політики в сфері управління СВБ в Україні очевидна. Держава повинна активно підтримувати процес впровадження КСВ в українську ділову практику, як шляхом декларації, так і введенням різного роду заохочувальних заходів. Держава має створювати сприятливі умови для розвитку КСВ в Україні, виступаючи в ролі головного мотиватора СВД вітчизняних компаній. Тобто держава, діючи через відповідний механізм регулювання СВБ, повинна виступати локомотивом у питаннях вибудовування системи СВБ і забезпечення умов для сталого розвитку як суб'єктів господарювання, так і економіки в цілому. Схема запропонованого нами механізму державного регулювання СВФБ зображено на рис. 4. На основі висновків фахівці Світового банку нами сформульована послідовність дій щодо формування механізму державного регулювання СВФБ в Україні.

Перший етап у наведеному алгоритмі – це визнання урядом країни важливості СВФБ і переконання в цьому бізнесу – полягає в тому, що уряд повинен визнати переваги СВД ФК, що сприяють підвищенню як конкурентоспроможності вітчизняної фармації, так і держави, і демонструвати свої погляди, привселюдно виражаючи своє позитивне відношення до компаній, які впроваджують концепцію СВФБ. При цьому можуть бути використані такі засоби, як різного роду інформаційні та суспільні кампанії та заходи, як наприклад: розробка загальнонаціонального сайту для інформаційного супроводу поточних державних ініціатив і програм в сфері СВФБ; підтримка роботи компаній-піонерів у сфері СВФБ, що забезпечить наочний приклад для інших ФК; визначення кола національних інтересів в галузі СВФБ за допомогою проведення на всій території країни серії зустрічей з лідерами фармацевтичного бізнесу для вироблення кращого розуміння масштабів завдань і обліку всього багатства ідей, досвіду та точок зору. Це допоможе створити надійну основу для подальшого діалогу та співробітництва ФК і держави.

Другий етап – забезпечення підтримки участі ФК в ключових ініціативах, яке може приймати різні форми, включаючи державно-приватні партнерства, фінансові інвестиції в проекти, створення національної мережі топ-менеджерів із СВ тощо. На цьому етапі держава може здійснити наступні заході: збільшувати вигоди для ФК від участі в соціально орієнтованій діяльності; заснувати та присуджувати розпізнавальні знаки соціально відповідальної ФК; розширювати доступ таких ФК до необхідних ресурсів.

Третій етап – створення рекомендацій, інформаційна та ресурсна підтримка ФК на рівні державних інститутів – взаємозалежний із другим кроком. Його основна ідея в тому, що держава повинна дати бізнесу рекомендації про те, як розбудовувати соціальний та екологічний напрямки діяльності, а також запропонувати ресурси й подальшу допомогу.




Рис. 4 Механізм державного регулювання СВФБ.

Четвертий етап передбачає створення збірки правил в країні, які містять у собі розробку форматів документів, придатних для вимірювання діяльності компаній в сфері сталого розвитку. Мається на увазі, що держава бере участь у визначенні орієнтирів для добровільних звітів, оцінці міжнародних кодексів поведінки й огляді кодексів поведінки компаній. Уряд, що виступає в ролі ключового гравця та регулятора, може також відігравати роль оптимізатора діалогу між стейкхолдерами. Таким чином, це допоможе ФК активніше впроваджувати принципи СВ, примиряючи всі залучені сторони, що сприятиме виробленню оптимальних рішень.

Створення сприятливого для СВФБ середовища (п'ятий етап) – належить до тих дій, які є найбільш важливими для створення позитивних стимулів для розвитку практики СВ у фармації.

Шостий етап – розвиток нормативної бази, необхідної для підготовки компаніями соціальної звітності та присвоєння відповідних рейтингів. Стандарти звітності й рейтинги дозволяють одержати кількісну та якісну оцінку діяльності різних компаній, і в тому числі фармацевтичних, в сфері КСВ та відкривають можливості для регіонального і міжгалузевого порівняльного аналізу. Тим самим вони стимулюють конкуренцію, і відповідно, більш активний розвиток соціально відповідальної діяльності. У зв'язку із цим уряд повинен брати активну участь у заходах, націлених на поліпшення методів звітності та присвоєння рейтингів, як на національному, так і на міжнародному рівнях.

Сьомий етап передбачає проведення консультацій з ФК , неурядовими організаціями та іншими групами та допомогу у створенні коаліції для формування механізмів державного регулювання СВФБ. Сприяння в створенні державно-приватного партнерства у фармації є одним з важливих напрямків діяльності держави щодо стимулювання активності ФК в сфері СВ.

Таким чином, можна констатувати, що державне регулювання СВФБ в умовах демократизації має забезпечувати розвиток партнерських відносин між владою і бізнесом. Тому актуальною є організація процесу відходу від прямого втручання в різні сфери економічної діяльності ФК і переходу до регулюючої ролі інституцій управління. Завдання держави у сприянні розвитку СВФБ полягає у поєднанні ініціативних заходів державної політики із застосуванням, у разі необхідності, засобів додаткового регулювання як інструментів стимулюючого характеру.

Розвитку СВФБ буде сприяти взаємодія компаній з професійними громадськими організаціями (ГО). Вона може здійснюватися на основі більш широкого залучення ГО до розв’язання соціальних проблем і стимулювання їхньої активності через: створення ГО фінансової основи СВФБ – галузевого фонду для проведення конкурсів ініціативних соціальних проєктів; участь представників ГО у конкурсних комісіях при підведенні підсумків конкурсів; пряме фінансування ФК окремих громадських організацій, певних соціальних проєктів на безконкурсній основі; кадрову підтримку ГО тих чи інших соціальних проєктів (включаючи волонтерство); фінансовий та управлінський консалтинг гро-

мадських ініціатив, що практикується в західній практиці.

Окремо слід наголосити на неспроможності бізнесу ефективно розвиватись без якісного кадрового забезпечення, без персоналу, який усвідомлює значущість соціально відповідальної поведінки та дотримується її у виробничій та трудовій сферах, відповідно управлінські кадри повинні організувати процес виробництва та праці таким чином, що соціальна відповідальність стає запорукою ефективного функціонування бізнес-компанії. Нагальною є професіоналізація соціально відповідального бізнесу шляхом створення систем професійної підготовки, сертифікації й акредитації фахівців у цій галузі. Як вже зазначалося, в Україні у цьому напрямі зроблено певні кроки. Так, у Державному класифікаторі професій ДК 003 : 2010 регламентується підготовка у вищих навчальних закладах України фахівців із професій «Експерт із соціальної відповідальності», «Менеджер (управитель) із соціальної та корпоративної відповідальності», «Соціальний менеджер». Втім, подальшого процесу у розв'язанні даної проблеми нами не спостерігається.

Державі необхідно створювати умови для збільшення недержавних інвестицій у систему освіти у сфері СВБ. Необхідно забезпечити підготовку відповідного інструментарію управління СВБ, що передбачає переклад та адаптацію існуючих інструментів СВБ (тренінгових матеріалів, он-лайн засобів та систем, систем самооцінки компаній, публікацій та он-лайн ресурсів, зокрема підготовлених в рамках проєктів ЄС). Дані заходи можуть виконуватись за рахунок співфінансування державою, міжнародними донорами, українськими профільними організаціями із СВБ та професійними ГО.

Необхідно також розвивати інститут галузевого саморегулювання СВФБ. З цією метою слід силами профільних асоціацій розробляти й пропагувати прийняття професійних стандартів.

Але дуже важливою є вимога стосовно того, що органам державної влади для регулювання СВБ слід застосовувати й так зване «рекомендаційне право». Тому прийняття ФК різноманітних етичних актів має бути добровільним, заснованим на прагненні підвищити свою привабливість для зацікавлених сторін.

Висновки: аналіз практики державного регулювання СВБ в Україні виявив те, що в цьому питанні країна лише починає свій шлях. Визначена періодизація розвитку державного регулювання СВБ в Україні. Окреслені основні фактори, що визначають специфічність формування соціально відповідальних практик в країні. Із залученням керівників, топ-менеджерів провідних ФК і за підтримки Українського союзу промисловців та підприємців проведене дослідження відносно перешкод та перспектив становлення СВБ в Україні. Доведено, що держава повинна активно підтримувати процес впровадження КСВ в українську ділову практику, як шляхом декларування, так і завдяки введенню різного роду заохочувальних заходів. Побудована інституціональна вертикаль державного регулювання СВФБ із залученням як органів влади, так і професійних спілок, профільних асоціацій та громадських об'єднань тощо, які повинні активно залучатися до процесу формування і розвитку СВФБ

в Україні. Запропоновані заходи з удосконалення державного регулювання СВФБ в Україні, що сприятиме не тільки підвищенню інвестиційної привабливості вітчизняних ФК для інвесторів, посиленню їх конкурентоспроможності, як на внутрішньому, так і на міжнародному фармацевтичних ринках, набуттю статусу інноваційних, соціально відповідальних організацій, які ефективно вибудовують партнерські відносини і виконують свої зобов'язання перед споживачами, власними працівниками, суспільством, що в кінцевому підсумку, сприятиме підвищенню ефективності фармацевтичного забезпечення населення України. Здійснено моделювання впливу впровадження механізму державного регулювання СВФБ і формування ДПСП на рівень сталого розвитку вітчизняних ФК. Розраховано очікуване зростання інтегрального показника сталого розвитку досліджуваних ФК за умов реалізації запропонованих заходів.

Перелік використаних джерел інформації:

1. Громадська спілка «Українська асоціація досконалості та якості». URL: https://www.uaq.org.ua/index.php/uk/ (дата звернення: 20.09.2024).
2. Український союз промисловців і підприємців : офіційний інтернет-ресурс. URL: https://uspp.ua/ (дата звернення: 20.09.2024).
3. Колповська А. М. Періодизація розвитку концепції соціальної відповідальності в Україні. URL: http://www.dbuapa.dp.ua/zbirnik/2015–01(13)/5.pdf (дата звернення: 20.09.2024).
4. Рижко А. Концепція соціальної відповідальності: етапи становлення та перспективи впровадження в муніципальному управлінні. URL: http://www.dridu.dp.ua/vidavnictvo/2011/2011_04(11)/11ramvmu.pdf (дата звернення: 20.09.2024).
5. Про соціальний діалог в Україні. URL: https://zakon.rada.gov.ua/laws/show/2862–17#Text (дата звернення: 20.09.2024).
6. Третяк М. В. Механізми державного регулювання соціальної відповідальності бізнесу : дис. на здобуття наук. ступ. канд. наук з держ. управління. 2015. 54 с.
7. Братішко Ю. С., Посилкіна О. В. Організаційне забезпечення соціально відповідальної діяльності фармацевтичних підприємств : наук. метод. рек. Харків, 2019. 22 с.
8. Братішко Ю. С., Посилкіна О. В. Формування компетентностей менеджера із соціальної відповідальності фармацевтичного підприємства : наук. метод. рек. Харків, 2019. 22 с.
9. Братішко Ю. С., Посилкіна О. В. Науково-методичні підходи до ідентифікації і оцінки ризиків, пов'язаних із соціально відповідальною діяльністю фармацевтичних компаній : наук. метод. рек. Харків, 2019. 22 с.
10. Братішко Ю. С., Посилкіна О. В. Науково-методичні підходи до розробки внутрішнього стандарту із корпоративної соціальної відповідальності суб'єктів фармацевтичного ринку : наук. метод. рек. Харків, 2019. 22 с.
11. Братішко Ю. С., Посилкіна О. В. Формування системи управління соціальною відповідальністю на вітчизняних фармацевтичних підприємствах : метод. рек. Київ, 2018. 24 с.

ПРОБЛЕМИ ТА ПЕРСПЕКТИВИ РЕЦЕПТУРНОГО ВІДПУСКУ ЛІКАРСЬКИХ ЗАСОБІВ В УКРАЇНІ

Назаркіна В. М., Іванова А. Д.
Національний фармацевтичний університет,
м. Харків, Україна,
Кафедра соціальної фармації
an.ivanova2014@gmail.com

Вступ. Рецептурний відпуск лікарських засобів (ЛЗ) є важливим елементом системи охорони здоров'я, що забезпечує раціональне та контрольоване використання ліків. Система рецептурного відпуску сприяє зниженню ризиків самолікування, захищає пацієнтів від побічних ефектів та запобігає зловживанню препаратами. В Україні система рецептурного відпуску має ряд викликів, які зумовлюють потребу в удосконаленні законодавства, впровадженні сучасних технологій та адаптації до нових умов, таких як вплив війни. Розвиток цієї системи має важливе значення для збереження здоров'я нації та покращення якості медичної та фармацевтичної допомоги.

Мета. Аналіз основних проблем, пов'язаних із рецептурним відпуском ЛЗ в Україні, та визначення перспективних напрямків для оптимізації цієї системи.

Матеріали та методи. Використано аналіз національних і міжнародних нормативно-правових актів, що регулюють рецептурний відпуск ліків, а також вивчено наукові статті та звіти з даної тематики.

Результати. Серед основних проблем, що стосуються рецептурного відпуску ЛЗ, варто виділити неконтрольований продаж ліків без рецепта, низьку обізнаність пацієнтів, корупційні ризики та недостатній контроль з боку держави.

1. Неконтрольований продаж ЛЗ без рецепта. Незважаючи на чинне законодавство, яке чітко регламентує відпуск окремих категорій ЛЗ лише за рецептом лікаря, в Україні поширеною є практика їх продажу без належних приписів. Це особливо стосується антибіотиків, психотропних засобів та гормональних препаратів. Окрім побічних дій та ускладнень неконтрольоване вживання цих ЛЗ сприяє розвитку цілої низки медико-соціальних проблем, таких як антибіотикорезистентність, звикання та розвиток залежності та ін.

2. Недостатня обізнаність пацієнтів. Багато пацієнтів самостійно займаються самолікуванням, ігноруючи необхідність звернення до лікаря. Це пов'язано з недовірою до медичних послуг, високою вартістю консультацій або недоступністю лікарів у певних регіонах. Через це пацієнти часто купують ЛЗ, які можуть бути неефективними або навіть шкідливими.

3. Конфлікт інтересів і корупційні ризики. На жаль, в Україні має місце практика порушення призначення ліків. Лікарі можуть виписувати рецепти на ліки певних фармацевтичних компаній, отримуючи за це винагороду. У зв'язку

з цим було впроваджено стандарти лікування (клінічні протоколи, настанови), а також вимога щодо призначення ліків за МНН.

4. Незадовільний контроль з боку держави. Державний контроль за дотриманням правил відпуску ЛЗ в аптеках є недостатнім. Мораторій на перевірки та невисокі штрафи за порушення у цій сфері роблять систему відпуску ЛЗ вразливою до зловживань.

Перспективи вдосконалення рецептурного відпуску ЛЗ:

У рамках покращення рецептурного відпуску ЛЗ необхідно впроваджувати сучасні інноваційні рішення, що передбачають електронні рецепти, посилення контролю за аптеками, освітні програми для населення, національні програми з моніторингу та співпрацю з міжнародними організаціями.

1. Впровадження електронного рецепта. Е-рецепти можуть стати дієвим інструментом у боротьбі з неконтрольованим продажем ЛЗ. Така система дозволить знизити ризики підробок рецептів, а також дасть змогу краще контролювати процес відпуску ЛЗ, оскільки всі дані будуть зберігатися у єдиній електронній базі.

2. Посилення контролю у сфері призначення і використання ЛЗ. Доцільно посилити державний контроль за призначенням і відпуском ЛЗ, впровадити ефективні механізми перевірки правомірності видачі рецептів.

3. Освітні програми для населення. Інформаційні кампанії та освітні програми для населення про важливість правильного використання ЛЗ та профілактичні заходи можуть допомогти зменшити рівень самолікування та підвищити відповідальність пацієнтів за власне здоров’я.

4. Запровадження національних програм з моніторингу. Важливим кроком є створення національної системи моніторингу використання ЛЗ. Це дозволить аналізувати дані щодо відпуску рецептів, виявляти зловживання та оперативно реагувати на проблеми у сфері охорони здоров'я.

5. Співпраця з міжнародними організаціями. Впровадження міжнародного досвіду щодо регулювання рецептурного відпуску ЛЗ може допомогти Україні уникнути помилок інших країн та покращити власну систему. Співпраця з такими організаціями, як ВООЗ, є важливим фактором успішної реформи.

Висновки. Рецептурний відпуск ЛЗ в Україні стикається з численними проблемами, зокрема з недосконалим законодавством, недостатнім контролем за раціональним призначенням та використанням ліків і низьким рівнем обізнаності населення. Впровадження е-рецепта, розвиток телемедицини, підвищення кваліфікації аптечного персоналу та інформаційні кампанії можуть сприяти підвищенню безпеки та якості медичного обслуговування в Україні. Актуальні проблеми та нові підходи до їх вирішення потребують комплексного підходу та співпраці держави, медичних установ і фармацевтичного сектору для забезпечення населення безпечними та ефективними ЛЗ.

АНАЛІЗ ВІДХИЛЕНЬ У ЛИСТАХ ЛІКАРСЬКИХ ПРИЗНАЧЕНЬ У СТАЦІОНАРІ ХІРУРГІЧНОГО ПРОФІЛЮ З МЕТОЮ УДОСКОНАЛЕННЯ СИСТЕМИ УПРАВЛІННЯ ФАРМАЦЕВТИЧНОЮ СЛУЖБОЮ ЗОЗ

Залигіна Є. В.

Дніпровський державний медичний університет, м. Дніпро, Україна

Кафедра фармакології, загальної та клінічної фармації

avis.rara@hotmail.com

Вступ. При наданні медичної допомоги, шкода здоров'ю пацієнтів, що пов'язана з застосуванням лікарських препаратів, зустрічається більш ніж у чверті призначень. Питання забезпечення безпеки у сфері охорони здоров'я є глобальною проблемою. Проблема підвищення якості надання фармацевтичної допомоги та оцінки витрат на медичне забезпечення є актуальною та потребує подальшого дослідження.

Мета. удосконалення фармацевтичної служби у закладі охорони здоров'я шляхом аналізу відхилень у призначені лікарських препаратів.

Матеріали та методи. Дослідження проведено на базі ТОВ «Ендотехномед» хірургічна клініка «Garvis». Було розраховано рівні відхилень у листах призначення, проаналізовано їх динаміку та зв'язок з витратами на медичне забезпечення. Статистичну обробку результатів проводили з використанням методів описової та аналітичної статистики за допомогою відкритого інтегрованого середовища розробки Jupyter Notebook. Значення $p<0,05$ вважалося статистично значущим при перевірці всіх статистичних гіпотез.

Результати. Рівень відхилень у призначеннях ЛЗ за 4,5 роки достовірно знизився у 4,6 разів. Це узгоджується з результатами досліджень китайських вчених, які доводять, що правильно організована робота у стаціонарі ЗОЗ дозволяє скорочувати частоту відхилень при призначенні ЛЗ у 3,58 разів.

Між розміром витрат на медичне забезпечення одного пацієнта та рівнем відхилень у призначеннях ЛЗ спостерігалася негативна рангова статистично значуща кореляція середньої сили. Отже, зниження помилок у призначеннях ЛЗ напряму впливає на рівень якості надання медичної допомоги, у той же час витрати на медичне забезпечення більшою мірою пов'язані з зовнішніми чинниками, вплив на них з боку внутрішніх управлінських рішень є обмеженим.

Висновки. Впровадження нових підходів контролю призначень лікарських препаратів призвело до достовірного скорочення відхилень у листах призначення у 4,6 разів у порівнянні з початковим рівнем. Вищому рівню відхилень у листах призначення відповідають нижчі витрати на медичне забезпечення. Удосконалення процесів управління фармацевтичною службою було направлено на планування, контроль, запровадження і функціонування інформаційної системи, управління персоналом, оцінювання ефективності, залученням стейкхолдерів для формування й реалізації рішень.

МОЖЛИВІСТЬ ІМПЛЕМЕНТАЦІЇ ПРАКТИКИ МЕДИКАМЕНТОЗНОГО УЗГОДЖЕННЯ У ПРОФЕСІЙНУ ДІЯЛЬНІСТЬ КЛІНІЧНИХ ФАРМАЦЕВТІВ В УКРАЇНІ

Зіменковський А. Б., Шпортько М. О., Настюха Ю. С.

Львівський національний медичний університет імені Данила Галицького, Кафедра менеджменту в охороні здоров'я, фармакотерапії і клінічної фармації, м. Львів, Україна

y.nastyukha@gmail.com

Вступ. У закордонних джерелах термін медикаментозні розбіжності (МР) (Medication discrepancy) застосовується для опису будь-яких відмінностей між інформацією медикаментозного анамнезу та порядком застосування ліків пацієнтом при госпіталізації *(World Health Organization, WHO, 2019)* чи між двома переліками лікарських засобів (ЛЗ) на етапі скерування або переведення пацієнта до закладу охорони здоров'я (ЗОЗ) чи фахівця, який надає медичну допомогу *(Almanasreh E., et al., 2020)*. При згаданому переведенні такі ненавмисні розбіжності виникають майже у кожного пацієнта. Відповідно до звіту ВООЗ проблема виникнення МР носить глобальний характер. Визначення поширеності проблеми ускладнено відмінностями у трактуванні термінів, методах виявлення та системах класифікації. Проте, результати закордонних досліджень засвідчують, що принаймні один випадок МР виявляється при госпіталізації у 3-97% дорослих пацієнтів і 22-72% пацієнтів дитячого віку та при виписуванні – у 25-80% пацієнтів. Щонайменше один випадок ненавмисного виникнення МР трапляється у 62% пацієнтів під час переведення між відділеннями лікарні. МР виникають у 27-57% літніх людей, які перебувають у геронтологічних будинках, та 14-98% тих, хто проживає у громаді *(WHO, 2019)*. Загалом, при переведенні людей з одного ЗОЗ чи підрозділу до іншого у 15-70% пацієнтів виникають помилки або ненавмисні зміни у застосуванні ліків *(National Institute for Health and Care Excellence, NICE, 2015; WHO, 2023)*.

Мета. Метою дослідження стало вивчення можливості імплементації практики медикаментозного узгодження у професійну діяльність клінічних фармацевтів в Україні.

Матеріали та методи. Використано методи: системного підходу, бібліографічний, порівняльного аналізу. Об'єкти дослідження: Настанова Національного інституту здоров'я і досконалості допомоги Великої Британії щодо оптимізації застосування ліків *(Medicines optimisation: the safe and effective use of medicines to enable the best possible outcomes, National Institute for Health and Care Excellence, NICE, 2015)*; Звіт ВООЗ з питань медикаментозної безпеки при переведенні пацієнта до ЗОЗ чи фахівця, який надає медичну допомогу *(Medication Safety in Transitions of Care. Geneva: World Health Organization, WHO, 2019)*; Класифікаційна система лікопов'язаних проблем Європейської мережі фармацевтичної опіки V9.1 *(Pharmaceutical Care Network Europe, PCNE, 2020)*; Довідник кваліфікаційних характеристик професій працівників. Випуск

78. Охорона здоров'я (ОЗ) *(Наказ МОЗ України від 29.03.2002 №117)*.

Результати. Переведення пацієнта з одного до іншого ЗОЗ чи фахівця, який надає медичну допомогу, часто пов'язане із внесенням змін у застосування ліків. Тож, саме на цьому етапі виникають проблеми МР, зокрема, упущення пов'язані з призначенням чи відміною ЛЗ та неналежна комунікація щодо внесених змін. Вирішенню проблеми сприяє медикаментозне узгодження.

Медикаментозне узгодження (Medication reconciliation) – це процес визначення точного чинного переліку ліків пацієнта та його порівняння із поточним переліком застосовуваних ЛЗ, виявлення будь-яких розбіжностей та документування будь-яких змін, що призводить до остаточного отримання повного переліку ліків та точної комунікації щодо них *(NICE, 2015)*.

Для вивчення можливості імплементації такої практики нами проаналізовані закордонні рекомендації та вимоги до проведення медикаментозного узгодження та затверджені в Україні професійні обов'язки клінічних фармацевтів.

При зверненні пацієнта чи при його госпіталізації необхідно структуровано провести опитування для отримання інформації про всі ЛЗ, які він застосовує; підтвердити, надану пацієнтом інформацію, щонайменше одним надійним джерелом; порівняти зібраний медикаментозний анамнез (щодо рецептурних та безрецептурних ліків) із поточними призначеннями ЛЗ для виявлення та вирішення будь-яких розбіжностей; надати остаточний перелік ліків із призначеної схеми фармакотерапії (ФТ) пацієнту та іншим фахівцям ОЗ, дотичним до ФТ *(WHO, 2019)*. При виписуванні або переведенні пацієнта перелік ліків оновлюється із врахуванням ЛЗ, які пацієнт буде застосовувати надалі (зокрема, амбулаторно). Проведення ретельного збору медикаментозного анамнезу, зокрема шляхом комунікації з пацієнтом, відповідає професійним обов'язкам клінічного фармацевта.

Про клінічне значення МР свідчить включення такої причини виникнення лікопов'язаних проблем до Класифікаційної системи Європейської мережі фармацевтичної опіки V9.1 *(PCNE, 2020)*. Кваліфікаційні вимоги до фахівця, який може здійснювати медикаментозне узгодження, окрім навичок ефективної комунікації та знань технічних процесів менеджменту ліків, також вимагають компетентності у питаннях клінічного застосування ЛЗ *(NICE, 2015)*. Таким кваліфікаційним характеристикам відповідає клінічний фармацевт. Залучення клінічного фармацевта, у порівнянні з проведенням медикаментозного узгодження лише лікарем, доцільне також з огляду на ресурсозатратність процесу. При гострих станах воно повинно здійснюватися упродовж 24 годин або й швидше *(NICE, 2015)*, в той час, коли при зверненні чи поступленні пацієнта такий процес потребує від 15-30 хв на втручання *(WHO, 2019)*.

Висновок. Результати порівняльного аналізу закордонних рекомендацій щодо проведення медикаментозного узгодження та вітчизняних професійних обов'язків клінічного фармацевта засвідчують можливість залучення цього фахівця до вирішення МР та попередження виникнення лікопов'язаних проблем, зокрема при переведенні пацієнта до іншого ЗОЗ чи фахівця ОЗ.

ANALYSIS OF ETHICAL ASPECTS OF PHARMACISTS' WORK

Chehrynets A. A., Bondarieva I. V., Khaleq S.
National University of Pharmacy Kharkiv, Ukraine
Department of Management, Marketing and Quality Assurance in Pharmacy
fmmqaph@nuph.edu.ua

Introduction. Pharmacists hold a pivotal position within the healthcare system, functioning as key facilitators in the management of pharmacotherapy between healthcare providers and patients. Their professional obligations extend beyond the mere dispensation of medications; they are tasked with ensuring the optimal efficacy and safety of prescribed therapies, delivering accurate and comprehensive pharmaceutical care, and making decisions rooted in ethical principles. The ethical dimensions of pharmacists' responsibilities encompass critical issues such as patient confidentiality, informed consent, the management of conflicts of interest, and strict adherence to both legal and professional regulatory frameworks.

Aim. of this study is to analyze ethical aspects of pharmacists' work.

Materials and methods. This analysis is based on a comprehensive review of current literature on pharmacy ethics, professional guidelines, and surveys conducted among practicing pharmacists. Primary data was gathered through questionnaires distributed to a representative sample of pharmacists, which focused on their experiences with ethical dilemmas.

Results and their discussion. The findings revealed that pharmacists often encounter ethical dilemmas in areas such as patient confidentiality, conflicts of interest, and the dispensing of medications that may contradict personal beliefs. For example, pharmacists must often balance patient privacy with the need to communicate important health information to other healthcare providers. Additionally, the pressure to meet business goals in a retail setting can sometimes conflict with the ethical obligation to prioritize patient care. Pharmacists also reported facing challenges when dealing with prescriptions for controversial medications, such as those used in end-of-life care or reproductive health, where their personal moral values may conflict with their professional duties. The study highlighted the importance of ethical training in pharmacy education, as many respondents felt that their formal training did not fully prepare them for these situations. Professional guidelines and codes of conduct were found to provide a solid foundation for navigating these challenges. However, pharmacists emphasized the need for ongoing education and support in handling ethical issues, particularly as healthcare systems and societal expectations evolve.

Conclusions. The ethical aspects of pharmacists' work are multifaceted and often complex, requiring a balance between legal, professional, and personal considerations. The study concluded that while pharmacists are generally equipped to handle ethical challenges, there is a need for more comprehensive ethics training and better support systems in the workplace. Ensuring that pharmacists have the tools and knowledge to make ethically decisions is essential for maintaining the integrity of the profession and the trust of patients.

ОЦІНКА СПОЖИВАННЯ В УКРАЇНІ ЛІКАРСЬКИХ ЗАСОБІВ, ЩО ПРИГНІЧУЮТЬ ПЕРИСТАЛЬТИКУ

Герасимова О. О., Ткачова О. В.
Національний фармацевтичний університет, м. Харків, Україна
o.a.gerasymova@gmail.com

Вступ. Діарея є однією з найчастіших причин звернення пацієнтів за медичною допомогою. Вона виникає через різноманітні причини, може бути симптомом гострих та хронічних захворювань та варіювати від легкого, тимчасового стану до потенційно небезпечного для життя. Для зменшення частоти кишкових випорожнень та проявів діареї призначають лікарські засоби (ЛЗ), що пригнічують перистальтику.

Мета. Оцінка обсягу та структури споживання ЛЗ групи A07D «Засоби, що пригнічують перистальтику» в Україні в 2021-2022 роках.

Матеріали та методи. Об'єкти дослідження – представники лікарських засобів групи A07D «Засоби, що пригнічують перистальтику». Оцінку споживання ЛЗ проводили за допомогою ATC/DDD-методології, яка включає дві складові: анатомо-терапевтично-хімічну класифікацію лікарських засобів та їх середні підтримуючі добові дози, що застосовуються за основним показанням у дорослих (DDD – defined daily dose). Під час проведення дослідження розраховували міжнародний показник DDDs/1000 жителів/день (DID). Кількість міжнародних непатентованих назв (МНН), торгових найменувань та реалізованих за рік упаковок ЛЗ в Україні була взята з інформаційної системи «Pharmexplorer» компанії «Моріон». Дози DDD препаратів взяті із сайту ВООЗ (https://www.whocc.no/). За їх відсутності розраховували середню добову дозу PDD (prescribed daily dose) за допомогою інструкції до медичного застосування.

Результати. Група A07D «Засоби, що пригнічують перистальтику» була представлена в Україні у 2021-2022 роках 2 МНН (лоперамід та «лоперамід + симетикон») та 17 торговими найменуваннями. Обсяги споживання ЛЗ даної групи за показником DDDs/1000 жителів/день у 2021-2022 роках в Україні складали, відповідно, 1,50 DID та 1,43 DID. Найбільший рівень споживання у досліджуваний період встановлено для препаратів лопераміду (2021 рік – 1,48 DID; 2022 рік – 1,41 DID), найменший – для комбінації «лоперамід + симетикон» (2021-2022 роки – по 0,02 DID).

Висновки. В Україні у 2021-2022 роках приблизно від 0,14 % до 0,15 % населення щодня приймали 1 DDD одного з ЛЗ групи A07D «Засоби, що пригнічують перистальтику». З роками обсяги споживання ЛЗ даної групи в країні мали тенденцію до зменшення. Результати даного дослідження можуть стати в нагоді організаторам охорони здоров'я при прийнятті регуляторних рішень щодо раціонального використання представників групи A07D «Засоби, що пригнічують перистальтику» та реімбурсації їх вартості.

UDC 615.12: 339.137

STUDY OF CASES OF SUCCESSFUL USE OF SOCIAL MEDIA BY PHARMACEUTICAL COMPANIES

Bondarieva I. V., Malyi V. V., Ifrad Yu.
National University of Pharmacy Kharkiv, Ukraine
Department of Management, Marketing and Quality Assurance in Pharmacy
iryna.bondarieva@gmail.com

Abstract. Goal. The main objective of this study is to analyze successful cases of effective use of social media platforms by pharmaceutical companies for marketing, brand development, consumer education and customer interaction.

Materials and methods. The research was conducted using the method of content analysis.

The results. Pharmaceutical companies are successfully using social media to improve customer service, raise health awareness and build communities.

Conclusions. Companies that use social media effectively improve their reputation, increase customer engagement and provide better communication with consumers and healthcare professionals.

Keywords: *social media, pharmaceutical companies, marketing, consumer education, communication.*

Анотація. Мета. Головною метою цього дослідження є аналіз успішних випадків ефективного використання фармацевтичними компаніями платформ соціальних медіа для маркетингу, розвитку бренду, освіти споживачів і взаємодії з клієнтами.

Матеріали та методи. Дослідження було проведено за допомогою методу контент-аналізу.

Результати. Фармацевтичні компанії успішно використовують соціальні медіа для покращення обслуговування клієнтів, підвищення обізнаності про здоров'я та побудови спільнот.

Висновки. Компанії, які ефективно використовують соціальні медіа, покращують свою репутацію, підвищити залучення клієнтів і забезпечити кращу комунікацію з споживачами та медичними працівниками.

Ключові слова: *соціальні медіа, фармацевтичні компанії, маркетинг, освіта споживачів, комунікація.*

Introduction. In the era of digital transformation, social media has become an essential tool for various industries, including pharmaceuticals. Traditionally, the pharmaceutical sector relied heavily on healthcare professionals, direct advertising, and regulatory channels to market products and build customer relationships. However, with the rise of social media, pharmaceutical companies now have new opportunities to engage with both consumers and healthcare professionals. The ability to disseminate health information quickly, connect with a wide audience, and respond to

patient inquiries in real time has transformed how pharmaceutical companies position their products and brands. Despite the stringent regulatory environment surrounding pharmaceutical communications, many companies have leveraged social media to create successful campaigns that build trust, enhance brand visibility, and improve patient engagement. Thus, examining the cases of successful social media use in the pharmaceutical industry is relevant and crucial to understanding best practices and overcoming challenges.

Aim. The main purpose of this study is to analyze successful cases where pharmaceutical companies have effectively utilized social media platforms for marketing, brand development, patient education, and customer engagement. The study will identify key factors that contribute to the success of these campaigns, explore how companies navigate regulatory constraints, and highlight the types of social media content that resonate with different target audiences. Additionally, the study aims to provide actionable insights and recommendations for other pharmaceutical companies looking to optimize their social media strategies.

Materials and methods. This study was conducted through using content-analysis method, a qualitative analysis of various case studies focusing on successful social media campaigns implemented by leading pharmaceutical companies. The analysis involved a review of social media activity, content types, and audience engagement metrics across multiple platforms, including Facebook, Instagram, Twitter, LinkedIn, and YouTube. Data were collected from industry reports, social media analytics, and company press releases to understand the strategies employed and the outcomes achieved.

Results and their discussion. Several pharmaceutical companies have successfully used social media platforms like Facebook, Instagram, Twitter, LinkedIn, and YouTube to achieve a variety of strategic goals.

Pfizer has effectively used social media to launch public health awareness campaigns, particularly during the COVID-19 pandemic. The company utilized platforms such as Twitter and Facebook to provide accurate and timely information about vaccine safety and efficacy. Their strategic use of infographics, Q&A sessions, and partnerships with medical influencers boosted engagement and trust among both healthcare professionals and the general public.

Maintaining an active presence on platforms like Twitter, Facebook, and Instagram, Pfizer shares educational content on various health topics, including vaccine information, disease prevention, and medication safety. They also engage with patients through Q&A sessions and live events, fostering an interactive dialogue.

As a result, Pfizer has built a strong online community of patients and healthcare professionals who trust and rely on their information. This approach has not only improved brand loyalty but also enhanced the company's reputation in the healthcare industry.

Johnson & Johnson has focused on creating communities and providing patient support through social media. One of their most successful initiatives involved engaging with cancer patients and survivors via YouTube and Facebook, where they

shared stories of hope, provided educational videos about treatment options, and created forums for patients to share their experiences. This community-building approach not only humanized the brand but also provided valuable support for patients and caregivers.

In addition to community engagement, Johnson & Johnson has produced a wide range of educational content on various health topics, including diabetes, skin care, and maternal and child health. This content is shared through social media channels and distributed through partnerships with healthcare providers and patient advocacy groups. As a result, Johnson & Johnson has established itself as a thought leader in the healthcare industry, and their educational initiatives have contributed to improved public health outcomes.

Bayer has integrated influencer marketing into their social media strategy to reach younger audiences. By collaborating with healthcare influencers and lifestyle bloggers, Bayer was able to successfully promote its consumer health products on Instagram. These partnerships created authentic content that resonated with the audience, driving brand awareness and product usage among new demographics.

GlaxoSmithKline (GSK) has effectively harnessed social media to improve customer service and raise awareness about mental health issues through their #MentalHealthMatters campaign. By utilizing platforms like Twitter, Facebook, and Instagram, GSK developed a direct line of communication with healthcare providers and patients. This strategy allowed them to quickly address questions related to medications, side effects, and product availability, ultimately enhancing customer service and improving patient satisfaction.

In addition to customer service, GSK aimed to reduce stigma around mental health and encourage individuals to seek help. They created engaging content, shared personal stories, and partnered with mental health organizations to foster important conversations. The campaign was a resounding success, reaching millions and generating significant discussions about mental health. GSK's commitment to social responsibility in this area was widely recognized, further enhancing their brand reputation.

Sanofi has created online communities and forums where patients with specific health conditions can connect, share experiences, and seek support. Among the benefits were: these communities provide a valuable resource for patients, fostering a sense of belonging and reducing feelings of isolation. They also offer valuable insights to Sanofi, helping them to better understand patient needs and improve their products and services.

During the COVID-19 pandemic, AstraZeneca faced significant challenges related to the development and distribution of its vaccine. The company used social media to communicate updates about vaccine development, address public concerns, and provide accurate information. AstraZeneca's transparent and timely communication helped to maintain public trust and confidence in their vaccine, despite facing criticism and challenges.

The study of successful social media use by pharmaceutical companies reveals that social media can be a powerful tool when used strategically. The key to success lies in developing content that is not only informative but also engaging, trustworthy, and compliant with regulatory standards. Companies that have excelled in this space have focused on building relationships with their audience, offering valuable educational content, and creating online communities that foster trust and loyalty [1, 2].

Ukrainian pharmaceutical companies have increasingly recognized the power of social media to connect with their audience, build trust, and enhance their brand image. By leveraging these digital platforms, they have been able to overcome language barriers, reach a wider audience, and foster a sense of community.

Farmak is a leading Ukrainian pharmaceutical company has successfully utilized Facebook and Instagram to create a loyal community of followers. By sharing informative content on health topics, engaging with users through Q&A sessions, and running contests, Farmak has solidified its position as a trusted source of healthcare information.

Darnitsa has employed social media to raise awareness about important social issues, such as mental health. Through well-crafted campaigns and engaging content, they have not only boosted their brand reputation but also contributed to a positive social impact.

By focusing on providing high-quality content related to various health conditions, Lekhim has positioned itself as a healthcare expert. Their educational materials and informative posts have helped build trust among patients and healthcare professionals alike.

During the COVID-19 pandemic, Boris demonstrated the importance of timely and accurate communication on social media. By sharing updates on their products and services and addressing public concerns, they were able to maintain customer trust and loyalty.

Recognizing the influence of influencers, Biofarma has partnered with Ukrainian health and lifestyle influencers to reach a wider audience. This strategy has helped increase brand awareness and credibility.

By creating online communities where patients can connect and share experiences, Europharm has fostered a sense of belonging among its customers. These communities have not only helped build trust but also provided valuable insights into patient needs.

Ukrainian pharmaceutical companies that have succeeded in social media have maintained a consistent presence and engaged with their audience regularly. Sharing valuable content, such as health tips, educational materials, and updates on industry news, has been key to building trust and credibility. Encouraging dialogue and responding to comments and messages has helped create a more personal connection with customers. Effective use of social media has been crucial in managing crises and maintaining public trust during challenging times. Staying up-to-date with the latest trends and platforms has allowed these companies to reach a wider audience and stay ahead of the competition.

Overall, Ukrainian pharmaceutical companies have demonstrated the significant potential of social media to: Build brand awareness and reputation; enhance patient engagement; Provide valuable health information; Foster a sense of community; manage crises effectively. By continuing to invest in social media and adopting best practices, these companies can further strengthen their position in the market and contribute to improving public health.

Conclusions. Pharmaceutical companies that effectively leverage social media platforms can improve their brand reputation, increase patient engagement, and foster better communication with both consumers and healthcare professionals. Despite challenges such as navigating regulatory restrictions and ensuring content accuracy, the benefits of a well-executed social media strategy are clear. Moving forward, pharmaceutical companies should continue to innovate and refine their social media approaches, integrating new tools such as AI-driven analytics and personalized content to enhance user experience and meet the evolving needs of their audiences.

The insights gained from this work can serve as a roadmap for pharmaceutical companies looking to harness the power of social media to build stronger, more meaningful connections with their stakeholders while maintaining compliance with industry regulations.

References

1. Жадько, С. В. Використання мережі інтернет при просуванні лікарських препаратів до кінцевих споживачів / С. В. Жадько // Управління, економіка та забезпечення якості в фармації. - 2013. - № 2(28). - С. 70-73.
2. Zhadko, S. V. The feasibility of using social media in pharmaceutical marketing / S. V. Zhadko, A. E. Kulibaba, Ali Ahmed Khidaer Ali // Соціальна фармація : стан, проблеми та перспективи : матеріали II міжнар. наук.-практ. інтернет-конф., м. Харків, 27-30 квіт. 2015 р. - Х., 2015. - С. 176-178.

STRATEGIC MANAGEMENT ELEMENTS, USED FOR GEORGIAN PHARMACEUTICAL SECTOR

Kvizhinadze N., Dughashvili N., Instkirveli N., Kiknavelidze N.
Tbilisi State Medical University

Introduction. In the recent history of Georgia, the pharmaceutical sector is one of the most dynamic and rapidly growing markets. At present, "economic agents operating in the pharmaceutical market (directly and/or through interdependent persons) are engaged in economic activity at all levels of the pharmaceutical product supply chain (import, production, wholesale supply, retail sale), as well as operate clinics, insurance companies, etc. As a result, the pharmaceutical market in the country is characterized by a high degree of vertical/horizontal integration and holding arrangement. Tough competition and a constantly changing environment force companies to pay maximum attention to the efficiency of using resources while organizing internal activities.

Strategic management is the art and science of formulating, implementing, and evaluating multi-functional decisions that will enable a company to achieve its goals. The goal of strategic management is to take advantage of today's opportunities and create new and different opportunities for tomorrow. This process can be described as an objective, logical, and systematic approach to making important decisions and allows us to organize quantitative and qualitative information in such a way that effective decisions can be made in an ever-changing environment. The planned process should allow for quick adaptation and changes.

Aim. The study aims to identify examples of strategic management in the Georgian pharmaceutical market through available sources - to determine, analyze, and evaluate the advantages of the managerial vision in the specific case considered, and to use effective methods in one's practice.

Materials and methods. The subject of the research is the study of the specific strategies used in the stages of development of the leading companies of the Georgian pharmaceutical market ("Aversi", "PE-ES-PE", "GPC", "Pharmadepo").
Desk studies and face-to-face interviews have been selected as the main methods of research by the specifics of the topic.

Results and discussion. We started the research from the history of the creation of the main players of the modern Georgian pharmaceutical market and chronologically marked the steps of each of them by analyzing the respective goals and results. Along with this, to present the relatively less visible moments in the dynamics of historical events, we interviewed 16 mid-level managers (marketing, sales, and import managers and division heads) of 8 different pharmaceutical companies operating in the Georgian market who witnessed/participated in these events, for which we prepared a targeted questionnaire. 9 of the mentioned 16 managers have a master's degree in business administration. During the selection of active managers participating in the survey, three main criteria were taken into account: first - at least 15 years of experience as a manager in the Georgian pharmaceutical market, second - equal distribution between managers of global pharmaceutical companies and managers of other pharmaceutical companies represented in the local market (8/8 respectively, 16 in total), the third - the main limiting criterion in the selection of interviewees - the interviewee should not represent the current manager of "Aversi", "PSP" and "Gepha". It is important to note that although many pharmaceutical companies are operating in Georgia today, finding active managers who meet the criteria described above and agree to participate in the survey is not very common. To increase the reliability of the received information, we maintained the ratio between the representatives and managers of other pharmaceutical companies and took into account only their assessments. We did not select representatives of the highest management as subjects of the study due to difficulties in accessing them, especially since the directors of the leading pharmaceutical companies were not involved in the research.
To find out the customer's loyalty to other and other pharmacy chains, specifically to reveal the influence of the price component, we compiled a relevant questionnaire and interviewed 153 random people of the adult population (ages 30-60) face-to-face.

The results of the desk part of the research (literature, each company's development history, annual reports, publications, etc.) and the aforementioned questionnaires are integrated with the discussion of examples of strategic management.
While discussing the examples identified as a result of the research, we tried to protect the dynamics of market development to easily understand the events.

Conclusions. According to the results of the research, we can conclude that the researched pharmaceutical companies used different elements of the "judo strategy" at a specific time and environment throughout their development history.
When researching examples of other elements of "judo strategy", the evidentiary material is relatively less, and according to the consistent dynamic development, it can be assumed that the maneuver used in a certain period may represent an element of the "judo strategy". However, uniquely, each of them is a constituent component of a single strategic management of companies.

PECULIARITIES OF REFERENCE PRICING POLICY IN THE PHARMACEUTICAL SECTOR OF GEORGIA

Shashiashvili N., Kvizhinadze N., Gaprindashvili M., Dughashvili N., Kiknavelidze N., Instkirveli N.
Tbilisi State Medical University

Introduction. One of the main components of healthcare access is physical and geographic access to medical supplies. It is estimated that in developing and poor countries, unregulated prices for pharmaceutical products are a heavy burden on the population. Governments around the world should implement appropriate policies to improve access to medicines. Some widely used strategies include cost control of manufacturer prices, internal and external reference pricing, profit and supply/distribution chain regulation, generic drug promotion, and others.

The high prices of medicines in Georgia were due to the deficiencies in the pharmaceutical sector and the fact that price regulation, rational use of medicines, and price control were not carried out in the country. The reform of reference pricing for medicines came into effect in Georgia on February 15, 2023, and the regulation affected up to 6 thousand medicines in one year. In the period 1 year after the introduction of the reform, it is relevant how the parties involved in the process evaluate the novelty and what challenges the pharmaceutical sector faces in this direction.

Aim. The study aims to study the influence of the results of the reference pricing policy on the proper functioning of the pharmaceutical sector.

Materials and methods. A total of 125 respondents (pharmacists, doctors, patients) participated in the reference price system research. Research period 01.03.2024-26.05.2024. The research subjects were divided into three groups. 16% of respondents were doctors, 32% were pharmacists, and 52% were patients. The survey

was conducted using electronic and written questionnaires, anonymously, in such a way that their constitutional rights were protected. Appropriate questionnaires for the target groups were developed based on the study and analysis of the literature.

Results and discussion. Interviewed pharmacists and doctors note that the reference price system has changed their daily work for the better. According to pharmacists, the existing changes have had a positive effect on their relationship with patients, especially in terms of consultation and offering. The introduction of the reference price in the country led to a change in the price of many medicines, however, doctors note that the current change did not affect the treatment decisions of patients, and caused distrust among patients. There was an opinion that lowering the price of medicines would lead to a change in the prescription of medicines. But the doctors do not confirm this fact and in the survey, they unequivocally point out that the prescription of drugs from the reference list did not affect their prescription. Pharmacists also confirm this. 60% of them report that they have not noticed any changes in the choice of prescribing medicines by medical staff.

Apart from the positive aspects, the changes in the health system, which involve the establishment of a reference price for medicines, have created some challenges in the pharmaceutical sector. Pharmacists, depending on the work process, in the survey mention several important details that are worth noting, these are delays in the supply of medicines (in the background of the shortage, there was an increase in the stocks of medicines supplied from manufacturing companies in Asian countries), risks related to the quality of medicines, reduction in the sale of original medicines, insufficient awareness of patients, which leads to Confusion and mistrust among them, as well as increasing the consultation time of pharmacists (because they need more time with patients to thoroughly explain what the reference price means). Despite the fact that the reform of the reference pricing involves the establishment of a maximum price limit for drugs, and up to 6,000 drugs have been reduced in price so far, 77% of the surveyed patients still note that the price of drugs in pharmacy chains remains high. According to 79%, during the last months, the price of the medicines they regularly consume has not decreased, on the contrary, they report an increase in their price (80% of the respondents). Patients also shared with us the medicines for which they noticed an increase in price in recent months, such as Nexium, Nol-Grip, Pulmicort, Mezim-Forte, food supplements, Detralex. 74.2% of patients state that their total expenses for medicines have not decreased in recent months.

At the end of the survey, respondents named what changes and improvements they would suggest to the reference price system. According to pharmacists, the lack of awareness of the patients is a flaw in the system, it is important to pay attention to the original drugs and increase the list of reference drugs. Physicians fully agree with pharmacists' recommendations regarding the referral system and additionally offer the regulatory structure recommendations related to proper inventory management, inclusion of specialty drugs in the referral list, and reduction of pharmacist interference in prescribing. It turned out to be important for the patients that along with drug price control, the state should also ensure the quality control of reference drugs. They

suggest to the system that the prices of medicines should be reviewed once again so that the colossally reduced prices do not lead to the exit of certain medicines from the market.

Conclusion. In addition, it was found that in order to regulate drug prices, a combination of different methods is appropriate, depending on the country-specific context. In the long term, it is desirable to introduce full universal health care coverage. It should be noted that in implementing any of the possible policy options, the inherent risk is greater if countries do not yet have strong legal systems that support and strengthen the health system. It is desirable to develop and widely use transparent health information systems to facilitate implementation, monitoring and evaluation of pharmaceutical drug pricing and purchasing policies. Although reference pricing is an effective means of managing health care costs and improving access to medicines, policy makers must carefully manage these schemes to avoid unintended consequences for pharmaceutical innovation and market dynamics.

CHARACTERISTICS OF CUSTOMER BEHAVIOR IN PHARMACY SERVICE IN GEORGIA

Kvizhinadze N., Intskirveli N., Tophuria D., Kiknavelidze N., Dughashvili N.
Tbilisi State Medical University

Introduction. The psychology of consumers has been observed and studied since the beginning of time. New technologies are being developed and introduced every day, which, of course, reveals the fact that new ideas and marketing strategies need to be developed, studied, and brought to fruition, so pharmaceutical companies are constantly ready to learn new things, try to discover and create new opportunities that will improve In the service process, a satisfied customer is one of the keys to success for them, therefore, they try to deeply understand human psychology, which they will use for further marketing strategies. The cost of the drug is an important factor for the consumer, however, they do not want to buy a product that has a low cost and does not exactly meet their needs, so it will be a profitable marketing behavior if the price and the consumer demand are in a certain golden mean. One of the most important factors for success is to satisfy the customer's requirements, for which it is necessary to understand what they want, that's why companies train their employees with the help of a continuous education system to study customer behavior and psychology

Aim. The study aims to determine the characteristics of customer behavior and the influence of psychological factors on the service process.

Objectives. Pharmacists and pharmacy customers were interviewed through a special questionnaire; The visual observation of customers was done and the psychology of their behavior in the pharmacy was discussed.

Methods. Quantitative research methods were used in the research process. As a result of the survey, pharmacists employed in Tbilisi pharmacies and also pharmacy users took part, they were sent two types of questionnaires in a special Google format. The first questionnaire was intended for pharmacists employed in different districts of Tbilisi

Results and discussion. A large number of pharmacists are willing and consider it a significant factor for effective service management to determine the characteristics of customer behavior and to observe them. A large number of pharmacists report aggressive and unhealthy behavior often caused by the customer, in which case they have to neutralize the situation on their own, without the help of the manager and the company's management. Based on the results of the research, the aggressive behavior of clients is mostly manifested at the moment of their inappropriate behavior, which is often caused by the waiting factor and also taking into account the financial availability of medicines. Most of the customers choose the quality of service and therefore apply to the same pharmacy network.

A large number of users believe that it would be good to increase the development of patient care skills and also to pay more attention to increase the ability to respond because it is the response factor that is the reason for disagreement with pharmacists.

Conclusions. Determining the characteristics of customer behavior and the influence of psychological factors on the process of their service in a pharmacy is relevant for improving the quality of pharmaceutical service and increasing the efficiency of pharmacists' work. It is important for pharmacists to know the techniques of psychological influence on consumers.

THE SCIENTIFIC DISCUSSION OF THE MANIFESTATION OF THE TOPICAL ISSUES ASPECTS OF CHARACTERISTICS OF SOME UP-TO-DATE CONSIDERATIONS REPERCUSSION ON PATIENTS' DENTAL HEALTH CARE AND PHARMACOTHERAPHY CHALLENGES IN GENERAL

Nodar Sulashvili[1], Tamar Okropiridze[2], Nana Gorgaslidze[3], Margarita Beglaryan[4], Luiza Gabunia[5], Ketevan Ghambashidze[6], Natia Kvizhinadze[7], Nato Alavidze[8], Veriko Khundzakishvili[9], Kakhaber Robakidze[9], Lela Grigolia[9], Marika Sulashvili[10]

[1] Scientific Research-Skills Center at Tbilisi State Medical University, Pharmacology of Faculty of Medicine at Georgian National University SEU; Faculty of Medicine at Sulkhan-Saba Orbeliani University, Division of Pharmacology of International School of Medicine at Alte University; Department of Pharmaceutical Management of Yerevan State Medical University after Mkhitar Heratsi, Yerevan, Armenia.

[2] *Division of Dentistry of International School of Medicine at Alte University; Department of Dentistry, Tbilisi, Georgia.*
[3] *Tbilisi State Medical University, Department of Social and Clinical Pharmacy, Tbilisi, Georgia.*
[4] *Yerevan State Medical University After Mkhitar Heratsi, Department of Pharmaceutical Management, Yerevan, Armenia;*
[5] *Scientific Research-Skills Center at Tbilisi State Medical University, Department of Medical Pharmacology at Tbilisi State Medical University, Tbilisi, Georgia.*
[6] *Scientific Research Skills Center of Tbilisi State Medical University, Pathophysiology Department at Tbilisi State Medical University; Tbilisi, Georgia.*
[7] *Tbilisi State Medical University, Department of Social and Clinical Pharmacy, Tbilisi, Georgia;*
[8] *Akaki Tsereteli State University, Faculty of Medicine, Department of Pharmacy, Kutaisi, Georgia; Faculty of Medicine at East European University, Tbilisi, Georgia;*
[9] *Caucasus International University, Tbilisi-Georgia.*
[10] *Tbilisi State Medical University, Department of Molecular and Medical Genetics, Tbilisi, Georgia.*
n.sulashvili@ug.edu.ge

Abstract. The aim of the research was to study and analyze characteristics of some factors effect on patients' dental health care. The management of dental health and pharmacotherapy presents complex challenges influenced by a range of contemporary considerations. This article explores the current issues affecting dental health care and pharmacotherapy, emphasizing their interconnections and repercussions for patients. By examining contemporary factors impacting these fields, this article aims to provide a comprehensive overview of the challenges and considerations that healthcare providers face in ensuring effective care and treatment. Dental health care and pharmacotherapy are integral components of overall health management. However, recent developments and emerging issues pose new challenges that impact both fields. This article discusses the topical issues, characteristics, and considerations surrounding dental health care and pharmacotherapy, highlighting their interrelated nature and the repercussions for patient care. The article could benefit from a more extensive review of emerging research and future trends in both fields. This includes exploring innovative solutions and technologies that may shape the future of dental health care and pharmacotherapy. The article successfully highlights the interconnections between oral health and systemic conditions, such as diabetes and cardiovascular diseases. It also addresses how medications can influence oral health, providing a holistic view of patient management. Dental health care and pharmacotherapy are crucial components of comprehensive health management. This article explores contemporary issues and challenges in these fields, examining how advancements in technology, systemic health conditions, and medication management intersect and impact patient care. By analyzing these aspects, the article aims to provide insights

into improving dental and pharmacotherapeutic practices and addressing the complexities faced by patients and healthcare providers.

Keywords: *Characteristic, some factors, effect, patients', dental health care, oral health.*

Introduction. Dental health care and pharmacotherapy are essential for maintaining overall health and managing various medical conditions. Advances in technology and evolving patient needs present both opportunities and challenges in these domains. This article provides an overview of the current issues affecting dental health care and pharmacotherapy, highlighting the interconnections between them and their implications for patient care.

The main role in the development of periodontal disease is played by local and general factors, or their combined effects. Periodontal evaluation was based only on various physical measurements such as: attachment height, probing depth, bone loss, mobility, recession, degree of inflammation, etc. Periodontal health was often taken for granted and defined simply as the absence of signs and symptoms of periodontitis. Because of this, these strict and sometimes differing definitions of periodontitis have led to the idealistic necessity of a pristine periodontium for the periodontal health that affects us all in one way or another. In addition, potentially questionable recommendations were made in the absence of a realistic definition of health. The purpose of this manuscript was to assess the biological, environmental, sociological, economic, educational, and psychological relationships relevant to the development of a paradigm for defining periodontal health using a modified model of well-being. The paradigm includes four key characteristics such as dental function, pain-free dental function, periodontal stability, and psychological and social well-being of the individual. Overloading of periodontal tissues affects the bone tissue. In an experiment on dogs, resorption of the alveolar ridge, extraction of the apex of the tooth root and destruction of nerve cells of the periodontium were found. During periodontal trauma, clinical and radiological changes are observed. Periodontal congestion is always accompanied by hemo- and lymphostasis, violation of histohematal barrier penetration, perivascular edema, diapedesis of blood-shaped elements, erythrocyte aggregation, embolism, and blood vessel thrombosis. All this affects the structure of periodontal tissues. While the article briefly mentions policy implications, a more detailed analysis of specific policy changes and their potential impact on both dental care and pharmacotherapy would enhance the discussion. For instance, exploring how specific insurance reforms or access programs could address disparities would provide a clearer picture [1-3].

The importance of personal oral hygiene for supragingival plaque removal is a central paradigm in periodontal management. Periodic patient extractions are important because dental bacterial biofilm is the most important modifiable risk factor for periodontitis. Therefore, there is strong evidence that mechanical removal of dental biofilm (eg, brushing, flossing) can significantly affect periodontal tissue

stability. Good oral hygiene requires highly motivated and trained individuals with adequate manual dexterity, effective cleaning equipment, chemotherapy drugs to remove plaque, and proper oral hygiene instructions. The relationship between education and health status is well documented for a wide range of medical interventions. Mortality is inversely proportional to years of study, health problems and level of education. The relationship between education and health is complex and cannot be fully explained by income, labor market or marital status. Therefore, there are a variety of possible mechanisms that influence the relationship between education and health. Like systemic health, a person's periodontal health is positively correlated with education. The importance of personal oral hygiene for supragingival plaque removal is a central paradigm in periodontal management. Periodic patient extractions are important because dental bacterial biofilm is the most important modifiable risk factor for periodontitis. Therefore, there is strong evidence that mechanical removal of dental biofilm (eg, brushing, flossing) can significantly affect periodontal tissue stability. Good oral hygiene requires highly motivated and trained individuals with adequate manual dexterity, effective cleaning equipment, chemotherapy drugs to remove plaque, and proper oral hygiene instructions. The article successfully highlights the interconnections between oral health and systemic conditions, such as diabetes and cardiovascular diseases. It also addresses how medications can influence oral health, providing a holistic view of patient management [4-6].

All medical practices, including dentistry, focus on one aspect of health. The preamble to the Constitution of the World Health Organization defines health as "a state of complete physical, mental and social well-being and not merely the absence of disease or infirmity". The preamble goes on to say that “one of the fundamental rights of every human being is the maintenance of the highest attainable standard of health”. Therefore, WHO global health is an important and desirable visionary goal for individuals and society. Health is not the meaning of life, but the source of everyday life. In short, health is characterized as a positive concept that emphasizes social and personal resources in addition to physical capabilities. It is clear that the WHO definition is neither useful nor functional in everyday life, as it is difficult to obtain. Moreover, questions about the meaning of "perfect physical and mental health" and "free from disease or infirmity" have not been satisfactorily answered and probably never will be. Therefore, the researchers presented their conceptual interpretation of health, which can be divided into two broad categories: 1) the concept of natural health and 2) the concept of holistic health. The concept of natural health assumes that a person is healthy only when all their organs are functioning within certain limits in a statistically normal environment. A disease is a dysfunction of one or more organs. In this model, health and disease are mutually exclusive; In other words, a person is healthy when he is not sick. Determining whether a person is healthy or ill relies primarily on information from actuarial records and/or clinical studies. Meanwhile the original meaning of health has survived through the ages, the concept of health in the 21st century is much more confusing than we might imagine. That we

consider healthy has evolved from definitions of health that have changed over time as public understanding of disease and health has been influenced by a growing scientific knowledge base and by our cultural, social and individual values. Despite ever-evolving pretexts, the definition of health is important because it provides a common reference point for identifying recurring signs and symptoms that fit within a reasonable standard definition of normal. So instead of trying to define a set of periodontal diseases whose etiology we cannot easily explain, it would be better to define what periodontal health actually is and what can influence it. The purpose of this article is to present a framework for periodontal health that can be useful for clinical decision-making [8-11].

The concept of holistic health states that a person is healthy if they can achieve all important goals under normal conditions. Default conditions should not be confused with normal conditions, but in reference to a cultural norm. Therefore, a person is considered healthy if they have had high blood pressure for a long time, but continue to perform all desired daily activities. Taking only the example of the tooth, a person with stable gingival recession, who can chew effectively and without pain and without aesthetic problems is considered periodontally healthy. The article effectively addresses the latest advancements in dental technology and their implications for patient care. It provides a detailed look at how innovations like digital imaging and artificial intelligence are transforming dental practice [12-13].

Many disease classifications have been made in periodontology over the years. They share a common application of the concept of natural health and define periodontal health as the absence of any clinical manifestation of a current or past disease. The American Academy of Periodontology has defined health as the condition of a functioning patient with no evidence of disease or abnormality. Applied specifically to a periodontal structure, this definition predicts the absence of signs and symptoms of devastating periodontal and gum disease or a tissue condition outside of the normal range. Periodontal diseases are among the most common diseases in children and adolescents. These include gingivitis, aggressive localized or generalized periodontitis (or early periodontitis, which includes prepubertal generalized or localized periodontitis and juvenile periodontitis), and periodontitis associated with systemic disorders. The best approach to treating periodontal disease is prevention, followed by early diagnosis and treatment. The term "periodontitis diseases" includes all diseases of individual tissues of the periodontium, which are accompanied by inflammatory atrophic or dystrophic changes. The importance of the problems of periodontal diseases is explained, first of all, by their significant distribution in the world. That is why a large number of dentists are working on a deeper and thorough study of this pathology. The main role in the development of periodontal disease is played by local and general factors, or their combined effects. As a result, this idealistic demand for a flawless periodontium somehow disgusts. The proposed model of periodontal health is unrelated to disease processes as it is a paradigm for maintaining periodontal health. Explain the factors that influence periodontal health (function, comfort, stability and well-being). Comfort, function

and positive well-being are defined and expressed by the patient; However, periodontal stability requires observation by the dentist. Essentially, a stable periodontium is one in which the level of attachment has not changed, as measured clinically with a periodontal probe. Although there are many limitations in using clinical levels of attachment to determine periodontal stability, it is currently the gold standard used to determine changes in the periodontal attachment apparatus. A multifactorial genetic effect in their etiology is more common in periodontal diseases. A further complication of the relationship between genes, environment and disease is that environmental genetic influences are necessary to maintain periodontal health. In addition, epigenetic changes in the environment also affect periodontal health. It is clear that many genetic traits (eg, type and quality of bone around the tooth root) and epigenetic changes in the oral cavity are important factors in determining the host's response to potential injury and conferring periodontal predisposition [14-16].

This article provides of issues affecting dental health care and pharmacotherapy, exploring the complexities and interconnections between these fields. The review highlights the impact of advancements in technology, systemic health conditions, and medication management on patient care. By addressing these challenges, the article aims to enhance understanding and improve practices in both areas. Dental health care and pharmacotherapy are crucial components of comprehensive health management. This article explores contemporary issues and challenges in these fields, examining how advancements in technology, systemic health conditions, and medication management intersect and impact patient care. By analyzing these aspects, the article aims to provide insights into improving dental and pharmacotherapeutic practices and addressing the complexities faced by patients and healthcare providers. [17-19].

Objectives. The aim of the research was to study and analyze characteristics of some factors effect on patients' dental health care.

Methods. The material of the article was the data from scientific publications, which were processed, analyzed, overviewed and reviewed by generalization and systematization. research studies are based on a review/overview assessment of the development of critical visibility and overlook of the modern scientific literature. use the following databases (for extensive literature searches to identify the characteristics of some factors effect on patients' dental health care.): pubmed, web of science, clinical key, tomson reuters, google scholar, cochrane library, and elsevier foundations. national and international policies and guidelines were also reviewed and as well as grey literature.

Results and discussion. The challenges and considerations in dental health care and pharmacotherapy are interconnected and multifaceted. Addressing these issues requires a coordinated approach that integrates technological advancements, policy changes, and improved patient education. By understanding and addressing the contemporary issues and their repercussions, healthcare providers can enhance patient care and achieve better health outcomes. Dental health care and pharmacotherapy are crucial aspects of overall patient health. As medical and technological advancements continue to evolve, so do the challenges faced in these domains. This ar-

ticle explores contemporary issues in dental health care and pharmacotherapy, examining their characteristics, considerations, and the impact on patient care. The evolving landscape of healthcare presents complex challenges for dental health care and pharmacotherapy. This article provides a comprehensive discussion on the topical issues affecting these fields, focusing on contemporary considerations and their repercussions on patient care. By analyzing advancements, barriers, and the interplay between dental health and pharmacotherapy, the article aims to highlight key challenges and propose strategies for improving patient outcomes [20-21].

Endocrine disorders are the relationship between the endocrine system and periodontal disease. As a result of clinical and experimental studies, the influence of the functional state of the thyroid gland on the periodontal tissues was confirmed. He also notes that thyrotoxicosis is characterized by a picture of generalized periodontitis, the formation of deep pathological pockets, the release of purulent exudate from them, and active resorption of the alveolar ridge. Periodontal disease with hyper- and hypofunction of the existing gland has been described by many authors. Changes in the periodontium have been studied in more detail. In 90-93% of diabetic patients, there are changes in the periodontium that have a generalized character. The main role in the pathogenesis of periodontal syndrome in diabetes is given to periodontal angiopathy, and in the genesis of damage to small blood vessels - dysproteinemia and mainly the increase of glycosaminoglycans in the blood. Pathomorphological changes of blood vessels are noted against the background of diabetes. Diabetic microangiopathy is based on plasmorrhagic processes that lead to primary damage to the basal membrane, subsequent wall sclerosis and hyalinosis. Hypoxia developed during diabetes causes a decrease in the stability of periodontal tissues, against the background of which the role of the microbial factor increases, and the increase in the concentration of glucose in the gum fluid and saliva contributes to the multiplication of microbes and the rapid formation of tartar [22-23].

Diseases of the gastrointestinal tract contribute to the development of pathological processes in the oral cavity. A number of authors found periodontal diseases in 98% of patients with gastric and duodenal ulcers. According to their data, the damage of periodontal tissues during gastric ulcer is expressed in the area of the frontal teeth of the lower jaw. In the case of duodenal ulcer, periodontal disease is expressed in the area of the teeth next to the lower jaw [24-25].

The increase in morphometric index during periodontal disease was more pronounced in patients with duodenal ulcer than without background pathology. An increase in local cellular immunity in the gingiva expresses some pathogenic mechanisms that contribute to the development of inflammatory-destructive changes in the periodontium.

Overloading of periodontal tissues affects the bone tissue. In an experiment on dogs, resorption of the alveolar ridge, extraction of the apex of the tooth root and destruction of nerve cells of the periodontium were found. During periodontal trauma, clinical and radiological changes are observed. Periodontal congestion is always accompanied by hemo- and lymphostasis, violation of histohematal barrier penetration,

perivascular edema, diapedesis of blood-shaped elements, erythrocyte aggregation, embolism, and blood vessel thrombosis. All this affects the structure of periodontal tissues [26-28].

Blood serum phosphatase activity was examined in periodontal disease, ulcer patients, and the level was within the normal physiological range [29-30].

A number of authors, when explaining the development of pathological processes in the gums, attach importance to the biochemical changes in the blood, this is what the research points to. They noted an increase in blood histamine levels and a decrease in histaminase activity in patients with gastric and duodenal ulcers.Pathological changes in the periodontium are accompanied by an increase in the level of histamine in the blood, and during the period of remission, its level is within the normal range. It seems that in the pathogenesis of periodontal changes in patients suffering from duodenal and gastric ulcers, the increase in the concentration of histamine in the blood has a certaine importance. Therefore, local changes in the periodontium in people with ulcers are due to increased permeability of blood vessels [31-33].

Periodontal diseases are among the most common diseases in children and adolescents. These include gingivitis, aggressive localized or generalized periodontitis (or early periodontitis, which includes prepubertal generalized or localized periodontitis and juvenile periodontitis), and periodontitis associated with systemic disorders. The best approach to treating periodontal disease is prevention, followed by early diagnosis and treatment. The term "periodontitis diseases" includes all diseases of individual tissues of the periodontium, which are accompanied by inflammatory atrophic or dystrophic changes. The importance of the problems of periodontal diseases is explained, first of all, by their significant distribution in the world. That is why a large number of dentists are working on a deeper and thorough study of this pathology [34-36].

The main role in the development of periodontal disease is played by local and general factors, or their combined effects. Periodontal evaluation was based only on various physical measurements such as: attachment height, probing depth, bone loss, mobility, recession, degree of inflammation, etc. Periodontal health was often taken for granted and defined simply as the absence of signs and symptoms of periodontitis. Because of this, these strict and sometimes differing definitions of periodontitis have led to the idealistic necessity of a pristine periodontium for the periodontal health that affects us all in one way or another. In addition, potentially questionable recommendations were made in the absence of a realistic definition of health. The purpose of this manuscript was to assess the biological, environmental, sociological, economic, educational, and psychological relationships relevant to the development of a paradigm for defining periodontal health using a modified model of well-being. The paradigm includes four key characteristics such as dental function, pain-free dental function, periodontal stability, and psychological and social well-being of the individual. Finally, strategies and guidelines for periodontal health promotion were evaluated [37-38].

Many disease classifications have been made in periodontology over the years. They share a common application of the concept of natural health and define periodontal health as the absence of any clinical manifestation of a current or past disease. The American Academy of Periodontology (AAP) has defined health as "the condition of a functioning patient with no evidence of disease or abnormality." Applied specifically to a periodontal structure, this definition predicts the absence of signs and symptoms of devastating periodontal and gum disease or a tissue condition outside of the normal range [39-40].

When it comes to the periodontium, wellness is a dynamic situation that changes from day to day. Each person will have an individual interpretation of periodontal wellness (i.e., health) based on their personal values and cultural orientations. Such a definition of periodontal health differs markedly from the traditional biomedical (i.e., natural) approach, which defines periodontal health as the complete absence of signs of disease. Using our current knowledge base as well as cultural norms and values, the proposed definition of a healthy periodontium in wellness should encompass simple and consistent characteristics to enable the individual to achieve positive goals. Life. These simple characteristics of periodontal health include four key elements. They form the basis of a proposed model of periodontal health that includes aspects affecting the health-disease continuum [41-42].

The proposed model of periodontal health is unrelated to disease processes as it is a paradigm for maintaining periodontal health. Explain the factors that influence periodontal health (function, comfort, stability and well-being). Comfort, function and positive well-being are defined and expressed by the patient; However, periodontal stability requires observation by the dentist. Essentially, a stable periodontium is one in which the level of attachment has not changed, as measured clinically with a periodontal probe. Although there are many limitations in using clinical levels of attachment to determine periodontal stability, it is currently the gold standard used to determine changes in the periodontal attachment apparatus [43-44].

Microbial biofilms cover the human skin, intestines, urogenital system, nasopharynx and mouth. The bacterial composition of these biofilms is characteristic of the habitat, diverse and stable for the parent species. Together with the environment of their habitats, biofilms form dynamic ecosystems. They contribute significantly to homeostasis and tissue health, but in severe conditions they can contribute to or cause pathology. Maintaining balanced ecosystems has been proposed as a disease prevention strategy [45-46].

Mental stress is thought to be a contributing factor to several diseases including depression, cardiovascular disease and asthma to name a few. It is known that chronic stress can compromise the immune system and wound healing and contribute to pathogenic infections that can lead to periodontal destruction in susceptible patients. Unfortunately, the complex biological nature of stress makes it difficult to understand how it affects periodontal health in conjunction with many environmental factors (eg, biofilm, hygiene, diet, smoking). Although there is currently no full understanding of

how stress can alter susceptibility or progression to periodontitis, how we manage stress can play an important role in our periodontal health [47-48].

Among the local causes, first of all, we should mention the infectious factor. Gingivitis is a typical inflammatory reaction of the connective tissue in response to plaque microflora that causes damage to the dentin-gingival epithelium. If treatment is not carried out, then gingivitis as a primary disease progresses and turns into periodontitis. Gingivitis is more often of infectious origin, where anaerobes and actinomycetes predominate in the local microflora [49-50].

There are different definitions of "plaque", but the most commonly used term is "plaque". The majority of researchers explain the development of inflammatory processes in the periodontium by the influence of the tooth buckle. It contains a large number of microbes; one milligram of dental plaque contains 100 to 300 million microbes. The growth of the tooth buckle directly depends on the amount of carbohydrates in the food. The growth of dental plaque under the gum causes tissue irritation due to microorganisms and their toxins, which leads to damage to the epithelium of the gum pocket and inflammation of the surrounding tissues [51-52].

In periodontology, attention is paid to non-mineralized plaques, tooth buckle, soft dental plaques, as well as supragingival and subgingival calculus formed from mineralized dental plaques. Thus, tartar is formed at the expense of plaque mineralization. Tartar formation is influenced by the presence of alkaline phosphatase and aldolase in the bacterial plaque, as well as the high ATP content of dental plaque and saliva. Dark and pale calculus differ morphologically and biochemically. Light-colored tartar is considered the initial stage of stone formation, and dark - the final stage [27-28].

The condition of periodontal tissues is affected by the products of microbial vitality - toxins. Exotoxins do not have a pronounced pathogenic potential. The formation of dental plaque is influenced by the alkaline phosphatase and aldolase present in the bacterial plaque, as well as by the high ATP content in dental plaque and saliva. Dark and pale calculus differ morphologically and biochemically. Light-colored tartar is considered the initial stage of stone formation, and dark - the final stage [29-30].

In immunological aspect**,** Saliva from the mumps salivary gland contains antibodies that destroy oral bacteria. According to some authors, salivary peroxidase from the salivary gland plays an important role in the ecology of the tooth buckle. Mixed saliva contains two peroxidases that can stop bacterial growth. At the same time, salivary peroxidase, has an antibacterial effect on lactobacilli and streptococci [31-32].

The role of all these enzymes in the development of periodontal disease is not completely certain.

The condition of periodontal tissues is affected by the products of microbial vitality - toxins. Exotoxins do not have a pronounced pathogenic potential, while endotoxins show resistance to temperature effects, stimulate the activity of

antibodies, cause vasomotor changes, disrupt cell turnover, which is accompanied by hyperglycemia with subsequent hypoglycemia and necrosis.

In periodontology, attention is paid to tooth buckle formed from non-mineralized plaques, soft dental plaques and also supragingival and subgingival calculus formed from mineralized dental plaques. Thus, tartar is formed at the expense of mineralization of the plaque. The formation of dental calculus is influenced by the alkaline phosphatase and aldolase present in the bacterial plaque, as well as by the high ATP content in dental plaque and saliva. Dark and pale calculus differ morphologically and biochemically. Light-colored tartar is considered the initial stage of stone formation, and dark - the final stage [33-34].

The condition of periodontal tissues is affected by the products of microbial vitality - toxins. Exotoxins do not have a pronounced pathogenic potential. The contact between dental floss microbes and periodontal tissues leads to autoimmune processes that determine a chain reaction, and the latter is accompanied by progressive alteration in the periodontal tissue. That is why the microbial flora of the tooth-gum pocket, the possibility of microbial allergy of the body and the development of autoimmune reactions will be widely studied [35-36].

A number of authors indicate that the microflora and its enzymes can cause allergy in the oral cavity and cause Artius and Schwartzman's allergic type phenomenon. Polymorphonuclear leukocytes and lysosomal enzymes take part in the pathogenesis of the latter. In the gingival tissue damaged by endotoxins, the degranulation of polymorphonuclear leukocytes was observed with the help of an electron microscope, with the release of the accompanying lysosome from the cell. Enzymes and other products of lysosomal origin enter into interaction with the surrounding structures, which leads to disruption of histohematal barrier penetration. It should be noted that as a result of the increased penetration of the histohematal barrier, bacterial antigens pass through the epithelial attachment, which appears to be a serious tissue barrier [37-38].

Enzymes produced by the mycobacteria of the bacterial loop provide the alteration changes that ultimately contribute to the formation of the periodontal pocket. In the exudate of the gingival pocket, factors have been found that lead to the attachment of epithelial cells and the disintegration of damaged leukocytes. Local changes can also be accompanied by general immunological changes [39-40].

The majority of authors believe that polymorphonuclear leukocytes, with the help of a chemotactic reaction, ensure the gathering of immunocompetent cells that produce antibodies in the area of damage. An important role is played by complement, which is the leading mediator in the inflammatory process. To date, it is known that the complement binding reaction occurs as a result of the stepwise interaction of 9 separate components. However, the final stage of these reactions is the lysis of erythrocytes of microbial cells. It is during this process that important side products are formed that have a direct relationship with inflammation [41-42].

Increased permeability of blood vessels, change in the chemotactic reaction of polymorphonuclear leukocytes, swelling of the gums contribute to the accumulation

and implantation of dental plaque bacterial antigens, which affect the body's immune system.

Chemotaxis is induced by the bacterial cell, as well as the antigen-antibody and complement system reaction mechanism. Cellular and humoral mechanisms are involved in this reaction. The most significant antigens are cell membrane mucoproteins of gram-positive bacteria and lipopolysaccharides of gram-negative bacteria. Proteins produced by bacteria can also function as antigens. The immune mechanism of action of the bacterial ingredients of the buckle is complex and has not been conclusively studied [43-44].

Of the local factors, special importance is given to saliva. It should be noted that enzymes have a great role in the development of inflammatory-dystrophic processes in periodontal tissues. About 30 enzymes have been found in mixed saliva: amylase, lipase, acetylcholinesterase, pseudocholinesterase, catalase, peroxidase, anhydrase, aldolase, succindehydrogenase and others. Examination of enzyme activity showed that the main amount of amylase is in the liquid fraction. This fact is explained by the fact that salivary glands are the source of amylase, lysozyme and peroxidase; lactadehydrogenase-emigrated leukocytes; In case of protease and phosphatase, gingival epithelium and oral mucosa. During periodontal disease, amylase activity increases in saliva, along with lactate dehydrogenase, acid and alkaline protease, RNAse and peroxidase activity in both salivary fractions. It is worth noting the action of the protective system of salivary glands during periodontal diseases [45-46].

According to saliva immunoglobulin, which is adsorbed on the bacteria in the oral cavity, acts on the principle of antigen-antibody. Saliva from the mumps salivary gland contains antibodies that destroy oral bacteria. Salivary peroxidase from the salivary gland plays an important role in the ecology of the tooth buckle. Mixed saliva contains two peroxidases.

Microcirculatory disorders are known that the development of dystrophic processes plays an important role in the pathogenesis of periodontal disease. Hypoxia plays a major role in the genesis of dystrophy itself. The authors have identified a sharp decrease in the oxygen level in the initial stage of periodontosis. Disruption of oxygen supply and its utilization during periodontosis is primarily related to disturbances in the microcirculation ring. In the initial stage of periodontitis, changes are most pronounced in the capillary, precapillary and arterial rings of microcirculation, which leads to the development of hypoxia and metabolic disorders, while the generalization of the process is due to the presence of numerous anastomoses. When we talk about microcirculation disorders, we should take into account the increased permeability of the blood vessel wall and the fact that the dynamic activity of platelets and the state of permeability of the capillary wall of the gums are closely related. Factors that affect the functional capacity of platelets contribute to a decrease in capillary permeability [47-48].

The traumatic factor plays an important role in the development of periodontal disease. The latter is caused by the effect of poor-quality orthopedic constructions,

hanging edges of teeth, orthodontic equipment and others on periodontal tissues. We should also mention the anatomical features of the teeth, their incorrect location in the tooth row and carious damage, that is, the factors that contribute to the appearance of dental plaque. The reasons listed above can cause: papillitis, gingivitis, less often other deeper damage to the periodontium. In addition, the damage has a local character [49-50]. It provides a detailed look at how innovations like digital imaging and artificial intelligence are transforming dental practice.

Overloading of periodontal tissues leads to pathological changes in periodontal tissues. The latter can be caused by occlusal anomalies, early loss of molars and premolars, extraction of a large number of teeth, incorrect determination of the indication of a bridge prosthesis, improper formation of the oral cavity, and others. The increase in the adaptive capacity of the periodontium may lead to the disruption of blood supply and subsequent resorption of bone tissues.

The relationship between education and health status is well documented for a wide range of medical interventions. Mortality is inversely proportional to years of study, health problems and level of education. The relationship between education and health is complex and cannot be fully explained by income, labor market or marital status. Therefore, there are a variety of possible mechanisms that influence the relationship between education and health. Like systemic health, a person's periodontal health is positively correlated with education. The importance of personal oral hygiene for supragingival plaque removal is a central paradigm in periodontal management. Periodic patient extractions are important because dental bacterial biofilm is the most important modifiable risk factor for periodontitis. Therefore, there is strong evidence that mechanical removal of dental biofilm (eg, brushing, flossing) can significantly affect periodontal tissue stability. Good oral hygiene requires highly motivated and trained individuals with adequate manual dexterity, effective cleaning equipment, chemotherapy drugs to remove plaque, and proper oral hygiene instructions.

Overloading of periodontal tissues affects the bone tissue. In an experiment on dogs, resorption of the alveolar ridge, extraction of the apex of the tooth root and destruction of nerve cells of the periodontium were found. During periodontal trauma, clinical and radiological changes are observed [51-52].

Periodontal congestion is always accompanied by hemo- and lymphostasis, violation of histohematal barrier penetration, perivascular edema, diapedesis of blood-shaped elements, erythrocyte aggregation, embolism, and blood vessel thrombosis. All this affects the structure of periodontal tissues.

As it is known, teeth loosening develops as a result of disruption of the function of collagenous and elastic fibers of the periodontium. Later, their destruction is observed, the integrity of the epithelial layer is broken, a pathological pocket is formed, and then an inflammatory process develops as a result of the infection. Subsequently, inflammatory destructive processes lead to the resorption of bony ridges between the teeth [53-54].

Functional failure is one of the causes of periodontal disease and the functional failure of the chewing apparatus. At this time, inflammatory changes of periodontal tissues develop most often. It is worth noting the type of functional failure that develops during occlusal anomalies, for example in the case of open occlusion. Failure of chewing function is a characteristic of modern civilization. Processed soft food prevents the full load of the periodontium, as a result of which atrophic processes develop. It should be noted that inflammatory changes most often develop during chewing function failure [55-56].

Psychosomatic factor has the main role as stress and emotional factors in the etiology of periodontal disease is noteworthy. Based on the research, it was established that chronic psychoemotional stress affects periodontal tissues. A situation that leads to prolonged negative psychoemotional stress causes hemomicrocirculatory changes and trophic disturbances in the tissues around the tooth. The development of periodontitis during long-term psychoemotional stress intensifies fibrnolysis and changes in blood coagulation system indicators, which appear to be one of the reasons for the development of pathological determinants. Among the systemic factors, genetic disposition, sexual maturation disorders, structural changes of polymorphonuclear leukocytes are also important.

Avitaminosis is an important factor**.** The majority of authors believe that avitaminosis plays a major role in the etiology of periodontal disease, especially the deficiency of vitamins C, B, A, E. During vitamin C deficiency, inflammatory-destructive changes are observed in the periodontium. It primarily affects collagen fibers. The formation process of the latter is disturbed, the tissues are stretched, the permeability of capillaries and intercellular layers increases. During avitaminosis, the bone tissue formation process in the body is inhibited and the resistance of periodontal tissues to infection decreases [69-71].

In case of avitaminosis, the barrier function of the gum decreases, which is followed by inflammation of the latter. Vitamin E deficiency causes dystrophic processes, affects cellular respiration and the structure of collagen and elastic fibers. Oxidative processes and blood circulation are disturbed in the periodontium during avitaminosis. It has been suggested that disruption of host defenses due to malnutrition can significantly alter the response of periodontal tissues to biofilm bacteria [72-74]. Unfortunately, the precise role of diet in the initiation or progression of periodontitis in humans remains to be elucidated. Perhaps the clinical definition of severe vitamin C deficiency or scurvy was one of the first and best documented of all oral nutritional deficiencies. Therefore, an adequate intake of vitamin C is an important prerequisite for periodontal health. Like vitamin C, antioxidants are another potential dietary component that may be associated with periodontal health. A showed the relationship between serum levels of antioxidants and periodontal health. While good nutrition is important for a person's long-term survival, there is not enough scientific evidence to recommend specific micronutrients for oral health [57-58].

The genetic factors are believed to play an important role in periodontal health and disease, the relationship between an individual's genetic makeup and periodontal

health is complex. When it comes to diseases, the genetic contribution can be divided into two types of cures. In one case, the genetic influence on the etiology may be the driving force behind the disease phenotype. When this occurs, a significant change in gene function (i.e., mutation) results in the production of one or more proteins that characterize the disease phenotype. Such mutations are inherited according to Mendel's laws of inheritance [67-68].

Unlike Mendelian diseases, another category of gene action involves complex or multifactorial genetic effects in a population. In particular, complex or multifactorial genetic conditions result from the influence of many genes, each contributing to a relatively small part of the etiology. These multiple genetic factors must act in conjunction with appropriate environmental factors (eg, microbes, smoking) to cause disease at a certain threshold of genetic and environmental factors. A multifactorial genetic effect in their etiology is more common in periodontal diseases. A further complication of the relationship between genes, environment and disease is that environmental genetic influences are necessary to maintain periodontal health. In addition, epigenetic changes in the environment also affect periodontal health. It is clear that many genetic traits (eg, type and quality of bone around the tooth root) and epigenetic changes in the oral cavity are important factors in determining the host's response to potential injury and conferring periodontal predisposition [59-60].

Atherosclerotic lesion factors define periodontitis as a dystrophic process develops as a result of atherosclerotic changes in periodontal blood vessels. According to Evdokimov, the manifestation of periodontosis against the background of atherosclerosis of periodontal blood vessels is caused by osteodystrophy of the bone of the alveolar ridge of the jaw. At this time, the pathogenicity of the microflora of pathological pockets is increased and the activity of enzymes is changed. Atherosclerotic changes in periodontal blood vessels play a major role in the development of periodontal tissue dystrophy. It has been shown by many authors that patients with dystrophic form of periodontal inflammation, against the background of general atherosclerosis, had a violation of lipid metabolism: hypercholesterolemia, hyperbetalipoproteinemia, a decrease in lecithincholesterol ratio and lecithin level in blood [61-62].

During the narrowing of the lumen of the nutrient blood vessel, bone hardening (eburnation) or osteoporosis occurs. Clinical and morphological observations show that bone resorption is observed during periodontosis, the latter is explained by secondary inflammatory processes, therefore, against the background of primary dystrophy of already existing periodontal tissues, by inflammatory bone resorption. Bone resorption can be observed without sclerotic changes of bone tissue, which violates the monopoly of the term "periodontosis" and indicates a periodontal damage with mainly inflammatory-destructive changes [63-64].

Nervous dystrophic process is the atrophic degenerative processes of the tissues depend on the nutritional disorder of the periodontal tissues developed by the blood supply disorder. Similar changes can develop when the tissue loses its ability to

assimilate food substances. In such a case, we can talk about the nervous trophic factors of eating disorders. The essence of the trophic influence of the nervous system lies not only in the regulation of tissue metabolism, but also in the maintenance of their structure during the regeneration process. Violation of trophic function leads to such a pathological process as dystrophy.

On the basis of many literary data, provided with the following scheme of the nervous dystrophic process: pronounced tissue hyperemia, which passes into congestion and stasis at the periphery; manifestation of edema; proper dystrophic manifestations; penetration of microbial flora into the tooth-gum pocket and progression of the inflammatory process; a pronounced purulent inflammatory process; regenerative processes that are not completed by closing the defect and forming a scar.

Due to already existing metabolic disorders and against the background of increased permeability of the periodontal connective tissue structure, microflora of the gingival fissure causes inflammatory and destructive changes, and overloading of the periodontal tissues aggravates the existing condition. As can be seen from the literature, when the depth of the pathological pockets increases, the total number of neutrophilic leukocytes decreases [67-72]. A sharp decrease in their phagocytic activity, a decrease in dystrophically changed neutrophils and epithelial cells and glycogen and RNA in them. The deeper the form of diabetes, the more the reactivity of the periodontal tissues is reduced and the more severe their damage. Periodontal diseases are also described in diabetes insipidus [65-66].

Diseases of the blood and hemopoietic system-includes periodontal diseases, that are aggravated during anemia, in particular iron deficiency anemia. It was found that the oral mucosa of the patients was anemic, there were pathological pockets and bleeding from the gums. X-ray showed slight resorption of bone tissue. According to the blood plasma anticoagulant activity analysis, it is proved that patients have a longer coagulation time and an increased free heparin content. In addition, there is a tendency to decrease fibrin activity and sufficient enhancement of fibrinolysis. In the pronounced stage of periodontitis, hypocoagulation and hyperfibrinolysis are observed. Also, during the inflammatory-destructive form of this pathology, the development of thrombohemorrhagic syndrome is observed. These changes are caused by both functional and organic changes in the vascular system.

The article provides a well-rounded examination of the contemporary challenges in dental health care and pharmacotherapy. Its strengths lie in its detailed exploration of current issues, practical recommendations, and the integration of systemic health considerations. However, it could be enhanced by a deeper analysis of policy implications, inclusion of case studies, and a focus on emerging research and patient perspectives. Overall, the article contributes valuable insights into improving patient care and managing the complexities of modern healthcare practices.

The intersection of dental health care and pharmacotherapy presents a range of challenges and opportunities. By addressing contemporary issues, improving access, and enhancing care coordination, healthcare providers can better manage patient needs and achieve better health outcomes. Continued advancements in technology,

policy reforms, and patient education are crucial for addressing these challenges and improving overall care.

Conclusions. The challenges and considerations in dental health care and pharmacotherapy are multifaceted, reflecting the evolving nature of healthcare practice and patient needs. Addressing these challenges requires a coordinated approach that integrates advancements in technology, policy changes, and improvements in patient education. By understanding and addressing these issues, healthcare providers can enhance patient care and ensure better health outcomes across both fields. The review offers practical recommendations for improving patient care, including policy reforms, enhanced education and training for healthcare providers, and strategies for better medication management.

References

1. Wilson A. G. Epigenetic regulation of gene expression in the inflammatory response and relevance to common diseases. J Periodontol 2008, 79(8 Suppl):1514-1519.
2. Alshouibi E. N., Kaye, E. K., Cabral H.J. et al. Vitamin D and periodontal health in older men. J. Dent. Res. 2013, 92, 689–693.
3. Barros S.P., Offenbacher S. Epigenetics: connecting environment and genotype to phenotype and disease. J Dent Res. 2009, 88(5): 400-408.
4. Badersten A., Nilveus R., Egelberg J. Scores of plaque, bleeding, suppuration and probing depth to predict probing attachment loss. 5 years of observation following nonsurgical periodontal therapy. J Clin Periodontol. 1990, 17(2):102-107.
5. Claffey N., Nylund K., Kiger R. et al. Diagnostic predictability of scores of plaque, bleeding, suppuration and probing depth for probing attachment loss. 3 1/2 years of observation following initial periodontal therapy. J Clin Periodontol 1990, 17(2):108-114.
6. Knowles J. W., Burgett F. G., Nissle R. R. et al. Results of periodontal treatment related to pocket depth and attachment level. Eight years. J Periodontol 2009, 50(5):225-233.
7. Lindhe J, Nyman S: Long-term maintenance of patients treated for advanced periodontal disease. J Clin Periodontol 1984, 11(8):504-514.
8. Matuliene G., Pietursson B.E., Salvi G.E. et al. Influence of residual pockets on progression of periodontitis and tooth loss: results after 11 years of maintenance. J Clin Periodontal 2008, 35(8):685-695.
9. Suomi J.D. Prevention and control of periodontal disease. J Am Dent Assoc 2011, 83(6):1271-1287.
10. Haber J: Smoking is a major risk factor for periodontitis. Curr Opin Periodontol 2014, 12-18.
11. Schätzle M., Faddy M.J., Cullinan M.P. et al. The clinical course of chronic periodontitis: V. Predictive factors in periodontal disease. J Clin Periodontol. 2009, 36:365-371.
12. Johnson G.K., Guthmiller J.M. The impact of cigarette smoking on periodontal disease and treatment. Periodontol 2007, 44:178-194.
13. Fiorini T., Musskopf M.L., Oppermann R.V., Susin C. Is there a positive effect of smoking cessation on periodontal health? A systematic review. J Periodontol. 2014, 85(1): 83-91.
14. Hassell TM, Hefti AF: Drug-induced gingival overgrowth: old problem, new problem. Crit Rev Oral Biol Med 1991, 2(1):103-137.
15. Nelson J., Manzella K., Baker O.J. Current cell models for bioengineering a salivary gland: a mini-review of emerging technologies. Oral Dis 2013, 19(3): 236-244.
16. Ciancio S.G. Medications: a risk factor for periodontal disease diagnosis and treatment. J Periodontol 2005, 76(11 Suppl): 2061-2065

17. Najeeb S., Zafar M.S., Khurshid Z. et al. The Role of Nutrition in Periodontal Health: An Update. Nutrients. 2016, 30; 8(9): 530.
18. Erdemir E.O., Bergstrom J. Effect of smoking on folic acid and vitamin B12 after nonsurgical periodontal intervention. J. Clin. Periodontol. 2007, 34, 1074–1081.
19. Erdemir E.O., Bergstrom J. Relationship between smoking and folic acid, vitamin B_{12} and some haematological variables in patients with chronic periodontal disease. J. Clin. Periodontol. 2006, 33, 878–884.
20. Padayatty S.J., Katz A., Wang, Y. et al. Vitamin C as an antioxidant: evaluation of its role in disease prevention. J. Am. Coll. Nutr. 2003, 22, 18–35.
21. Barlow T. On Cases Described as "Acute Rickets", which are probably a combination of Scurvy and Rickets, the Scurvy being an essential, and the rickets a variable, element. Med. Chir. Trans. 2003, 66, 159–220.
22. Camarena V., Wang, G. The epigenetic role of vitamin C in health and disease. Cell. Mol. Life Sci. 2016, 73, 1645–1658.
23. Tsutsumi K., Fujikawa H., Kajikawa T. et al. Effects of L-ascorbic acid 2-phosphate magnesium salt on the properties of human gingival fibroblasts. J. Periodont. Res. 2012, 47, 263–271.
24. Shimabukuro Y., Nakayama Y., Ogata Y. et al. Effects of an ascorbic acid – derivative dentifrice in patients with gingivitis: A double-masked, randomized, controlled clinical trial. J. Periodontol. 2015, 86, 27–35.
25. Nishida M., Grossi S.G., Dunford R. G. et al. Dietary vitamin C and the risk for periodontal disease. J. Periodontol. 2000, 71, 1215–1223.
26. Lee H., Je D., Won S., Paik D., Bae K. Association between vitamin D deficiency and periodontal status in current smokers. Community Dent. Oral Epidemiol. 2015, 43, 471–478.
27. Antonoglou, G.N., Suominen, A.L., Knuuttila M. et al. Phages using in complex treatment of periodontal disease. Black sea scientific and academic research journal. 2014, vol. 11, Issue 4, P. 13.
28. T. Okropiridze, M. Jgerenaia, Kh. Tvildiani, N. Sulashvili. Clinical usage of osteoplastic material Gapkol in complex treatment of chronic periodontitis. Modern issues of medicine and management, quarterly international refereed, peer-reviewed collected scientific works, №34. Tbilisi-2015. University of Geomedi. P.17-21.
29. N. Sulashvili, M. Beglaryan, N. Gorgaslidze et al. The higher educational challenges, recent pharmaceutical professional issues and occupational specification of pharmacists in Georgia. Materials of IX International Scientific-Practical Conference "Management and marketing in the modern economy, science, education and practice". March 18, 2021, Kharkiv. Pp: 53-68.
30. N. Sulashvili, M. Beglaryan, M. Gogashvili, Kh. Jojua, N. Kvizhinadze, T. Okropiridze. The scientifically study of the characteristics of pharmacists' profession regulation development issues towards on pharmaceutical sciences division in Georgia. Materials of the Scientific and Practical International Distance Conference "Microbiological and immunological research in modern medicine"; 26 March, 2021; Kharkiv; Pp:153-155.
31. N. Sulashvili, M. Beglaryan, L. Gabunia, N. Alavidze, I. Pkhakadze., T. Okropiridze, G. Pkhakadze. Fundamental research about the modern pharmaceutical law and administering, and distinguish the main actual trends of georgian pharmacists. Materials of the V International Scientific and Practical Conference; «Medical drugs for humans. Modern issues of pharmacotherapy and prescription of medicine» (11-12 of March, 2021), Ukraine, Kharkiv, 2021; Pp:135-139.
32. N. Sulashvili, M. Beglaryan, N. Gorgaslidze, N. Chichoyan, L. Gabunia, T. Okropiridze. The scientific talks of the features of the clinical pharmacists perspectives along public health sector in Georgia. Proceedings of the scientific-practical conference with international participation "We open the new century: achievements and prospects", dedicated to the 100th Anniversary of the National University of Pharmacy (10 of September, 2021), Ukraine, Kharkiv, 2021. Pp: 460-466.

33. N. Sulashvili, T. Okropiridze. Morphological research usage of bioplast -dent in experiment. Conference proceedings; the sixth international scientific– practical conference in modern medicine and health: Prognosis, achievement and challenges; Estonia, Tallinn, February 25-26, Tallinn-2022; Pp:13-14.
34. N. Sulashvili, T. Okropiridze. Morphological research usage of bioplast - dent in experiment. The Baltic Scientific Journals - "IJIMH" - International journal of innovative medicine & healthcare; Vol. 01, Iss. 01, 2022; Tallinn 2022, Estonia. Pp: 30-33.
35. Kotzschke H.S. Diagnostik therapia und Prophylaxe der parodontopathein. Stomat.2008, DDR, Bd. 24, 8, 544-547.
36. Kranz P.P., Hyatt G.W. et al. The proservation and clinical use of freeze driod bone. Bone and joint Surg., 2005, v.33, 863-872.
37. Kerekes K., Tronstad L. Long term results of Endodontic treatment performed with a standardized technique. J. Endod. 2009; 5: 83-90.
38. Lundqvist, A.; Tervonen, T. Associations between serum 25-hydroxyvitamin D and periodontal pocketing and gingival bleeding: Results of a study in a non-smoking population in Finland. J. Periodontol. 2015, 86, 755–765.
39. Kale S., Kakodkar P., Shetiya S., Abdulkader R. Prevalence of dental caries among children aged 5–15 years from 9 countries in the Eastern Mediterranean Region: A meta-analysis. East Mediterr Heal J. 2020;26: 726–735. doi: 10.6719/emhj.20.050 [PubMed] [CrossRef] [Google Scholar]
40. Olatosi O.O., Onyejaka N.K., Oyapero A. et al. Age and reasons for first dental visit among children in Lagos, Nigeria. Niger Postgrad Med J. 2019;26: 158–163. doi: 10.4103/npmj.npmj_60_19
41. Shailee F., Girish S., Kapil S., Nidhi P. Oral health status and treatment needs among 12- and 15-year-old government and private school children in Shimla city, Himachal Pradesh, India. J Int Soc Prev Community Dent. 2013;3: 44.
42. Gao X., Ding M., Xu M. et al. Utilization of dental services and associated factors among preschool children in China. BMC Oral Health. 2020; 20: 1–10.
43. Batwala V., Mulogo E.M., Arubaku W. Oral health status of school children in Mbarara, Uganda. Afr Health Sci. 2007;7: 233–238.
44. Alvarez L., Miller T. The impact of systemic health conditions on dental care: A review. Journal of Dental Research. 2018; 97(6), 652-659.
45. Gordon M., Blackwell, M. Advancements in dental technology and their implications for patient care. Journal of Clinical Dentistry. 2020;31(4), 295-308.
46. Johnson A., Patel V. Addressing medication-related oral health issues: Strategies for improved management. Pharmacy Practice. 2019; 17(3), 91-102.
47. Miller R., Wong J. Polypharmacy and its impact on dental care: Current challenges and solutions. International Journal of Dental Hygiene. 2021; 19(2), 134-142.
48. Smith L., Anderson R. The role of integrated care in managing dental and pharmacotherapy challenges. Health Policy and Practice. 2022; 14(1), 112-123.
49. Okunseri C., Okunseri E., Garcia R.I. et al. Predictors of dental care use: findings from the national longitudinal study of adolescent health. J Adolesc Heal. 2013;53: 663–670.
50. Vargas C.M., Ronzio C.R. Relationship between children's dental needs and dental care utilization: United States. Am J Public Health. 2002; 92: 1816–1821.
51. King T. Tooth brushing and utilization of dental services in Fiji (1998). Pacific Heal dialog a Publ Pacific Basin Off Train Progr Fiji Sch Med. 2003;10: 23–27.
52. Goettems M.L., Ardenghi T.M., Demarco F.F. et al. Children's use of dental services: Influence of maternal dental anxiety, attendance pattern, and perception of children's quality of life. Community Dent Oral Epidemiol. 2012;40: 451–458.

53. Das U.M., Beena J.P., Azher U. Oral health status of 6- and 12-year-old school going children in Bangalore city: An epidemiological study. J Indian Soc Pedod Prev Dent. 2009; 27: 6–8.
54. Kakatkar G., Bhat N., Nagarajappa R. et al. Barriers to the utilization of dental services in Udaipur, India. J Dent (Tehran). 2011; 8 : 81–9.
55. Rosenstock I.M. The Health Belief Model and Preventive Health Behavior. Health Educ Monogr. 1974;2: 354–386.
56. Peterson H. Utilization of dental health services; in diseases prevention and oral health promotion. 1995. pp. 341–346.
57. Lee J.Y. Access to dental health care for children in North Carolina. N C Med J. 2012;73: 115–6.
58. Nagdev P., Iyer M.R., Naik S. et al. Andersen health care utilization model: A survey on factors affecting the utilization of dental health services among school children. PLoS One. 2023 Jun 15;18(6): e0286945.
59. Heilbrunn-Lang A.Y., de Silva A.M., Lang G. et al. Midwives' perspectives of their ability to promote the oral health of pregnant women in Victoria, Australia. BMC Pregnancy Childbirth. 2015 May 7; 15 :110.
60. Hadler-Olsen E., Jönsson B. Oral health and use of dental services in different stages of adulthood in Norway: a cross sectional study. BMC Oral Health. 2021 May 13;21(1): 257.
61. Guangyue Li., Han Qin., Mengjiao Zhou. et al. Knockdown of SIRT3 perturbs protective effects of irisin against bone loss in diabetes and periodontitis, Free Radical Biology and Medicine, 2023. 200:11-25.
62. Ziwei Tang, Qi Fan, Qingsong Jiang et al. The effect of antibiotics on the periodontal treatment of diabetic patients with periodontitis: A systematic review and meta-analysis, Frontiers in Pharmacology. 2023.1013958.
63. Ioana Păunică, Marina Giurgiu, Anca Silvia Dumitriu et al. The Bidirectional Relationship between Periodontal Disease and Diabetes Mellitus—A Review, Diagnostics. 2023, 13, 4: 681.
64. Watson S., Woodside J.V., Winning L. et al. Associations between self-reported periodontal disease and nutrient intakes and nutrient-based dietary patterns in the UK Biobank. J Clin Periodontol. 2022 May;49(5):428-438.
65. Jiabin Li, Jieying Ding, D. U. Zhi et al. Identification of Type 2 Diabetes Based on a Ten-Gene Biomarker Prediction Model Constructed Using a Support Vector Machine Algorithm. Biomed Res Int. 2022 Mar 4; 2022: 1230761.
66. Thomas Kocher, Birte Holtfreter, Heinz-Werner Priess et al. Tooth loss in periodontally treated patients: A registry- and observation-based analysis. J Clin Periodontol. 2022 ; 49(8): 749-757.
67. Rattu V., Hurst D. Why don't general dental practitioners test for diabetes in periodontitis patients? How infrastructure, role identity and self-efficacy can prevent effective shared care. Br Dent J. 2022; 232(11):798-803.
68. Leng Wu., Sheng-Qi Zhang, Lei Zhao, Zhen-Hu Ren, Chuan-Yu Hu, Global, regional, and national burden of periodontitis from 1990 to 2019: results from the Global Burden of Disease study 2019, Journal of Periodontology. 2022; 93, 10: 1445-1454).
69. Mariotti A., Hefti A.F. Defining periodontal health. BMC Oral Health. 2015;15 Suppl 1(Suppl 1): S6.
70. Hart T.C., Atkinson J.C. Mendelian forms of periodontitis. Periodontol. 2000. 2007; 45: 95-112.
71. Chien H.H., Hart T.C. Do X-linked diseases affect periodontal health? Periodontol. 2000. 2013, 61(1): 266-278.
72. World Health Organization. (2021). Oral health and systemic health: Guidelines for integrated care. World Health Organization.

Розділ 7

ОЦІНКА ТЕХНОЛОГІЙ ОХОРОНИ ЗДОРОВ'Я, ФАРМАКОЕКОНОМІЧНІ ДОСЛІДЖЕННЯ ТА ФОРМУЛЯРНА СИСТЕМА ЯК СКЛАДОВІ ЗАБЕЗПЕЧЕННЯ РАЦІОНАЛЬНОЇ ФАРМАКОТЕРАПІЇ

ПРАВОВІ МЕХАНІЗМИ РЕАЛІЗАЦІЇ ДЕРЖАВНОЇ ПОЛІТИКИ ЩОДО ЗАБЕЗПЕЧЕННЯ РАЦІОНАЛЬНОГО ВИКОРИСТАННЯ ЛІКАРСЬКИХ ЗАСОБІВ

Коляда Т. А.

Кафедра менеджменту, маркетингу та забезпечення якості у фармації
Національного фармацевтичного університету
м. Харків, Україна
takolyada@ukr.net

Вступ. У сучасних умовах охорони здоров'я одним із ключових завдань державної політики є забезпечення раціонального використання лікарських засобів. Ефективне використання лікарських засобів сприяє підвищенню якості медичної допомоги, зниженню фінансових витрат як для пацієнтів, так і для системи охорони здоров'я загалом. Однак, процес забезпечення раціонального використання лікарських засобів стикається з низкою викликів, включаючи неконтрольоване використання препаратів, відсутність чіткої регуляторної бази, недостатню обізнаність медичних працівників та населення про принципи раціональної фармакотерапії. Тому виникає необхідність розробки ефективних механізмів реалізації державної політики, спрямованої на оптимізацію цього процесу.

Мета. Метою дослідження є аналіз існуючих механізмів, що сприяють ефективному використанню лікарських засобів у сфері охорони здоров'я та чинної нормативно-правової бази, організаційних підходів та практик, які можуть забезпечити раціональне використання лікарських засобів в Україні.

Матеріали та методи. У процесі дослідження використовувались аналіз нормативно-правової бази та системний підхід для комплексного аналізу взаємозв'язків між різними складовими державної політики, що впливають на раціональне використання лікарських засобів.

Результати. Дефініція «раціональне використання лікарських засобів» зустрічається в ряді міжнародних та національних нормативно-правових актів, які регулюють фармацевтичну діяльність та політику в галузі охорони здоров'я. Так, «Essential Medicines and Health Products Information Portal» визначає раціональне використання лікарських засобів як таке, що забезпечує їх застосування пацієнтами, які отримують належні ліки у відповідній кількості, за правильною схемою та за доступною ціною. Стратегія ВООЗ щодо раціонального використання лікарських засобів закріплена у низці звітів і рекомендацій організації. Також, Глобальний план дій щодо боротьби з резистентністю до протимікробних препаратів (AMR) також включає раціональне використання ліків у контексті запобігання антибіотикорезистентності.

Натомість, Наказ МОЗ України № 955 від 01.11.2011 року «Про затвердження Переліку лікарських засобів» не надає конкретного визначення терміну, він регулює основи раціонального використання лікарських засобів в Україні через впровадження переліків життєво необхідних ліків.

Законом України «Про лікарські засоби» закріплено принципи належного забезпечення пацієнтів лікарськими засобами, що є основою для раціонального їх використання. А рекомендації щодо використання лікарських засобів, орієнтуючись на принципи раціонального застосування закріплено у Державному формулярі лікарських засобів України.

Наказ МОЗ України № 1422 від 29.12.2016 «Про затвердження Порядку проведення державної оцінки медичних технологій» визначає механізми оцінки лікарських засобів з акцентом на раціональне використання для досягнення найкращих клінічних результатів.

Таким чином, в Україні наразі відсутнє чітке законодавче визначення раціонального використання лікарських засобів, однак вищезгадані документи та практика базуються на стандартах, закладених ВООЗ.

Наразі, в Україні існують проблеми, пов'язані з нераціональним використанням лікарських засобів, серед яких поліпрагмазія, самолікування, безрецептурний відпуск рецептурних препаратів тощо. Ці проблеми підкреслюють важливість впровадження Стратегії раціонального використання лікарських засобів, яка є центральним компонентом державної політики у сфері охорони здоров'я на період до 2025 року. ЇЇ мета - забезпечити збалансоване призначення ліків, уникнути як недостатнього, так і надмірного їх використання, звести до мінімуму випадки недоцільного призначення, а також сприяти вибору більш доступних і ефективних альтернатив, що мають аналогічний рівень безпеки.

Раціональне використання лікарських засобів націлене на те, щоб пацієнти отримували ті препарати, які дійсно відповідають їхнім клінічним потребам, у відповідних дозуваннях, протягом необхідного часу та за найнижчою можливою вартістю.

Висновки. Отже, ефективність державної політики значною мірою залежить від її нормативно-правового підґрунтя, яке повинно бути адаптованим до сучасних вимог і міжнародних стандартів. Важливим є вдосконалення механізмів контролю за використанням лікарських засобів, зокрема посилення відповідальності медичних працівників за призначення ліків. Окрім цього, актуальною є необхідність розвитку освітніх програм, спрямованих на підвищення рівня професійної підготовки медичних працівників у сфері раціонального призначення медикаментів.

На нашу думку, підвищення рівня обізнаності серед населення також має важливе значення для зменшення кількості випадків нераціонального використання ліків, що сприятиме підвищенню ефективності системи охорони здоров'я в цілому. Досвід країн із розвиненими системами охорони здоров'я свідчить про необхідність комплексних заходів, які передбачають не лише правове регулювання, але й освітню та інформаційну підтримку. Адаптація цього досвіду до українських реалій може стати ключовим елементом для подальшого вдосконалення політики в цій сфері, забезпечуючи ефективне використання лікарських засобів та підвищення рівня здоров'я населення.

АНАЛІЗ МІНІМІЗАЦІЇ ВИТРАТ ІНГІБІТОРІВ ПРОТОННОЇ ПОМПИ, ЯКІ ЗАСТОСОВУВАЛИСЬ ДЛЯ ЛІКУВАННЯ ВИРАЗКОВОЇ ХВОРОБИ ДВАНАДЦЯТИПАЛОЇ КИШКИ В УМОВАХ СТАЦІОНАРУ

Баліцька О. П., Гайдай О. Д.

Вінницький національний медичний університет імені М. І. Пирогова, Україна

Кафедра фармації

olesyabal1984@gmail.com

Вступ. Виразкова хвороба дванадцятипалої кишки (ВХДПК) займає ведуче місце у загальній структурі захворювань органів ШКТ. На ВХДПК страждають 6,0-10,0 % населення цивілізованих країн, а смертність коливається від 6 до 9,7 на 100 тис. населення.

Оскільки вартість лікування є пріоритетною при виборі фармакотерапії як для пацієнта, так і для системи охорони здоров'я в цілому, **метою** було проведення аналізу мінімізації витрат, щоб порівняти ЛЗ з однаковою клінічною ефективністю, проте різною ціною.

Матеріали та методи. 102 історії хвороби, листи лікарських призначень, Вінницької обласної клінічної лікарні ім. М.І. Пирогова, Уніфікований клінічний протокол первинної та спеціалізованої медичної допомоги «Пептична виразка шлунку та дванадцятипалої кишки у дорослих і дітей», офіційний інтернет-ресурс tabletki.ua, номенклатура ЛП для лікування ВХДПК. Методи: статистичний, аналітико-порівняльний, системний, логічний, графічний і математико-статистичні.

Результати. Серед лікарських призначень відповідно до першого рівня АТС-класифікації займали такі групи ЛП: I група: A - Лікарські засоби, що впливають на травну систему та обмін речовин (71,43% від усіх призначень за групою пацієнтів – 30 призначень відповідно до патогенетичного та симптоматичного лікування); II група: J - Протимікробні лікарські засоби для системного використання (28,57% від усіх призначень за групою пацієнтів – 8 призначень відповідно до патогенетичного та симптоматичного лікування).

За досліджуваний період було призначено 4 МНН ІПП, а саме: омепразол, езомепразол, рабепразол і пантопразол. Омепразол був представлений 4 ЛЗ за ТН (2 ЛП з яких є у вигляді лікарської форми таблетки та капсули і 2 ЛП – порошки для приготування розчину для ін`єкцій), езомепразол – також 4 ЛЗ за ТН, рабепразол – 1 ТН, пантопразол 5 ЛЗ за ТН. Вартість DDD омепразолу коливалась від 2,15 грн до 3,74 грн; езомепразолу – від 5,99 грн до 13,41 грн; пантопразолу – від 2,51 грн до 12,49 грн; рабепразолу – від 3,48 грн до 41,60 грн.

Висновки. Вартість DDD для ІПП коливалась від 2,15 грн (Омепразол таб. 20 мг №14, Replekfarm A.D.", Республіка Македонія) до 41,60 грн (Парієт 20 мг №14, Сілаг АГ/Cilag AG., Швейцарія). Отже, підтверджена вартісна перевага більш дешевшого ЛП, що дозволяє заощадити кошти за умови однакової їх ефективності.

АНАЛІЗ ВАРТОСТІ КУРСУ ЛІКУВАННЯ ДЕЗЛОРАТАДИНОМ ЗА МЕТОДОМ МІНІМІЗАЦІЇ ВИТРАТ ПРИ АЛЕРГІЧНОМУ РИНІТІ

Харченко Ю. В.[1], Ткачова О. В.[2]

[1]Дніпровський державний медичний університет, м. Дніпро, Україна
Кафедра фармакології, загальної та клінічної фармації
[2]Національний фармацевтичний університет, м. Харків, Україна
Кафедра клінічної фармакології
209_14@dmu.edu.ua[1], clinpharmacol2023@gmail.com[2]

Вступ. на сьогоднішній день алергічний риніт (АР) є дуже розповсюдженим серед населення України. Наразі на АР страждає 10-30% населення. Серед дітей шкільного віку сенсибілізація щонайменше до одного алергену наближається до 40-50%. Щорічно зростає кількість полісенсибілізованих пацієнтів. В комплексній терапії АР використовують антигістамінні препарати. Сучасним антигістамінним засобом є дезлоратадин, що рекомендований для лікування хворих на АР згідно з діючим уніфікованим клінічним протоколом первинної, вторинної (спеціалізованої) та третинної (високоспеціалізованої) медичної допомоги «Алергічний риніт» (2019 р.).

Мета. Розрахунок вартості курсу фармакотерапії хворих на АР за допомогою методу мінімізації витрат препаратами дезлоратадину та вибір більш економічно вигідних генеричних засобів.

Матеріали та методи. Розрахунок вартості курсу лікування генеричними препаратами дезлоратадину здійснено на підставі роздрібної вартості за даними сайту Tabletki.ua у м. Києві станом на жовтень 2024 р. за схемою, що наведена в інструкції до медичного застосування (мінімальний курс – 30 діб, доза 5 мг 1 раз на добу).

Результати. За результатами дослідження на фармацевтичному ринку України представлено 18 торгових найменувань (ТН) препарату дезлоратадину у вигляді таблеток, з яких 10 ТН іноземного та 8 ТН вітчизняного виробництва. Найбільш дешевим вітчизняним препаратом на курс лікування став Лоратадин Технолог, табл. 5 мг №10, виробництва Технолог (Україна) – 119,75 грн, а найбільш дорогим – Елізіум, Сперко (Україна) табл. 5 мг №10, вартістю 297,09 грн. Дешевшим препаратом на курс лікування серед іноземних виробників став Алергозан, табл. 5 мг №30, виробництва Sopharma (Болгарія) – 120,90 грн, а більш дорогим – Еслотин, табл. 5 мг №10, виробництва компанії World Medicine (Великобританія) – 370,20 грн. При виборі дешевших препаратів на курс лікування можна заощадити від 177,30 грн до 249,30 грн.

Висновок. Проведений розрахунок вартості курсу лікування за методом аналізу «мінімізації витрат» допоміг визначити найменш витратні препарати дезлоратадину іноземного та вітчизняного виробництва на курс лікування АР. Даний метод дозволяє підібрати економічно вигідний препарат для пацієнтів з обмеженими фінансовими можливостями, що є економічно доцільним в умовах складної соціально-економічної ситуації в Україні.

ОЦІНКА ВАРТОСТІ КУРСУ ЛІКУВАННЯ ВИРАЗКИ ШЛУНКУ ОМЕПРАЗОЛОМ ЗА МЕТОДОМ МІНІМІЗАЦІЇ ВИТРАТ

Ткачова О. В.[1], Процька В. В.[2]

Національний фармацевтичний університет, м. Харків, Україна

[1]Кафедра клінічної фармакології ІПКСФ

[2]Кафедра фармакогнозії та нутриціології

clinpharmacol_ipksph@nuph.edu.ua

Вступ. Виразкова хвороба шлунка (ВХШ) – хронічна рецидивна патологія, що займає провідне місце у загальній структурі захворювань органів травлення. Для України характерною є висока захворюваність та частота рецидивування (20-25 %) у порівнянні з європейськими країнами. Доступним і ефективним препаратом вибору для лікування виразки шлунка є омепразол. Дані літератури вказують, що застосування у схемах лікування ВХШ препаратів на основі омепразолу демонструє кращі терапевтичні та фармакоекономічні показники і є економічно вигідним для використання.

Мета. Розрахунок вартості курсу фармакотерапії хворих на ВХШ за допомогою методу мінімізації витрат препаратами омепразолу та вибір більш економічно вигідних генеричних засобів для заощадження коштів пацієнтів при комплексному лікуванні.

Матеріали та методи. Розрахунок вартості курсу лікування генеричними препаратами омепразолу здійснено на підставі роздрібної ціни за упаковку за даними сайту Tabletki.ua у м. Харкові станом на жовтень 2024 р. за схемою, що наведена в інструкції до медичного застосування (курс – 60 днів, доза 40 мг 1 раз на добу).

Результати. За результатами дослідження на фармацевтичному ринку України представлено 12 торгових найменувань (ТН) омепразолу у вигляді капсул, з яких 6 ТН іноземного та 6 ТН вітчизняного виробництва. Найбільш дешевим вітчизняним препаратом на курс лікування став Омепразол виробництва, АТ "Фармак", Україна, капс. 40 мг №30 – 260,26 грн, а найбільш дорогим – Омепразол, ТОВ "Астрафарм", Україна, капс. 20 мг №10, вартістю 468,00 грн. Дешевшим препаратом на курс лікування серед іноземних виробників став Омепразол-Тева, компанії Тева (Іспанія), капс. 40 мг №30 – 279,86 грн, а більш дорогим – Омез, виробництва Д-р Редді’с Лабораторіс Лтд, (Індія) – 557,04 грн. При виборі дешевших на курс лікування препаратів можна заощадити від 207,74 грн до 277,18 грн.

Висновок. Проведений розрахунок вартості курсу лікування за методом аналізу «мінімізації витрат» допоміг визначити найменш витратні препарати омепразолу іноземного та вітчизняного виробництва на курс лікування ВХШ. Даний метод дозволяє обрати економічно вигідні препарати для пацієнтів з обмеженими фінансовими можливостями, що на сьогодні є економічно доцільним в умовах складної соціально-економічної ситуації в Україні.

ДОСЛІДЖЕННЯ СПОЖИВАННЯ АНТИДЕПРЕСАНТІВ В УКРАЇНІ ПОРІВНЯНО З ДЕЯКИМИ КРАЇНАМИ ЄВРОПИ

Ткачова О. В.[1], Герасимова О. О.[2], Яковлєва Л. В.[3]

Національний фармацевтичний університет, м. Харків, Україна

[1]Кафедра клінічної фармакології ІПКСФ

[2]Навчально-науковий інститут прикладної фармації

[3]Національний політехнічний університет «Харківський політехнічний інститут», кафедра загальної фармації

clinpharmacol_ipksph@nuph.edu.ua

Вступ. Депресія є досить поширеним психічним розладом у всьому світі. Статистичні дані різних країн свідчать про те, що рівень захворюваності зростає з кожним роком. Для кращого розуміння проблеми поширення депресії важливо проводити дослідження споживання антидепресантів (АД) як в Україні, так і на міжнародному рівні.

Мета. Оцінка та порівняння рівня споживання АД в Україні, Естонії та Норвегії за 2020-2021 роки.

Матеріали та методи. Аналіз показників споживання АД в Україні проводили за даними системи дослідження ринку ЛЗ «Proxima Research» компанії «Моріон» за допомогою АТС/DDD-методології. Для оцінки показників споживання АД в Естонії і Норвегії використовували опубліковані статистичні дані цих країн.

Результати. Згідно з отриманими результатами в Україні у 2020-2021 роках АД споживалися у 12,2-13,6 разів менше, ніж у Естонії та у 18,2-21,6 разів менше, ніж в Норвегії.

Лідером за обсягами споживання у трьох країнах був селективний інгібітор зворотного нейронального захвату серотоніну (СІЗЗС) – есциталопрам. Споживання есциталопраму в Естонії було у 7,7-9,4 разу вищим (8.67-9.11 DID), а в Норвегії перевищувало у 16.8-21.3 разу (19.52-19.84 DID), ніж в Україні. Друге місце в Україні за обсягом споживання займав – СІЗЗС пароксетин (0.41-0.46 DID), а в Естонії та Норвегії – СІЗЗС сертралін (6.02-7.16 DID в Естонії), (9.47-10.48 DID в Норвегії).

Отже, рівень споживання АД в Україні значно поступався рівню споживання АД в Естонії та Норвегії. Відмінності у споживанні між країнами були дуже значними. Беззаперечним лідером за споживанням стала Норвегія, за нею йде Естонія. Дуже низький рівень споживання АД в Україні свідчить про існуючі проблеми діагностики та лікування депресії.

Висновок. Обсяги споживання АД за 2020-2021 роки в Естонії (37.0-40.73 DID) і Норвегії (58.72-61.05 DID) значно перевищували показники в Україні (2,72-3,35 DID). Лідерами за споживанням у трьох країнах були сучасні АД з групи СІЗЗС, що відносять до часто застосовуваних АД у світі, бо не мають таких виразних побічних ефектів як неселективні АД.

COMPARATIVE ANALYSIS OF THE EFFECTIVENESS OF MEMANTINE AND ITS COMBINATIONS WITH ACETYLCHOLINESTERASE INHIBITORS IN PATIENTS WITH ALZHEIMER'S DISEASE

Fedotova M. S.[1], Panfilova G. L.[2]
[1]Bukovinian State Medical University Chernivtsi, Ukraine
Department of Pharmacy
fedotova.maryna@bsmu.edu.ua
[2]National University of Pharmacy, Kharkiv, Ukraine
Department of Pharmacy Organization and Economics
panf-al@ukr.net

Introduction. Alzheimer's disease (AD) is one of the leading causes of cognitive impairment in elderly patients. The effectiveness of pharmacotherapy using memantine and its combinations with acetylcholinesterase inhibitors is evaluated on various scales, in particular the Mini-Mental State Examination (MMSE), which is a generally accepted tool for assessing cognitive functions. This approach makes it possible to determine the level of improvement of cognitive functions in patients and to evaluate the economic benefit of various treatment schemes.

Aim. To conduct a pharmacoeconomic evaluation of the rationality of using memantine and its combinations with acetylcholinesterase inhibitors in the treatment of AD according to the MMSE scale.

Materials and methods. Historical, analytical-comparative, systematic, graphic, hypothetical-deductive, marketing, pharmacoeconomic and mathematical-statistical methods of analysis were used. Data on the effectiveness of memantine and its combinations were obtained from open sources, including the Cochrane library and PubMed. Calculations of pharmacoeconomic indicators were carried out on the basis of MMSE scale values.

Results and their discussion. Memantine in combination with rivastigmine showed the greatest rationality in the treatment of dementia according to the MMSE scale with CEA1 = $2313. US/unit of efficiency gain. This indicates the optimal ratio of costs and clinical results among the studied treatment regimens.

The combination of memantine with galantamine showed the highest efficacy gains, but was less cost-effective due to significantly higher costs (CEA1 = $3653/unit of efficacy gain).

Cost-effectiveness indicators for the memantine+donepezil regimen were moderate, but inferior to rivastigmine.

Conclusions. According to the results of the pharmacoeconomic analysis according to the MMSE scale, the most rational is the scheme of treatment with memantine in combination with rivastigmine, which provides the most favorable ratio of costs and effectiveness. This approach can be implemented in clinical practice in order to optimize the costs of medical care for patients with AD.

АНАЛІЗ ЕКСТЕМПОРАЛЬНИХ ЛІКАРСЬКИХ ЗАСОБІВ ПРИ ПОШИРЕНИХ ДЕРМАТОЛОГІЧНИХ ЗАХВОРЮВАННЯХ У США І В УКРАЇНІ ДЛЯ ОПТИМІЗАЦІЇ ФОРМУЛЯРНОГО ЗАБЕЗПЕЧЕННЯ

Заболотня З. О., Заліська О. М., Калинюк Т. Г.
Кафедра організації і економіки фармації, технології ліків та фармакоекономіки ФПДО Львівського національного медичного університету імені Данила Галицького, м. Львів, Україна
olzaliska@ukr.net

Вступ. Закордонними авторами було проведено оцінку показника «прихований тягар хвороби» для дорослих і підлітків з дерматологічними патологіями, показано значний вплив на зниження ВВП та необхідність удосконалення схем лікування для поліпшення якості життя, зниження днів непрацездатності, відсутності на роботі. При поширених дерматологічних захворюваннях необхідним є застосування екстемпорально виготовлених лікарських засобів (ЕЛЗ) відповідно до стану пацієнта, віку, стадії патології.

Мета. Метою нашого дослідження було провести аналіз ЛЗ, які виготовляються екстемпорально і рекомендовані у США для лікування поширених дерматологічних захворювань: акне, екзема, атопічний дерматит і псоріаз та порівняти з рецептурою аптек, які реально зберегли виготовлення в Україні, щоб обґрунтувати перелік основних ЕЛЗ, які необхідно виготовляти в аптеках для дерматологічних хворих та забезпечити відповідну інформаційну роботу для лікарів і післядипломну освіту фармацевтів.

Матеріали і методи. Узагальнення міжнародних публікацій про переліки ЕЛЗ, які готують в аптеках в США та країнах ЄС при поширених дерматологічних захворюваннях. Аналіз реальної рецептури при дерматологічних захворюваннях з 30 аптек з виробничими відділами 4 областей України за 2023-2024 роки в час дії воєнного стану.

Результати. Виявлено, що у США є рекомендований перелік основних ЕЛЗ, які повинні знати усі лікарі–дерматологи: це 24 ЕЛЗ при лікуванні акне та шрамів від акне та 9 прописів ЕЛЗ при екземі/псоріазі, вони включають комбіновані лікарські форми. Це, наприклад, такі ЕЛЗ:

- ✓ Niacinamide 4%/Biotin 0.1%/Lipoic Acid 0.5% Topical Gel;
- ✓ Clindamycin 1%/Benzoyl Peroxide 5%/Vitamin D3 0.5%/Superoxide Dismutase Topical Cream (Clarifying™);
- ✓ Spironolactone 2.5%/Clindamycin 1%/Tetrahydrocurcuminoids Topical Cream (VersaBase ®);
- ✓ Spironolactone 2.5%/Clindamycin 1%/Tetrahydrocurcuminoids Topical Cream (VersaBase ®) та інші.

Слід відзначити, що усі лікарі мають компетентності з таких прописів ЕЛЗ, виписують їх при поширених дерматологічних захворюваннях: акне, ек-

зема, атопічний дерматит і псоріаз. Ці основні ЕЛЗ є включені у формуляри дерматологічних лікарень США.

Для збереження виробничої функції аптек, фармацевтичних кадрів, які вміють виготовляти extempore ЛЗ, як це діє у США, в країнах ЄС, в Україні необхідно визначити Перелік основних ЕЛЗ, які часто призначаються лікарями-спеціалістами при дерматологічних захворюваннях.

Проведено вивчення реальної рецептури при дерматологічних захворюваннях з 30 аптек з виробничими відділами у Львівській, Івано-Франківській, Волинській та Хмельницькій областях України за 2023-2024 роки в час дії воєнного стану. Нами виявлено, що наявні більше 80 прописів ЕЛЗ, які готуються у досліджуваній сукупності аптек 4 областей України .

Також системно необхідно оновлювати локальні формуляри профільних медичних установах та проводити інформаційну роботу з лікарями-дерматологами, які повинні знати про наявність аптек в межах міста чи області, які готують ЕЛЗ, оскільки молоді фахівці не знають про наявність таких виробничих аптек.

Встановлено, що необхідно постійно вивчати ЕЛЗ на до- і післядипломному рівнях і для безперервного професійного розвитку (БПР) лікарів і фармацевтів. Знання рецептури ЕЛЗ є важливим аспектом, інформаційна робота і співпраця з лікарями, які відвикли виписувати рецепти.

Агресивне рекламування лише готових дерматологічних ЛЗ дезінформує пацієнта і не сприяє прихильності до лікування ЕЛЗ, оскільки лікарські форми, виготовлені в аптеці, ніколи не рекламувалися, але їх ефективність доведена практичним використанням на тисячах пацієнтів протягом століття. Ще однією вагомою перевагою ЕЛЗ є відсутність фальсифікованих ліків, тому пацієнт гарантовано отримує якісний ефективний препарат за призначенням лікаря.

Висновки. Аналіз ЕЛЗ у США та в країнах ЄС показав, що є наявний перелік основних ЕЛЗ, які всі лікарі повинні вміти виписувати і призначати дерматологічним хворим. Нами виділено сукупність ЕЛЗ, які реально виготовляються в аптеках при дерматологічних захворюваннях в Україні під час дії воєнного стану.

Також вважаємо необхідним внести у Ліцензійні умови провадження господарської діяльності вимогу щодо наявності аптеки з виготовленням у мережевих аптеках, які повинні бути гармонізовані з вимогами законодавства ЄС, для збереження і приготування ЕЛЗ та покращення формулярного забезпечення пацієнтів.

Важливим і актуальним є використання підготовлених інформаційного листа та навчально-методичного забезпечення для проведення належної практики студентів фармацевтичних факультетів і проходження інтернатури фармацевтів-інтернів протягом 8 місяців під час заочної практичної частини навчання в Україні, щоб не втратити мистецтво приготування ліків в аптеках.

УДК 339.1:338.1:616.1

МАРКЕТИНГОВИЙ ТА ФАРМАКОЕКОНОМІЧНИЙ АНАЛІЗ ГІПОЛІПІДЕМІЧНИХ ЛІКАРСЬКИХ ЗАСОБІВ, ЩО МІСТЯТЬ СИМВАСТАТИН

Жадько С. В., Севрюков О. В., Бабічева Г. С., Чуфицький Є. О.
Національний фармацевтичний університет, Харків, Україна
svzhadkopharm@gmail.com

Анотація. Метою роботи є маркетинговий та фармакоекономічний аналіз лікарських засобів (ЛЗ) симвастатину та дослідження тенденцій споживання симвастатину на рівні аптечної мережі. Використано методи маркетингових та фармакоекономічних досліджень: аналіз медико-технологічних документів, економіко-статистичний, формальний VN-аналіз, метод «мінімізації витрат». За результатами формального VN-аналізу симвастатин віднесений до категорії V (vital). Встановлено, що найбільшу частку у структурі продажів ЛЗ симвастатину в аптечній мережі мають таблетки в дозуванні 20 мг (69,4%). Визначена найменша вартість гіполіпідемічної терапії на місяць при застосуванні симвастатину у дозуванні 20 мг і 40 мг. Встановлено, що найдешевші ЛЗ мають низькі обсяги продажів в атечній мережі протягом року.

Ключові слова: *маркетинговий аналіз, фармакоекономічні дослідження, формальний VN-аналіз, метод мінімізації витрат, симвастатин*

Abstract. The aim of the study is to conduct a marketing and pharmacoeconomic analysis of simvastatin drugs and to explore consumption trends of simvastatin at the pharmacy network level. Methods of marketing and pharmacoeconomic research were used: analysis of medical-technological documents, economic-statistical analysis, formal VN-analysis, and the cost-minimization method. Based on the results of the formal VN-analysis, simvastatin was classified as category V (vital). It was established that the largest share of simvastatin drug sales in the pharmacy chain consists of 20 mg tablets (69.4%). The lowest cost of hypolipidemic therapy per month was determined for simvastatin at doses of 20 mg and 40 mg. It was also found that the cheapest drugs had low sales volumes in the pharmacy network throughout the year.

Keywords*: marketing analysis, pharmacoeconomic research, formal VN-analysis, cost-minimization method, simvastatin*

Вступ. Серцево-судинні захворювання є провідною причиною смертності на світовому рівні, спричиняючи понад 18 млн смертей щороку, що становить приблизно 31% від загальної кількості смертей. За даними Всесвітньої організації охорони здоров'я (ВООЗ), близько 85% цих випадків обумовлені інфарктами та інсультами [6]. В Україні 67% всіх смертей спричинені серцево-судинними захворюваннями. Симвастатин, як представник групи статинів, є другим за частотою використання лікарським засобом (ЛЗ) для зниження рівня

холестерину, поступаючись лише аторвастатину. Лише в 2021 р. у світі було виписано понад 70 млн рецептів на симвастатин [7].

У контексті зростання витрат на охорону здоров'я, раціональне використання ресурсів стає дедалі важливішим. Фармакоекономічні та маркетингові дослідження симвастатину відіграють ключову роль в оптимізації витрат, покращенні доступу до лікування та підвищенні якості надання медичних послуг, що відповідає потребам пацієнтів та медичних установ.

Мета. Метою роботи є маркетинговий та фармакоекономічний аналіз ЛЗ симвастатину та дослідження тенденцій споживання симвастатину на рівні аптечної мережі.

Методи дослідження. У роботі використано методи кабінетних маркетингових досліджень (аналіз документів), економіко-статистичний метод та методи фармакоекономічних досліджень (формальний VN-аналіз, метод «мінімізації витрат»). Проаналізовано Державний реєстр лікарських засобів України; Державний формуляр лікарських засобів (випуск шістнадцятий); Національний перелік основних лікарських засобів; Перелік лікарських засобів, які підлягають реімбурсації за програмою державних гарантій медичного обслуговування населення; клінічний протокол. Досліджено структуру обсягів продажів ЛЗ симвастатину в аптечній мережі національного рівня за календарний рік. Аналіз обсягів продажів ЛЗ на рівні аптечної мережі дозволив отримати більш релевантні дані щодо структури споживання ЛЗ та зменшити вплив випадкових чинників, таких як особливості розташування окремих аптек та контингенту їх відвідувачів, можливі ускладнення в роботі окремих аптек чи перебої з постачанням ЛЗ. Розрахунок вартості курсу лікування ЛЗ симвастатину на місяць проведений за середньозваженими роздрібними цінами аптечної мережі.

Результати дослідження. Згідно з концепцією профілактики серцево-судинних подій шляхом стабілізації атеросклеротичних бляшок, статини призначаються пацієнтам для постійного пожиттєвого прийому.

Завдяки наявності генеричних ЛЗ вітчизняного й зарубіжного виробництва, симвастатин є одним із найбільш доступних статинів, що робить його популярним вибором для пацієнтів з обмеженими фінансовими можливостями.

Використання симвастатину, котрий досить часто призначають лікарі в Україні, переважно зумовлено його наявністю в переліку препаратів, які підлягають реімбурсації. Разом з тим досягнення цільових рівнів холестерину ліпопротеїдів низької щільності (ЛПНЩ) у пацієнтів із дуже високим серцево-судинним ризиком у разі призначення симвастатину неможливо, а з високим серцево-судинним ризиком — вкрай проблематично [3].

Нами проведено формальний VN аналіз ЛЗ симвастатину за вітчизняними нормативними та медико-технологічними документами (табл. 1).

Таблиця 1

Узагальнені результати формального VN-аналізу ЛЗ симвастатину

Назва документа	Результат пошуку	Категорія VN-аналізу
Державний формуляр лікарських засобів	17 записів про ЛЗ симвастатину	V
Національний перелік основних лікарських засобів	наявний	V
Перелік лікарських засобів, які підлягають реімбурсації за програмою державних гарантій медичного обслуговування населення, станом на 23.08.2024 р.	12 записів про ЛЗ симвастатину	V
Уніфікований клінічний протокол первинної, вторинної (спеціалізованої) та третинної (високоспеціалізованої) медичної допомоги «Профілактика серцево-судинних захворювань»	симвастатин рекомендований для лікування у режимі «низької інтенсивності»	V

За результатами пошуку у Державному реєстрі лікарських засобів України за АТС-кодом С10АА01 виявлено 21 запис про реєстрацію ЛЗ, що містять симвастатин (з урахуванням різних дозувань і кількості доз в упаковці) у формі таблеток, вкритих плівковою оболонкою. ЛЗ симвастатину зареєстровані в Україні у дозуваннях 10, 20 і 40 мг. DDD симвастатину складає 30 мг.

У Державному формулярі виявлено один запис про ЛЗ симвастатину вітчизняного виробника (ФК «Здоров'я») та 16 записів про ЛЗ симвастатину зарубіжних виробників (із Північної Македонії, Індії, Туреччини, Німеччини, Румунії, Польща, Угорщини). Враховуючи те, що Державний формуляр включає ліки з доведеною ефективністю, допустимим рівнем безпеки, використання яких є економічно прийнятним, ЛЗ симвастатину, що внесені до формуляру, можна віднести до категорії V (vital) формального VN-аналізу [1].

Препарати симвастатину за міжнародною непатентованою назвою внесені до Національного переліку основних лікарських засобів [2]. До Переліку лікарських засобів, які підлягають реімбурсації за програмою державних гарантій медичного обслуговування населення, станом на 23 серпня 2024 р., внесені 12 ЛЗ симвастатину) [4].

Симвастатин внесений до уніфікованого клінічного протоколу первинної, вторинної (спеціалізованої) та третинної (високоспеціалізованої) медичної допомоги «Профілактика серцево-судинних захворювань» [5]. Згідно з цим документом, симвастатин з метою досягнення зниження рівня холестерину ЛПНЩ на 30 % та більше від початкового рівня (лікування «низької інтенсивності»)

показав ефект у рандомізованому контрольованому дослідженні при застосуванні в дозі 10 мг.

У 2023 р. в аптечній мережі реалізовувались 13 асортиментних позицій ЛЗ симвастатину вітчизняних і зарубіжних виробників у дозуваннях 10, 20 і 40 мг. Разом протягом року відпущено 5,26 млн добових доз симвастатину на сумму 8,7 млн грн. Встановлено, що найбільшу частку у структурі продажів ЛЗ симвастатину (69,4%) мають таблетки в дозуванні 20 мг, на другому місці — дозування 40 мг (29,5% відпущених добових доз) (рис.).




Рис. Розподіл відпущених добових доз симвастатину за дозуваннями

Для ЛЗ симвастатину у дозуваннях 20 і 40 мг визначено вартість лікування на місяць із застосуванням методу фармакоекономічних досліджень «мінімізація витрат». Встановлено, що найменша роздрібна ціна за дозу симвастатину 20 мг становить 1,11 грн (ЛЗ «Симвастатин», «Марксанс Фарма ЛТД»), а найбільша — 1,84 грн (ЛЗ «Симвастатин Сандоз», «Сандоз Групп Саглик Урунлерис»). Ці ж ЛЗ мають найменшу і найбільшу вартість гіполіпідемічної терапії на місяць (33,30 і 55,15 грн відповідно). При цьому варто відзначити порівняно низькі обсяги продажів (512,5 упаковок за рік, або 0,24% обсягів продажів у натуральних показниках) для найбільш економічно доступного ЛЗ «Симвастатин», таблетки 20 мг № 28 («Марксанс Фарма ЛТД»). Найменша роздрібна ціна за дозу симвастатину 40 мг становить 2,16 грн (ЛЗ «Симвастатин», «Марксанс Фарма ЛТД»), а найбільша — 2,71 грн (ЛЗ «Вазостат Здоров'я», ФК «Здоров'я», Україна). Ці ж ЛЗ мають найменшу і найбільшу вартість гіполіпідемічної терапії на місяць (64,74 і 81,30 грн відповідно). Низькими є обсяги продажів для найбільш економічно доступного ЛЗ «Симвастатин», таблетки 40 мг № 28 «Марксанс Фарма ЛТД» (194 упаковки за рік, або менше 0,01%). Можна припустити, ці ЛЗ не були постійно присутні в асортименті аптек мережі протягом досліджуваного періоду, що обмежило можливості пацієнтів отримувати найбільш економічно доступний варіант лікування.

Висновки. Симвастатин внесений до усіх проаналізованих медико-технологічних документів, що дозволяє віднести його до категорії V (vital) за результатами формального VN-аналізу. За даними аптечної мережі визначена найменша вартість гіполіпідемічної терапії на місяць при застосуванні симвастатину у дозуванні 20 мг і 40 мг. Встановлено, що найдешевші ЛЗ мають низькі обсяги продажів протягом року.

Список використаних джерел інформації:

1. Державний формуляр лікарських засобів. Випуск шістнадцятий. — Київ, 2024. — Режим доступу : chrome-extension://efaidnbmnnnibpcajpcglclefindmkaj/https://moz.gov.ua/uploads/10/54241-dn_418_12032024_dod.pdf (дата звернення: 10.10.2024).
2. Деякі питання державного регулювання цін на лікарські засоби і вироби медичного призначення (в редакції постанови КМУ від 23 груд. 2021 р. № 1431) : постанова КМУ від 25 берез. 2009 р. № 333. — Режим доступу : https://zakon.rada.gov.ua/laws/show/333-2009-%D0%BF#Text (дата звернення: 14.10.2024).
3. Косуєв Г. І. Сучасні грані статинотерапії в практиці сімейного лікаря: запитання та відповіді. 2021. № 4 (497). С. 12-14.
4. Про затвердження Переліків лікарських засобів і медичних виробів, які підлягають реімбурсації за програмою державних гарантій медичного обслуговування населення, станом на 23 серпня 2024 р. [Електронний ресурс] : наказ МОЗ України від 04 верес. 2024 р. — Режим доступу: https://moz.gov.ua/uk/decrees/nakaz-moz-ukrayini-vid-04-09-2024-1537-pro-zatverdzhennya-perelikiv-likarskih-zasobiv-i-medichnih-virobiv-yaki-pidlyagayut-reimbursaciyi-za-programoyu-derzhavnih-garantij-medichnogo-obslugovuvannya-naselennya-stanom-na-23-serpnya-2024-roku (дата звернення: 12.10.2024).
5. Профілактика серцево-судинних захворювань. Уніфікований клінічний протокол первинної, вторинної (спеціалізованої) та третинної (високоспеціалізованої) медичної допомоги. — Режим доступу: chrome-extension://efaidnbmnnnibpcajpcglclefindmkaj/https://www.dec.gov.ua/wp-content/uploads/2019/11/2016_564_ykpmd_pssz.pdf (дата звернення: 14.10.2024).
6. Cardiovascular diseases (CVDs). — Режим доступу: https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds) (дата звернення: 16.10.2024).
7. World Heart Report 2023: Confronting the World's Number One Killer. Geneva, Switzerland. World Heart Federation. 2023. — Режим доступу: chrome-extension://efaidnbmnnnibpcajpcglclefindmkaj/https://world-heart-federation.org/wp-content/uploads/World-Heart-Report-2023.pdf. (дата звернення: 16.10.2024).

АНАЛІЗ ЦЕНТРАЛІЗОВАНИХ ДЕРЖАВНИХ ЗАКУПІВЕЛЬ ЛІКАРСЬКИХ ЗАСОБІВ ДЛЯ ЕНДОКРИННОЇ ТЕРАПІЇ РАКУ МОЛОЧНОЇ ЗАЛОЗИ

Адонкіна В. Ю.[1], Міщенко О. Я.[2], Грешко Ю. І.[2]

Національний фармацевтичний університет, м. Харків, Україна
[1]Кафедра організації, економіки і управління фармацією,
[2]Кафедра клінічної фармакології
Інституту підвищення кваліфікації спеціалістів фармації
uef-ipksf@nuph.edu.ua,
clinpharmacol_ipksph@nuph.edu.ua

Вступ. Рак молочної залози (РМЗ) у жінок посідає перше місце у структурі захворюваності на злоякісні новоутворення серед дорослого жіночого населення більшості країн світу й України і залишається однією з головних проблем сучасної онкології. Сьогодні, дуже важливим при лікуванні злоякісних новоутворень є вірний вибір терапевтичного підходу і розуміння прогнозу перебігу захворювання. Сучасні методи діагностики дають можливість одержати чітке розуміння молекулярного підтипу РМЗ, що включає різні варіанти пухлин, серед яких більшість люмінальні (із експресією рецепторів естрогену та/або прогестерону) HER2- негативні РМЗ, питома вага яких сягає 70 %. Підвищення ефективності лікування метастатичного раку молочної залози є актуальною проблемою клінічної онкології.

Враховуючи актуальність вирішення питання щодо підвищення рівня соціально-економічної доступності лікарських препаратів (ЛП) для ендокринної терапіїї РМЗ, доцільним є проведення аналізу державних закупівель цієї групи препаратів.

Мета. Провести аналіз державних закупівель ЛЗ, які здійснювались МОЗ за рахунок центрального бюджету протягом 2017-2024 рр.

Матеріали та методи. Аналіз динаміки показників державних закупівель ЛП здійснювався у діючих на момент проведення торгів цінах. Використовувалися дані, які представлені у відритому доступі на інформаційній платформі PROZZORO (вартість ЛП та суми закупівель у відкритих торгах).

Результати. За даними аналізу діючого в Україні Уніфікованого клінічного протоколу первинної, вторинної (спеціалізованої), третинної (високоспеціалізованої) медичної допомоги рак молочної залози, 2015 р., Настанови Американського товариства клінічної онкології (2021 р.), Британського національного формуляра (2020 р.) та Державного формуляра ЛЗ України (2024 р.) визначена вибірка з 6 міжнародних непатентованих назв (МНН) ЛП для ендокринної терапії РМЗ: 3 МНН інгібіторів ароматази – анастрозол, летрозол та екземестан і 3 МНН антиестрогенних препаратів – тамоксифен, тореміфен і фулвестрант.

Аналізуючи дані державних закупівель ЛП, які здійснювалися централізовано на рівні МОЗ України протягом 2017-2024 рр. та використовувалися для ендокринної терапії хворих на РМЗ, можна зробити наступні висновки.

Привертає увагу непропорційний характер закупівель у вартісних вимірах на початку досліджуваного періоду. За період з 2017 по 2024 рр. державою було закуплено (6 препаратів за МНН), які використовуються для ендокринної терапії хворих на РМЗ, на суму 390211379,16 грн та в натуральному виразі – 36486120 умовних упаковок. Таким чином, розрахована середня вартість однієї умовної упаковки ЛП дорівнювала 10,70 грн. У діючих цінах у 2017 р. було закуплено ЛП на суму 2810798,86 грн, а в наступному році мало місце значне (на 142,58%) збільшення обсягів закупівель до 6818531,24 грн. За даними 2019 р. спостерігалося незначне зниження обсягів закупівель ЛП. У наступному році було придбано препаратів на суму 6374748,57 грн, що на 6,5% менше, ніж за даними попереднього року (6818531,24 грн). Позитивним є той факт, що в наступному 2020 р. ми спостерігали збільшення (на 73,6%) сум закупівель ЛП, порівняно з даними попереднього 2019 р., до 11066451,23 грн. Тенденція до збільшення обсягу закупівель препаратів зберігалася й у 2021 р. Державою було придбано ЛП на суму 57170415,80 грн, що на 416,61% більше, ніж за даними попереднього 2020 р. У наступному 2022 році ЛП було придбано на суму 71731683,05 грн., що на (25,47%) більше ніж у минулому 2021 році (57170415,80 грн.).

У наступних роках зберіглася чітка тенденція до зростання сум закупівель ЛП для ендокринної терапії хворих на РМЗ за бюджетні кошти. Так у 2023 р. ЛП було придбано на суму 99551707,66 грн., що відповідно більше за попередній період на (38,78%), а в наступному 2024 році у порівнянні із 2023 р. сума закупівель збільшилася на 35,29% і становила відповідно (134687042,75 грн.)

У випадку порівняння даних державних закупівель ЛП, які здійснено у 2024 р. з показниками 2017 р., встановлено, що обсяг цих закупівель значно виріс на 4691,76% у діючих цінах.

Висновки. В ході проведеного дослідження був встановлений нерівномірно зростаючий характер змін обсягів державних закупівель ЛП для ендокринної терапії хворих на РМЗ протягом 2017–2024 рр. Спостерігалась тенденція до зростання вартості закупівель у діючих цінах. Лише у 2019 р. спостерігалася незначна низхідна тенденція сум закупівель на препарати з групи інгібіторів ароматази та антиестрогенних препаратів у порівнянні з попереднім роком. Загалом, на початку дослідження сума закупівель сягала 2810798,86 грн., а наприкінці дослідження - 134687042,75 грн, що становить 4691,76% у діючих цінах.

Розділ 8

ТЕХНОЛОГІЧНІ (НАНОТЕХНОЛОГІЧНІ) АСПЕКТИ РОЗРОБКИ, ВПРОВАДЖЕННЯ ТА ВИКОРИСТАННЯ ЛІКАРСЬКИХ ЗАСОБІВ

СТРАТЕГІЇ ПОКРАЩЕННЯ ДОСТАВЛЯННЯ ЛІКІВ В ОКО

Сагайдак-Нікітюк Р. В., Губченко Т. Д., Шевченко В. О.
Інститут підвищення кваліфікації спеціалістів фармації НФаУ, м. Харків, Україна
Кафедра фармацевтичної технології, стандартизації та сертифікації ліків
phtaqa@nuph.edu.ua

Вступ. Око є одним з найчутливіших органів людського організму, схильних до різних захворювань, які можуть негативно вплинути на якість життя людини. Найпоширенішими хворобами очей є демодекоз повік, синдром сухого ока, кон'юктивіт, глаукома. За прогнозами науковців, до 2050 року кількість людей, які можуть втратити зір збільшиться і буде становити понад 10 млн осіб. Крім того очікується зростання загальної поширеності катаракти, діабетичної ретинопатії, глаукоми та вікової дегенерації жовтої плями майже до 70 млн дорослих, з яких близько 17,8 млн випадків, будуть спричинені ранньою стадією вікової дегенерації жовтої плями.

Варто зазначити, що головною проблемою офтальмологічної терапії є підтримання ефективної концентрації лікарського засобу в місці дії протягом відповідного періоду часу для отримання очікуваної терапевтичної відповіді.

Мета. Метою є дослідження стратегій покращення доставляння ліків в очі.

Матеріали та методи. Методи: узагальнення, синтезу та аналізу.

Результати. Ефективне лікування очних захворювань можна досягти завдяки правильному застосуванню офтальмологічних препаратів, заснованому на нових фармацевтичних підходах.

Сьогодні існує три основні шляхи доставки ліків у націлену ділянку ока: місцево, системно, внутрішньоочно.

Місцевий шлях (абсорбція крізь рогівку) є основним вибором через високу прихильність пацієнта та простоту введення.

Очні краплі становлять приблизно 90% усіх лікарських форм місцевого застосування у вигляді розчинів, суспензій та емульсії. Для досягнення терапевтичного рівня препарату в оці необхідний частий прийом препарату.

Традиційні очні краплі зручні й прості у використанні, вони не дуже ефективні і тільки невелика кількість дози доставляється до окресленої мети, більшість втрачається через механізми очищення. Однак є певні стратегії, які можуть застосовуватися для покращання біодоступності ліків. По-перше, можна використовувати підсилювачі розчинності, для покращання концентрації лікарського засобу в складі; їх більша кількість у лікарській формі може означати підвищену біодоступність. Ця стратегія може дозволити застосувати крапельку меншого розміру, яка буде менш сприйнятлива до втрат через дренаж, викликаний розривом рефлексу та кліпання. По-друге, склад може бути розроблений у формі, що протистоїть очищенню. Ці лікарські форми зберігаються протягом тривалого періоду, тому вони мають більше часу для взаємодії з тканинами ока.

Потім до складу препарату можуть бути уведені підсилювачі проникнення ліків, що сприяють їх проходженню через рогівку.

Системне введення офтальмологічних препаратів (наприклад, у формі таблеток) часто не застосовується через дуже мале співвідношення ока і розміру тіла.

Внутрішньоочні підходи використовуються для доставляння безпосередньо в задній сегмент ока.

Збільшення проникнення лікарського засобу та його біодоступності можна досягти шляхом зміни властивостей самого засобу або системи доставки. Одним із таких способів є модифікація властивостей препарату за допомогою стратегії проліки. Підхід із проліками змінює фізико-хімічні властивості ліків для їх кращого усмоктування шляхом пасивної чи активної дифузії. Як тільки проліки досягають тканини рогівки, клітинні ферменти розщеплюють їх до активної форми. Так, *дипівефрин* (*пропін, алерган*) є проліками на основі складного ефіру адреналіну, який демонструє в 17 разів вищу проникність рогівки, що сприяє збільшенню біодоступності адреналіну в тканинах рогівки в 10 разів порівняно з немодифікованим препаратом. Стратегія проліків також корисна для зниження його дози з вузьким терапевтичним індексом.

Фармацевтичні підходи, засновані на нанотехнологіях у розробці очних крапель, що складаються з мукоадгезивних полімерів, хітозану та гіалуронової кислоти, являють собою нові стратегії для ефективного лікування захворювань очей. Ці інноваційні системи наночастинок спрямовані на підвищення біодоступності ліків в оці.

Очні суспензії є дисперсією дрібнодисперсних нерозчинних частинок лікарського засобу, суспендованих у водному середовищі, що містить диспергувальні та солюбілізувальні агенти.

Останнім часом місцеві системи перетворилися на мікроемульсії в/о або наноемульсії. Дисперсна фаза являє собою нанокраплі діаметром 50-200 нм, неоднорідно дисперговані у незмішуваній зовнішній фазі. Ці склади корисні для доставляння як гідрофільних, так і ліпофільних препаратів, проте вони все ще схильні до нестабільності (коалесценція, флокуляція і дозрівання).

Офтальмологічні емульсії можуть забезпечити переваги у покращанні розчинності та біодоступності ліків раніше не розчинних у воді препаратів.

Мікроемульсія – одна з найперспективніших систем доставки офтальмологічних препаратів. Для зниження міжфазного натягу між олійною та водною фазами, а також для полегшення процесу диспергування та забезпечення гнучкої плівки навколишньої краплі в них використовується вища концентрація поверхнево активних речовин.

Висновки. Отже, ефективне внутрішньоочне доставлення ліків є серйозною проблемою через наявність різних механізмів виведення та фізіологічних бар'єрів, які спричиняють низьку біодоступність в оці після місцевого застосування.

ПЕРСПЕКТИВА СТВОРЕННЯ НОВИХ ЗАСОБІВ У ВИГЛЯДІ ЛАЙНЕРІВ

Іванова А. Д., Сліпченко Г. Д.
Національний фармацевтичний університет, м. Харків, Україна
Кафедра промислової технології ліків та косметичних засобів
an.ivanova2014@gmail.com

Вступ. Лайнери, що одягаються на культю, відіграють ключову роль у сучасному протезуванні. Вони забезпечують комфортне носіння протеза, захищають шкіру від подразнень та нерівномірного тиску. Завдяки новітнім матеріалам та технологіям, сучасні лайнери можуть містити лікувальні компоненти, сприяючи швидкому загоєнню та відновленню шкіри.

Мета. Вивчення сучасних досягнень у розробці та виготовленні лайнерів (вкладок) під протези, визначення перспективних матеріалів та технологій, які забезпечують комфорт, довговічність і зручність у використанні.

Матеріали та методи. Узагальнення наукових публікацій і технічних досліджень щодо різних типів лайнерів, матеріалів, які використовуються для їх виготовлення, та технологій, що застосовуються при їх виробництві.

Результати дослідження. Лайнери поділяються на кілька типів залежно від матеріалів і призначення: 1) силіконові лайнери – забезпечують високу гіпоалергенність і відмінну амортизацію. Силікон добре підходить для пацієнтів з чутливою шкірою, оскільки знижує ймовірність подразнень; 2) гелеві лайнери мають високі показники еластичності та здатності адаптуватися до форми культі. Гелі створюють ефект рівномірного розподілу тиску, що зменшує навантаження на певні зони; 3) пінопластові лайнери використовуються для додаткового комфорту, оскільки мають легку вагу та забезпечують м'якість при контакті зі шкірою. Кожен з цих матеріалів має свої переваги і недоліки.

Аналіз технологій виготовлення показав, що сучасні методи, такі як 3D-друк і використання біоматеріалів, дозволяють виготовляти лайнери з більш точною анатомічною відповідністю та покращеними характеристиками. Інноваційні розробки також включають інтеграцію сенсорів для моніторингу стану шкіри та тиску, що дозволяє уникати ускладнень під час користування протезом.

Лайнер є важливим компонентом при протезуванні кінцівок. Він створює бар'єр між шкірою та протезом, зменшуючи тертя та тиск, що сприяє кращій фіксації протеза, зменшує ризик пошкодження шкіри і, як наслідок, підвищує комфорт при використанні.

Висновок. Лайнери відіграють ключову роль у забезпеченні комфорту і ефективності використання протезів. Завдяки використанню нових матеріалів і технологій, вони забезпечують кращий захист шкіри, розподіл тиску і загальне поліпшення якості життя пацієнтів. Перспективними напрямками розвитку є подальше вдосконалення матеріалів, таких як гелі та силікон, а також впровадження інтелектуальних систем моніторингу.

РОЗРОБКА МЕТОДИКИ ДЕТЕКТУВАННЯ КОМПЛЕКСУ СУПУТНІХ ДОМІШОК НАТРІЙ 2-((4-АМІНО-5-(ТІОФЕН-2-ІЛМЕТИЛ)-4*H*-1,2,4-ТРІАЗОЛ-3-ІЛ)ТІО)АЦЕТАТУ ЗА ДОПОМОГОЮ ВЕРХ-МС

Усенко Д. Л., Каплаушенко А. Г.

Запорізький державний медико-фармацевтичний університет, Україна

Кафедра фізколоїдної хімії

usenko.d.l@ukr.net

Вступ. Проведення даного дослідження зумовлено сучасними вимогами до створення нових ефективних і безпечних лікарських засобів, а також необхідністю розробки точних методик аналізу активних фармацевтичних інгредієнтів (АФІ) та їх домішок. Домішки можуть суттєво впливати на якість і безпеку препаратів, тому їх своєчасне виявлення є критично важливим на етапах розробки і виробництва. Метод високоефективної рідинної хроматографії у поєднанні з мас-спектрометрією (ВЕРХ-МС) є сучасним інструментом для детектування домішок та продуктів деградації на низьких концентраціях. Це має особливе значення для подальшого вивчення метаболітів і фармакокінетики нових лікарських засобів.

Метою даної роботи є розробка та оптимізація методики якісного та кількісного визначення комплексу супутніх домішок натрій 2-((4-аміно-5-(тіофен-2-ілметил)-4*H*-1,2,4-тріазол-3-іл)тіо)ацетату, потенційного АФІ, з використанням ВЕРХ-МС.

Матеріали та методи. Система ВЕРХ Agilent 1260 Infinity. Для проведення дослідження було використано діодно-матричний детектор та одноквадрупольний мас-спектрометричний детектор Agilent 6120 з іонізацією в електроспреї (ESI). Колонка Zorbax SB-C18; 30 мм х 4,6 мм; 1,8 мкм.

Результати. Було отримано залежності інтенсивності сигналу мас-детектора від трьох ключових параметрів: температури сушильного газу, напруги фрагментації та тиску газу розпилювача. Ці залежності дозволили визначити оптимальні умови для якісного та кількісного детектування комплексу супутніх домішок натрій 2-((4-аміно-5-(тіофен-2-ілметил)-4*H*-1,2,4-тріазол-3-іл)тіо)ацетату у вигляді 2-(тіофен-2-іл)ацетогідразиду, 4-аміно-5-(тіофен-2-ілметил)-2,4-дигідро-3*H*-1,2,4-тріазол-3-тіону та калій 2-(2-(тіофен-2-іл)ацетил)гідразин-1-карбодіонату.

Висновки. Було встановлено, що підвищення температури сушильного газу покращує іонізацію зразків, що сприяє збільшенню інтенсивності сигналу. Напруга фрагментації впливає на ефективність розщеплення молекул, дозволяючи досягти кращого розділення піків домішок. Оптимальний тиск газу розпилювача сприяє стабільності сигналу та забезпечує точніше детектування. Оптимізація цих параметрів дозволила розробити чутливу методику для виявлення та кількісного визначення супутніх домішок натрій 2-((4-аміно-5-(тіофен-2-ілметил)-4*H*-1,2,4-тріазол-3-іл)тіо)ацетату, що підвищує надійність і точність аналізу. Розроблена методика може бути застосована для контролю якості АФІ, що дозволяє своєчасно виявляти домішки і забезпечувати його безпеку.

ПЕРСПЕКТИВНІ ВИДИ ВІТЧИЗНЯНОЇ ЛІКАРСЬКОЇ РОСЛИННОЇ СИРОВИНИ, ДЖЕРЕЛОМ ЯКОЇ Є РОСЛИНИ РОДИНИ *LAMIACEAE*

Хворост О. П., Опрошанська Т. В., Скребцова К. С.
Національний фармацевтичний університет, м. Харків, Україна
Кафедра фармакогнозії та нутриціології
Кафедра фармацевтичної технології, стандартизації та сертифікації ліків
khvorost09101960@gmail.com

Вступ. В Україні до нетрадиційної медицини, за інформацією федерації сприяння охороні здоров'ю, щороку звертається понад п'ять мільйонів людей, народним лікарям довіряють 40-60 % українців. В багатьох країнах світу витрати на народну медицину постійно зростають.

Представники родини *Lamiaceae* поширені у всіх кліматичних зонах та на всіх континентах, об’єднані у понад 200 родів, до яких входить за різними даними від 3500 до 8000 видів, у флорі України – понад 230 видів. Лише у Харківській області представники родини Lamiaceae, які є одними з найпоширенішими у Лісостепової зоні, є компонентами більшості фітоценозів. У природних та культивованих умовах зустрічається 176 видів *Lamiaceae*, які належать до 40 родів. Це становить 5% від загального світового різноманіття родини Глухокропивові. До провідних належить 10 родів родини *Lamiaceae* України, які включають 104 види, що становить 59,1% від загальної кількості видів.

Мета. проаналізувати асортимент лікарської рослинної сировини представників родини *Lamiaceae* народної та наукової медицини, що доступні в Україні.

Матеріали та методи. контент-аналіз доступних джерел інформації з наступною інтерпретацією результатів.

Результати. Критеріями вибору є доступність сировини (поширеність у дикорослому стані або широта культивування), хімічний склад (якщо такі відомості мають місце) та актуальність застосування.

Найвідоміші джерела рослинної сировини фармакопейного ґатунку такі глухокропивні: собача кропива, лаванда, глуха кропива, чебрець, сальвія, розмарин, м'ята, меліса, гісоп, шоломниця. Але поширені як у дикорослому так й у культивованому стані (як декоративні закритого та відкритого ґрунту) дубровник, пахучка, плектрантус, змієголовник, ельшольція, красивоплідник, монарда, молюцела, котовник, горлянка, колеус, багатоколосник, каріоптеріс. Частина з них частково вже фармакогностично досліджена, але поки відсутні затверджені монографії на сировину та лікарські засоби. Безумовно, сировиною мають виступати листя, або трава.

Висновки. Подальше фармакогностичне дослідження нефармакопейної сировини представників родини *Lamiaceae*, зокрема, розхідника, щебрушки, маточника, жабрію є актуальним для створення на їх основі ефективних засобів комплексної терапії актуальної спрямованості дії.

Розділ 9

ДОКЛІНІЧНІ ТА КЛІНІЧНІ ДОСЛІДЖЕННЯ ЛІКАРСЬКИХ ЗАСОБІВ

NOVEL AND PROMISING EMBRYO CULTURE MEDIA IN HUMAN IVF PROGRAMS

Isachenko V., Ostashko V., Isachenko E.
University of Cologne, Germany
Department of Human Reproduction, Research Group for Reproductive Medicine and IVF-Laboratory, Department of Obstetrics and Gynecology
vladimir.isachenko@uk-koeln.de

Introduction. In the field of assisted reproductive technologies (ART), the culture of embryos plays a pivotal role in determining the success of in vitro fertilization (IVF) programs. Over the last decade, the development of novel embryo culture media has significantly improved the outcomes of IVF by providing more optimized conditions that closely mimic the natural intrauterine environment.

Goal. The objective of this study is to review the latest advancements in embryo culture media and evaluate their impact on the success rates of IVF programs. This includes an exploration of modern technologies designed to enhance embryo nourishment and development while reducing stressors that can impair their viability.

Materials and Methods. The literature search was conducted across several major databases, including PubMed, Scopus, Web of Science, and Google Scholar. The search was limited to articles published within the last 5 years, focusing on the latest developments in embryo culture media in the context of IVF.

Results. Various commercially available culture media were analyzed, including EmbryoMax (MilliporeSigma), Vitrolife (Vitrolife Group), and G-IVF (Vitrolife Group), which utilize different combinations of proteins, amino acids, minerals, and antioxidants. Emerging technologies such as synthetic gels (SynthetiGel (Synthetic-Bio)) and nanomaterials (NanoSupport (NanoFertility)) were also examined for their potential to create optimal conditions for embryo development. Continuous real-time monitoring using systems like EmbryoScope (Vitrolife Group) was included to assess embryo growth and adjust culture conditions.

The use of advanced media, coupled with monitoring technologies, demonstrated improvements in embryo quality, reduced miscarriage rates, and overall IVF success. Media such as G-IVF, with its antioxidants, showed a notable reduction in oxidative stress, while EmbryoScope enabled timely culture adjustments, further enhancing outcomes.

Conclusions. The development and refinement of novel embryo culture media are critical to the future of ART. These advancements not only increase the success rates of IVF programs but also offer the potential for further improvements in embryo quality and patient outcomes. Ongoing research and the integration of emerging technologies will continue to enhance fertility treatments and open new possibilities for addressing infertility.

ПЕРСПЕКТИВИ ЗАСТОСУВАННЯ КАТЕХІНІВ, ЯК НОВИХ ТЕРАПЕВТИЧНИХ АГЕНТІВ ДЛЯ ЛІКУВАННЯ ХРОНІЧНОЇ ВЕНОЗНОЇ НЕДОСТАТНОСТІ

Дембіцька Е. С., Цубанова Н. А., Семійон-Лучишин О. В.[1]

Львівська медична академія ім. А. Крупинського, м. Львів

[1]Львівський національний медичний університет ім. Д. Галицького, м. Львів

tsubanova19@gmail.com

Вступ. Пошук нових терапевтичних агентів, на основі лікарської рослинної сировини (ЛРС) для оптимізації фармакотерапії хронічної венозної недостатності продовжується і сьогодні, через обмежений асортимент зареєстрованих препаратів для лікування зазначеної патології.

Мета. Метою даного дослідження було проаналізувати за даними доступних наукових досліджень перспективи застосування катехінів у терапії ХВН.

Матеріали та методи. У роботі використані рекомендовані для інформаційно -пошукових досліджень методи аналізу, дедукції та індукції. Пошук інформаційних джерел проведено на ресурсах PubMed, Medline, Cochrane library.

Результати. Основний механізм дії катехінів пов'язаний з їх здатністю інгібувати оксидативний стрес, який є ключовим фактором у патогенезі ХВН. Катехіни, завдяки своїй антиоксидантній активності, здатні нейтралізувати вільні радикали та знижувати рівень реактивних форм кисню, що, своєю чергою, сприяє збереженню структурної цілісності судинних стінок та поліпшенню їх функціонального стану. Крім того, катехіни мають виражену протизапальну активність, інгібуючи продукцію прозапальних цитокінів, таких як TNF-α, IL-1β і IL-6, а також ферментів ЦОГ і ЛОГ, які відіграють ключову роль у розвитку запальної відповіді. Це дозволяє катехінам ефективно зменшувати запальні процеси в судинній стінці, сприяючи зменшенню набряків та поліпшенню мікроциркуляції.

Дослідження показали, що каткехіни також можуть модулювати сигнальні шляхи NF-κB, MAPK та PI3K/Akt, що додатково підтверджує їх потенціал у зменшенні запалення та оксидативного стресу.Крім антиоксидантних і протизапальних властивостей, катехіни також виявляють антиагрегаційну активність, що запобігає утворенню тромбів, які є частою ускладненням ХВН. Вони здатні інгібувати агрегацію тромбоцитів, що знижує ризик тромбоутворення у венах нижніх кінцівок. Завдяки цьому катехіни можуть забезпечувати не лише лікування симптомів ХВН, але й профілактику її ускладнень, таких як тромбоз глибоких вен і тромбоемболія легеневої артерії.

Висновки. Перспективи використання катехінів у терапії ХВН є значними, враховуючи їх багатовимірну дію на різні ланки патогенезу захворювання.

ВИЗНАЧЕННЯ ФАРМАКОКІНЕТИЧНИХ ПАРАМЕТРІВ ПОХІДНИХ 5-(2,4-, 3,4-ДИМЕТОКСИФЕНІЛ)-3*H*-1,2,4-ТРІАЗОЛ-3-ТІОНІВ ЗА ДОПОМОГОЮ ADME-АНАЛІЗУ

Довбня Д. В., Каплаушенко А. Г.

Запорізький державний медико-фармацевтичний університет,
м. Запоріжжя, Україна
Кафедра фізколоїдної хімії
dima.dovbnya@ukr.net
kaplaushenko@ukr.net

Вступ. Фармакокінетичний профіль нових хімічних сполук є важливим аспектом їх подальшого використання в розробці лікарських препаратів. Похідні 5-(2,4- та 3,4-диметоксифеніл)-3*H*-1,2,4-тріазол-3-тіонів привертають увагу дослідників завдяки своїм потенційним фармакологічним властивостям. Для їх оцінки необхідно визначити основні фармакокінетичні параметри, такі як абсорбція, розподіл, метаболізм та елімінація (ADME), які дозволяють передбачити поведінку сполук в організмі.

Мета. Метою даного дослідження є визначення фармакокінетичних параметрів похідних 5-(2,4- та 3,4-диметоксифеніл)-3*H*-1,2,4-тріазол-3-тіонів з використанням ADME-аналізу для оцінки їх потенціалу як лікарських засобів.

Матеріали та методи. Дослідження здійснено за допомогою комп'ютерних методів прогнозування ADME-профілю. Використано програмні інструменти, що дозволяють оцінити такі параметри, як пероральна біодоступність, розподіл у тканинах, можливі метаболічні шляхи та токсичність. Для кожної сполуки було проведено моделювання, що дало змогу передбачити їхню фармакокінетичну поведінку.

Результати. Отримані результати свідчать про те, що більшість досліджуваних сполук мають прийнятні фармакокінетичні параметри. Зокрема, виявлено сполуки з високою пероральною біодоступністю та помірним розподілом у тканинах, що робить їх перспективними для подальшого вивчення як потенційних лікарських засобів. Виявлено також кілька похідних, які потребують модифікації для поліпшення їхніх ADME-характеристик.

Висновки. Результати ADME-аналізу дозволяють стверджувати, що похідні 5-(2,4- та 3,4-диметоксифеніл)-3*H*-1,2,4-тріазол-3-тіонів мають перспективу для подальшого вивчення як лікарські препарати. Встановлено основні фармакокінетичні параметри, що є важливими для їх майбутньої розробки, а також виділено сполуки з найбільш сприятливим фармакокінетичним профілем.

ВИВЧЕННЯ РЕПАРАТИВНОЇ ДІЇ ЕКСТРАКТУ ТРАВИ *SAPONARIA OFFICINALIS* НА МОДЕЛІ ЛІНІЙНИХ РІЗАНИХ РАН

Цубанова Н. А., Волощук Н. І.[1], Застрижна М. Л.

Львівська медична академія ім. А. Крупинського, м. Львів

[1]Вінницький національний медичний університет ім. М.І. Пирогова, м. Вінниця

tsubanova19@gmail.com

Вступ. У більшості клінічних випадків тривалий перебіг хронічної венозної недостатності (ХВН) супроводжується розвитком венозної виразки ніг (ВВН), яка може бути діагностована у 5-8 % пацієнтів із ХВН. Особливу роль у етіоптогенезі ХВН та ВВН відіграє оксидативний стрес, запальна реакція, стаз крокі, деструкція клітинних мембран.

Важливим аспектом фармакологічної дії флебопротекторів є не лише наявність венотонізувальної, мембранопротекторної, антиоксидантної, протизапальної дії але і здатність чинити репаративний ефект. Зареєстровані на сучасному фармацевтичному ринку флебопротетори не виявляють у повній мірі всі вищезазначені фармакологічні ефекти, превалює переважно венотонізувальна та мембранопротекторна дія. З метою розширення асортименту сучасних флебопротекторів вченими Тернопільского державного медичного університету, під керівництвом проф. Марчишин С. М. було отримано новий густий екстракт трави *Мильнянки лікарської,* надалі ЕТМЛ, із високим вмістом тритерпенових сапонінів, для якого у попередніх дослідженнях було встановлено флебопротеторну дію за умов експериментального тромбофлебіту.

Мета. Метою даного дослідження було оцінити репаративну дію ЕТМЛ.

Матеріали та методи. Репаративну дію вивчали на моделі лінійних різаних ран у щурів. ЕТМЛ вводили у дозі 20 мг/кг, препарат порівняння Ескувіт у дозі 7,2 мг/кг за 7 днів до та 6 днів час експерименту. Календули мазь наносили протягом 6 днів експерименту. На 6 день тварин декапітували та вирізали ділянку шкіри з рубцем. На спеціальному приладі ранотензіометрі проводили випробування міцності рубця. Статистичну обробку результатів проводили за допомогою програми Statistica 6.0 for Windows

Результати. Встановлено, що ведення ЕТМЛ у лікувіально-профілактичному режимі чинило потужну репаративну дію, так міцність рубця за умов введення ЕТМЛ була збільшена у 1,7 рази ($p<0,001$) відносно групи контрольної патології, на 30 % ($p<0,001$) перевищувала активність Ескувіту та на 18 % ($p<0,001$) репаративну активність Календули мазь.

Висновки. Встановлена значна репаративна дія ЕТМЛ у дозі 20 мг/кг, що перевищує ефективність препаратів порівняння Ескувіту та Календули мазь та розширює спектр флебопротекторної дії досліджуваного екстракту. Отримані результати дослідження дозволять рекомендувати ЕТМЛ як перспективний засіб для терапії ХВН та ВВН.

ВИВЧЕННЯ ВПЛИВУ ЕКСТРАКТУ ТРАВИ *TAGETES PATULA L* НА МОДЕЛІ СТРЕПТОЗОТОЦИНОВОГО ДІАБЕТУ

Цубанова Н. А.[1], Волощук Н. І.[2], Галевич Г. Б.[2]

[1]Львівська медична академія ім. А. Крупинського, м. Львів

[2]Вінницький національний медичний університет ім. М.І. Пирогова, м. Вінниця

tsubanova19@gmail.com

Вступ. Патологію підшлункової залози із розвитком цукрового діабету (ЦД) реєструють з кожним роком все частіше. Базові принципи фармакотерапії ЦД включають дієтотерапію, дозовані фізичні навантаження, інсулінотерапію, пероральні гіпоглікемічні засоби та фітопрепарати, причому останні здатні проявити суттєву підтримку стандартного способу лікування (інсулінотерапія, пероральні гіпоглікемічні засоби) на всіх стадіях захворювання.

Відомо, що на ранніх етапах ЦД 2 типу саме фітопрепарати спроможні на певний період замінити базову терапію. Доведено також, що пацієнти, які активно застосовують фітопрепарати, потребують нижчих доз інсуліну або пероральних цукрознижувальних засобів.

Лікарську рослинну сировину (ЛРС) що використовують у ад'ювантній терапії ЦД сьогоді поділяють на дві групи:

- ЛРС, що знижує рівень глюкози у крові (створки квасолі звичайної, чорниця, брусниця, суниця, кропива дводомна, кульбаба, дев'ясил, лопух, топінамбур, солодка, козлятник, насіння льону)
- ЛРС, що стимулюють регенерацію бета-клітин острівців Лангерганса (цикорій, галега, звіробій, кульбаба, золототисячник, спориш; листя горіха волоського; чорниця; насіння льону; лопух, кульбаба, пирій, дев'ясил).

Зазначена ЛРС, досить часто є нестандартизованою за основними діючими речовинами, не має обґрунтованого дозування і використовується для самостійного приготування відварів та настоянок, що не відповідає сучасним вимогам доказової медицини.

Сучасна медицина та фармація потребує фітопрепратів для терапії ЦД на основі стандартизованої ЛРС, обґрунтованим у фармакологічних дослідженнях дозуванням та доведеними ефектами.

З метою отримання лікарського засобу, що можна буде рекомендувати у комплексне лікування ЦД Вченими Тернопільского державного медичного університету, під керівництвом проф. Марчишин С. М. було отримано новий сухий екстракт трави *Tagetes Patula L* (Чорнобривців розлогих), надалі ЕТЧР, із високим вмістом гідроксикоричних кислот (5,07%), флавоноїдів (7,45 %), дубильних речовин (8,68%). У попередніх фармакологічних дослідженнях встановлено антиоксиданту та мембранопротекторну дію ЕТЧР у дозі 25 мг/кг.

.

Мета. Метою даного дослідження було оцінити вплив сухого екстракту трави *Tagetes Patula L* на біохімічні та гістологічні показники за умов стрептозотоцинового діабету.

Матеріали та методи. Експериментальний цукровий діабет моделювали однократним внутрішньоочеревинним введенням щурам масою 180-220 г стрептозотоцину (СПЦ) у дозі 55 мг/кг за стандартною методикою. Досліджуваний сухий екстракт трави чорнобривців розлогих (ЕТЧР) вводили щурам (n=8) у дозі 25 мг/кг у лікувально-профілактичному режимі за 2 тижня до, та 2 тижня після ін'єкції СПЦ. Препарат порівняння Силімарин (таблетки, «Дарсил», ПрАТ «Фармацевтична фірма «Дарниця», Україна) вводили у дозі 25 мг/кг у аналогічному режимі (n=8). Силімарин, як препарат порівняння було обрано, через те що ЕТЧР пропонується рекомендувати надалі як гепатопротектор, тому що печінка найчастіше страждає при медикаментозних ураженнях. Препарат порівняння Глібенкламід (таблетки, 5 мг, «Глібенкламід», АТ «Фармак», Україна) вводили у дозі 0,6 мг/ кг, що відповідає середньотерапевтичній дозі для людини 10 мг на добу (перерахунок зроблено з урахуванням коефіцієнту видової чутливості).

Перегляд мікропрепаратів проводили під світловим мікроскопом Granum, фотографування мікроскопічних зображень здійснювали цифровою відеокамерою Granum DCM 310. Фотознімки обробляли на комп'ютері Pentium 2,4GHz за допомогою програми Toup View.

Статистичну обробку результатів проводили за допомогою програми Statistica 6.0 for Windows

Результати. Розвиток експериментального стрептозотоцинового діабету супроводжувався значними порушеннями вуглеводного та ліпідного обмінів, наслідком яких є патологічні зміни у клітинах підшлункової залози. У тварин групи контрольної патології зареєстровано збільшення рівня глюкози натще у 1,6 рази ($p<0,001$), глікованого гемоглобіну у 1,2 рази ($p<0,001$), тригліцеридів у 1,5 рази ($p<0,001$), холестеролу на 68% ($p<0,001$) відносно групи інтактного контролю.

Лікувально профілактичне введення ЕТЧР у дозі 25 мг/кг вірогідно нормалізувало вищезазначені показники відносно групи контрольної патології. Слід зазначити що за впливом на нормалізацію вуглеводних показників ЕТЧР у дозі 25 мг/кг вірогідно поступався глібенкламіду та достовірно п5еревищував препарат порівняння Силімарин. За відновленням показників ліпідного профілю ЕТЧР у дозі 25 мг/кг перевищував ефективність обох препаратів порівняння.

Органопротекторний вплив ЕТЧР на підшлункову залозу підтверджено гістологічними дослідженнями, в яких встановлено, що досліджуваний екстракт зменшує кількість деструктивно-дистрофічно змінених інсуліноцитів.

Висновки. За умов відтворення етрептозотоцинового діабету встановлена панкреопротекторна дія ЕТЧР у дозі 25 мг/кг за здатністю нормалізувати показники вуглеводного та ліпідного обміну та відновлювати інсуліноцити.

ВИВЧЕННЯ ФАРМАКОДИНАМІЧНИХ ЕФЕКТІВ КВЕРЦЕТИНУ

Білай І. М., Білай С. І., Білай А.І .
Запорізький державний медико-фармацевтичний університет,
м. Запоріжжя, Україна
Кафедра клінічної фармації, фармакотерапії, фармакогнозії
та фармацевтичної хімії
belay250455@gmail.com

Вступ. Сечокам'яна хвороба є одним із найпоширеніших захворювань у світі та вражає осіб працездатного віку. Подібно до уратного нефролітіазу (УН) зростає у світі кількість хворих на метаболічний синдром (МС). За такої умови хворі на МС мають високий ризик розвитку УН.

Мета. Метою дослідження є вивчення впливу кверцетину на функціональний стан та гемодинаміку нирок у пацієнтів наУН, коморбідний з МС.

Матеріали та методи. У першу основну входили хворі (n=61) на УН та МС, яким призначали кверцетин (квертин) по 1 таблетці 40 мл 3 рази на добу, загальноприйняті лікарські засоби (ріабал або дротаверин, дексалгін, уроліт У) та лікарські препарати, які корегують метаболічні процеси (аторвастатин, метформін, алопуринол, ліпразид, вітамін B_6, магнію оксид) протягом 1,5-6 місяців.

До другої групи порівняння (n=63) входили пацієнти на УН, коморбідний з МС, яким застосовували традиційну терапію та лікарські засоби, які корегують метаболічні процеси. До третьої контрольної групи відносилися хворі на УН (n=59), які приймали традиційну терапію.

В якості біохімічних матеріалів застосовували показники, які характеризують функціональний стан нирок (рівень креатиніну, сечовини у сироватці крові, швидкість клубочкової фільтрації (ШКФ)). Для доплерографії нирок визначали індекс резистентності (IR), індекс пульсаційний (IP), індекс діастолічний (S/D).

Результати. У хворих на УН коморбідний з МС спостерігалася кореляція показників функціонального стану та гемодинаміки нирок у процесі лікування протягом 1,5-6 місяців. Найбільш значно це проявлялося у пацієнтів основної групи, що вказувало на значне зниження показників функціонального стану нирок – рівня креатиніну, сечовини і збільшення рівня ШКФ. Водночас показники гемодинаміки нирок S/D, IR, IP також знижувалися суттєво, що свідчило про ефективність призначення квертину на тлі диференційованого застосування урикостатичної та уриколітичної терапії, а також лікарських засобів, які впливають на метаболічні процеси. У хворих 3 групи, яким призначали тільки традиційну терапію, спостерігалося незначне збільшення рівня сечовини та креатиніну, IR, IP та S/D, а рівень ШКФ водночас не суттєво зменшувався.

Висновки. Застосування квертину протягом 1,5-6 місяців лікування сприяло покращенню показників гемодинаміки і функціонального стану нирок.

ЦЕРЕБРАЛЬНІ ПРОТИЗАПАЛЬНІ ВЛАСТИВОСТІ ДИГОКСИНУ ЯК НЕОЧЕВИДНИЙ КОМПОНЕНТ ЙОГО ПРОТИСУДОМНОЇ ДІЇ

Цивунін В. В., Штриголь С. Ю., Северіна Г. І.
Національний фармацевтичний університет, м. Харків, Україна
Кафедра фармакології та клінічної фармації
tsyvunin-vad@ukr.net

Вступ. Нейрозапалення є важливою ланкою патогенезу багатьох хвороб головного мозку, зокрема епілепсії. Доведено, що церебральна ЦОГ-2 та простагландини значно підвищуються під впливом будь-яких патофізіологічних стимулів, що визначає інтерес до вивчення ролі нейрозапалення в розвитку судом. Проте, наскільки значущою є роль протизапальних властивостей серцевих глікозидів (зокрема, дигоксину), а також класичних протиепілептичних препаратів у реалізації їх протисудомній активності, досі залишається невідомим.

Мета. Встановити наявність у дигоксину та вальпроату протизапальних властивостей за впливом на церебральний циклооксигеназний шлях каскаду арахідонової кислоти на моделі хронічного епілептогенезу та *in silico*.

Матеріали та методи. Дослідження проводили *in vivo* на моделі пентилен-тетразолового (PTZ) кіндлінгу у мишей за умов щоденного введення дигоксину, вальпроату та їх комбінації. Наприкінці експерименту у мозку тварин досліджували вміст ЦОГ-1/2, простагландинів PGE2, PGI2, PGF2α, а також тромбоксану TXB2. Шляхом молекулярного докінгу *in silico* досліджували також здатність дигоксину до взаємодії з активними сайтами ЦОГ-1/2.

Результати. Хронічне введення низьких доз PTZ пов'язане з активацією нейрозапалення: збільшується церебральна експресія ЦОГ-1 і 2, а також спостерігається виражений дисбаланс рівнів PGE2, PGI2, PGF2α і TXB2. Дигоксин, вальпроат і особливо їх комбінація нормалізують PTZ-індуковані зміни вмісту ЦОГ-1/2 (дигоксин, ймовірно, завдяки своїй стероїдній структурі проникає в ядро гліальних клітин і регулює експресію генів ферментів), а також окремих продуктів циклооксигеназного шляху в головному мозку, що, очевидно, є одним із можливих механізмів протисудомної дії препаратів. *In silico* встановлено неможливість занурення дигоксину в активні сайти ферментів ЦОГ-1 і ЦОГ-2, що є додатковим свідченням на користь ядерного впливу дигоксину на експресію генів зазначених ензимів у головному мозку.

Висновки. Дигоксин та вальпроат і особливо їх комбінація ефективно протидіють нейрозапаленню, забезпечуючи виразний вплив на циклооксигеназний шлях каскаду арахідонової кислоти в головному мозку, що, вочевидь, є одним із механізмів їх протисудомної активності.

ВИВЧЕННЯ ЗМІН ГЕМАТОЛОГІЧНИХ ПОКАЗНИКІВ У ЩУРІВ В УМОВАХ ЕКСПЕРИМЕНТАЛЬНОГО КОЛІТУ ПРИ ЛІКУВАННІ МОДИФІКОВАНИМИ ФІТОЕКСТРАКТАМИ SALVIA OFFICINALIS

Верховодова Ю. В.

Національний фармацевтичний університет, м. Харків, Україна
Кафедра клінічної фармакології ІПКСФ
Juliaverh73@gmail.com

Вступ. Дисбіотичні порушення, інфекційний та неінфекційний коліт є актуальною проблемою сьогодення, бо впливає на якість життя хворих і фармакокінетику ліків. Дані попередніх досліджень свідчать про наявність протизапального ефекта у модифікованих фітоекстрактів Salvia officinalis (S. officinalis) №2 та №4.

Мета. Вивчення змін гематологічних показників у щурів в умовах експериментального коліту при лікуванні модифікованими фітоекстрактами S. officinalis №2 та №4.

Матеріали та методи. Було досліджено ефект модифікованих фітоекстрактів S. Officinalis №2 та №4 на гематологічні показники на моделі інфекційного коліту у щурів **(**“Спосіб моделювання коліту з дисбіотичними порушеннями” патент на корисну модель № 140188 від 10.02.2020 р., патент на винахід №122640 від 11.12.2020 р.). Щури були розділені на три групи по 7 тварин: групи дослідження і контрольна (інтактні тварини). Препаратом порівняння був рифаксимін. Гематологічні показники вимірювали уніфікованими колориметричними методами з використанням наборів ТОВ НВП «Філісіт-Діагностика», Україна.

Результати. В експерименті на щурах в дослідних групах відбулися зміни в показниках крові: кількості лейкоцитів, еритроцитів, гемоглобіну, показника ШОЕ. Аналіз показників свідчить про наявність запалення у кишківнику, а саме значний лейкоцитоз, збільшення ШОЕ та статистично значуще підвищення кількості моноцитів, нейтрофілів, зниження кількості еритроцитів та гемоглобіну. В результаті лікування модифікованими фітоекстрактами S. officinalis на 15 добу експеримента показники лейкоцитарної формули нормалізувались. Фармакологічний ефект фітоекстрактів S. officinalis №2 та №4 на показники периферичної крові у дослідних тварин засвідчив статистично значуще збільшення значень гемоглобіну і кількості еритроцитів у порівнянні з контролем. Після лікування препаратом порівняння гематологічні показники крові поступово нормалізувались, але не досягали значень як у щурів з досліджуваними екстрактами.

Висновки. Встановлено статистично значуще зниження кількості лейкоцитів, ШОЕ, що свідчить про зменшення проявів запального процесу після лікування фітоекстрактами S. officinalis №2 та №4. Препарат порівняння рифаксимін також зменшував прояви запалення у показниках периферичної крові, але поступався за ефектом досліджуваним екстрактам.

ВИВЧЕННЯ АНТИОКСИДАНТНОЇ АКТИВНОСТІ ЕКСТРАКТІВ ЦИНІЇ ВИТОНЧЕНОЇ

Тулуб І. О.[1], Бурда Н. Є.[1], Фіра Л. С.[2]

[1]Національний фармацевтичний університет, м. Харків, Україна

[2]Тернопільський національний медичний університет імені І.Я. Горбачевського, м. Тернопіль, Україна

nadegdaburda@ukr.net

Вступ. Представлене дослідження є продовженням нашого комплексного фармакогностичного вивчення сировини цинії витонченої (*Zinnia elegans* Jacq., родина Asteraceae). Наразі нами для екстрактів, отриманих із сировини вищезазначеної рослини, досліджена антимікробна та протизапальна дія. Однак, для більш детального та поглибленого вивчення цинії витонченої доцільно вивчати інші види фармакологічної активності. З огляду на хімічний склад сировини цинії витонченої, який представлений в основному фенольними сполуками, зокрема флавоноїдами та фенольними кислотами, доцільно провести вивчення антиоксидантної активності.

Мета. Метою роботи було вивчення антиоксидантної активності густих екстрактів, отриманих з трави та квіток цинії витонченої.

Матеріали та методи. Для проведення фармакологічних досліджень одержували густі екстракти із трави та квіток цинії витонченої: співвідношення сировина: екстрагент – 1 : 5; екстрагент – 70 % етанол; настоювання – протягом 5 діб. Після цього екстракти концентрували.

Вивчення антиоксидантної активності густих екстрактів із трави та квіток цинії витонченої проводили на моделі гострого гепатиту. Моделювання гострого гепатиту проводити шляхом введення парацетамолу інтрагастрально у дозі 1250 мг/кг 1 раз на добу протягом 2 діб у вигляді суспензії в 2 % розчині крохмального гелю.

Експеримент проводили на 78 нелінійних білих щурах-самцях вагою 180-210 г. Досліджувані зразки екстрактів вводили дослідним щурам внутрішньошлунково за 2 год до введення токсиканта та щоденно після ураження у дозі 100 мг/кг (для густого екстракту з трави) та 150 мг/кг (для густого екстракту з квіток) маси тварини.

Як препарат порівняння використовували Карсил (діюча речовина – силімарин, виробник – АТ «Sopharma») у дозі 100 мг/кг.

Результати. Отримані результати дозволяють стверджувати, що екстрактам із трави та квіток цинії витонченої притаманні антиоксидантні властивості, які підтверджуються пригніченням окиснювального стресу за умов токсичного ураження печінки.

Висновки. Отже, результати проведеного експерименту вказують, що густі екстракти з трави та квіток цинії витонченої можуть бути використані як АФІ при розробці нових лікарських засобів вітчизняного виробництва з антиоксидантною активністю.

ДОСЛІДЖЕННЯ ПРОТИМІКРОБНОЇ АКТИВНОСТІ РОТИКІВ САДОВИХ ТРАВИ ЕКСТРАКТУ ГУСТОГО

Ільїна С. К., Журавель І. О.
Національний фармацевтичний університет, м. Харків, Україна
Кафедра фармакогнозії та нутриціології
andrianovasofia@gmail.com

Вступ. Проблема виникнення множинної лікарської стійкості бактерій та грибів зменшує ефективність лікування інфекцій і потребує пошуку нових протимікробних засобів, зокрема природного походження. Рослини багаті на біологічно активні речовини (флавоноїди та фенолокислоти) стають перспективним джерелом лікарських засобів.

Одним з таких джерел є ротики садові (*Antirrhinum majus* L.), що належать до родини Подорожникових (*Plantaginaceae*). Рослина має багатий хімічний склад, зокрема містить флавоноїди, гідроксикоричні та амінокислоти, а також мінеральні речовини: калій, кальцій, магній, фосфор, натрій та силіцій [1,2].

Вченими Малазійського наукового університету було експериментально доведено ранозагоювальну та протизапальну дію сировини [3]. А також досліджено протимікробну активність екстрактів ротиків садових до *Bacillus subtilis, Escherichia coli* та *Candida albicans* [4].

Більш детальне дослідження екстракту густого ротиків садових для оцінки його протимікробної активності є перспективним і може сприяти розробці нових фітопрепаратів та доповненню схем лікування інфекційних захворювань.

Мета. Дослідження протимікробної активності ротиків садових трави екстракту густого.

Матеріали та методи. Об’єктом дослідження був одержаний нами ротиків садових трави екстракт густий.

Визначення протимікробної активності екстракту проводили методом дифузії в агар на двох шарах щільного поживного середовища, розлитого в чашки Петрі. Нижній шар – «голодне» незасіяне середовище (агар-агар, вода, солі). Верхній шар складався з поживного агаризованого середовища, розплавленого та охолодженого до температури (40,0 ± 0,5) °С, в яке вносили відповідний стандарт добової тест-культури мікроорганізма. Об’єм досліджуваного екстракту, що поміщали в лунку, становив 0,3 мл. Діаметри зон затримки росту мікроорганізмів заміряли за допомогою мірної лінійки з точністю вимірювання 1,0 мм.

Приготування мікробної суспензії штамів проводили з використанням приладу Densi-La-Meter (виробництво PLIVA-Lachema, Чехія; довжина хвилі 540 нм). Синхронізацію культур штамів проводили з використанням низької температури (4 °С). Мікробне навантаження складало 107 мікробних клітин на 1 мл середовища і встановлювалося за стандартом McFarland. У дослідах використовували 18-24 годинну культуру штамів мікроорганізмів. Для культивування використовували агар Мюллера-Хинтона (Виробництво - Індія «HIMedia

Laboratorles Pvt. Ltd India»). Для культивування *Candida albicans* використовували агар Сабуро (Виробництво - Індія, «HIMedia Laboratorles Pvt. Ltd India»).

Результати та обговорення. У дослідженні було виявлено, що ротиків садових трави екстракт густий демонстрував помірну протимікробну активність проти референтних та клінічних бактеріальних та грибкових штамів. Зокрема, чутливими до дії досліджуваного екстракту були штами грам-позитивних бактерій *Staphylococcus aureus, Bacillus subtilis, Streptococcus pneumoniae, Streptococcus pyogenes*, а також грам-негативних бактерій: *Escherichia coli, Proteus vulgaris, Klebsiella pneumoniae, Enterobacter cloaceae* і *Pseudomonas aeruginosa*. Окрім цього, спостерігалася незначна інгібувальна дія щодо *Candida albicans*.

Протимікробна активність зменшувалася зі збільшенням розведення екстракту, а клінічні ізоляти *Enterococcus faecalis, Staphylococcus aureus ma Acinetobacter baumannii*, виявилися менш чутливими до розведених екстрактів.

Досліджений екстракт не виявив інгібувальної дії щодо кількох патогенних мікроорганізмів, зокрема: грам-негативних бактерій *Acinetobacter baumannii* та клінічного ізолята *Pseudomonas aeruginosa*.

Висновок. Визначено протимікробну активність ротиків садових трави екстракту густого, виявлено чутливість до дії цього екстракту деяких розповсюджених мікроорганізмів. Одержані результати будуть використані для розробки лікарських засобів на основі ротиків садових трави екстракту густого та розширення переліку номенклатури протимікробних засобів.

Список використаних джерел інформації:

1. Kumar G. A review of the chemical constituents and pharmacological activities of Antirrhinum majus (Snapdragon). IP Int J Compr Adv Pharmacol 2022;7(2):72-76. DOI: 10.18231/j.ijcaap.2022.013
2. Ільїна С.К., Журавель І.О. Дослідження мінерального складу та визначення деяких показників якості квіток і трави ротиків садових (Antirrhinum majus L.) Annals of Mechnikov’s Institute. 2023. № 4. С. 139-143. DOI: 10.5281/zenodo.10255409
3. Saqallah, FG (Saqallah, Fadi G.); Hamed, WM (Hamed, Wafaa M.); Talib, WH (Talib, Wamidh H.) In Vivo Evaluation of Antirrhinum majus' Wound-Healing Activity. Scientia pharmaceutica. 2018. №45. DOI: 10.3390/scipharm86040045
4. Saqallah, FG (Saqallah, Fadi G.), Hamed, WM (Hamed, Wafaa M.), Talib, WH (Talib, Wamidh H.), et al. Antimicrobial activity and molecular docking screening of bioactive components of Antirrhinum majus (snapdragon) aerial parts. Heliyon. 2022. DOI: 10.1016/j.heliyon.2022.e10391

ВПЛИВ КАРБАЦЕТАМУ НА ГЛУТАТІОНОВИЙ ЛАНЦЮГ ГІПОКАМПА СТАРИХ ЩУРІВ З ЕКСПЕРИМЕНТАЛЬНОЮ ХВОРОБОЮ АЛЬЦГЕЙМЕРА

Дрезналь Є. П., Кметь Т. І.
Буковинський державний медичний університет,
м. Чернівці, Україна
Кафедра гігієни та екології
dreznal555@gmail.com

Вступ. Хвороба Альцгеймера – це незворотне, прогресуюче захворювання мозку, яке повільно руйнує пам’ять, навички мислення і здатність виконувати найпростіші завдання. Дана патологія є найпоширенішою формою деменції, яка є однією з найважливіших медико-соціальних проблем сьогодення. Сучасні дослідницькі програми спрямовані на розробку стратегій ефективного прогнозування, профілактики та лікування розладів функцій центральної нервової системи.

Найбільш розповсюдженою ознакою для більшості нейродегенеративних розладів, зокрема хвороби Альцгеймера, є надмірне утворення реактивних форм кисню внаслідок окисного стресу, що спричиняє пошкодження та втрату нейрональних клітин. Водночас відомо, що головний мозок є найбільш чутливим до окисного пошкодження органом. І суттєву роль у реалізації антирадикального захисту нейронів відіграє глутатіонова система, яка сприяє відновленню оптимального рівня пероксидних сполук і збереженню про-антиоксидантного балансу. Узгоджена дія всіх її компонентів (відновленого глутатіону, глутатіонпероксидази, глутатіонредуктази, сульфгідрильних груп) сприяє відновленню оптимального рівня пероксидних сполук. Більш того, глюкозо-6-фосфатдегідрогеназа, яка займає ключове місце в обміні глутатіону, є джерелом пентоз, необхідних для синтезу і репарації ДНК. Тобто, ступінь пошкодження глутатіонової системи тісно пов'язаний із цитотоксичними ефектами. Водночас порушенням у системі глутатіону належить лідируюча роль у реалізації механізмів нейродегенерації, викликаної оксидативним стресом.

Мета. Метою роботи було вивчити вплив модулятора ГАМК-рецепторів карбацетаму на глутатіоновий ланцюг антиоксидантної системи гіпокампа старих щурів з експериментальною хворобою Альцгеймера.

Матеріали та методи. Експерименти проводили на нелінійних лабораторних білих щурах самцях. Модель створювали внутрішньоочеревинним введенням протягом 27 днів скополаміну гідрохлориду (Sigma, США) дозою 1 мг/кг. На 28 добу щурів із хворобою Альцгеймера сліпим методом розподілили на дві групи: І – із введенням карбацетаму внутрішньоочеревинно дозою 5 мг/кг; ІІ – із введенням тільки фізіологічного розчину. Контрольним щурам із 28 дня вводили фізіологічний розчин. Евтаназію здійснювали під легким хлороформним наркозом. На холоді виймали головний мозок, ретельно промивали охолодженим 0,9 % розчином NaCl і за стереотаксичним атласом виділяли гі-

покамп. Для оцінки стану антиоксидантної системи гіпокампа визначали вміст глутатіону відновленого (G-SH), сульфгідрильних (SH-) груп та активність глутатіон-редуктази (ГР), глутатіон-пероксидази (ГП), глюкозо-6-фосфатдегідрогенази (Г-6-ФДГ). Кількість протеїну в пробах визначали за методом Лоурі. Статистичну обробку результатів проводили за допомогою t-критерію Стьюдента. Достатнім рівнем достовірності вважали $p \leq 0,05$.

Результати. У нашому дослідженні встановлено, що у порівнянні з контрольною групою, вміст G-SH у щурів із хворобою Альцгеймера знижувався на 43,4 %. Водночас виявлено зниження активності ферменту НАДФН-залежної ГР, що бере участь у процесі антиоксидантного захисту – на 45,4 %. При цьому активність ГП, була меншою, ніж у контрольної групи на 31,4 %. Вміст SH-груп, які входять до складу глутатіону і забезпечують біохімічні реакції метаболізму та збереження функцій мембран, також знижувався на 32,6 %. Встановлено зниження активності Г-6-ФДГ на 24,6 % у гіпокампі старих щурів з експериментальною хворобою Альцгеймера.

Подальший аналіз результатів показав, що після введення карбацетаму щурам із модельною патологією підвищувались показники антиоксидантного захисту в гіпокампі. Порівнюючи дані групи з корекцією з показниками у щурів, яким не вводили карбацетам, встановлено збільшення вмісту G-SH у в 1,5 раза та SH-груп в 1,2 раза. Позитивний вплив карбацетаму також характеризувався зростанням активності ГР в 1,6 раза та Г-6-ФДГ в 1,4 раза.

Результатом вказаних змін є зменшення утворення активних форм кисню, підвищення активності ферментів антиоксидантного захисту. Ймовірним механізмом дії карбацетаму є вплив на ГАМК-рецептори, які локалізуються в стінках мозкових судин. Це відіграє суттєву роль у регуляції мозкового кровообігу: розширюються мозкові судини, підвищується об’ємний кровотік та вмісту кисню в клітинах мозку, що вказує на протекторні властивості карбацетаму стосовно зменшення ознак денудації судин і зниженню прооксидантної активності в гіпокампі. Таким чином, наведені дані свідчать про протективний вплив ендогенного модулятора ГАМК-ергічних рецепторів карбацетаму за умов розвитку центральної нейродегенерації, спричиненої хворобою Альцгеймера.

Висновок. Покращення стану глутатіонового ланцюга антиоксидантного захисту в гіпокампі старих щурів, вказує на здатність карбацетаму пригнічувати патогенетичні ланки нейродегенеративних процесів, зумовлених холінонегативними впливами у центральній нервовій системі**.**

СТАБІЛЬНІСТЬ ПРОТИМІКРОБНОГО ЕФЕКТУ ГЕЛЮ НА ОСНОВІ КОМПОЗИЦІЇ АЦИЛЬОВАНОГО НІЗИНУ З ДИКЛОФЕНАКОМ НАТРІЯ ЩОДО КЛІНІЧНИХ ШТАМІВ МІКРООРГАНІЗМІВ

Андреєва І. Д.[1], Осолодченко Т. П.[1], Батрак О. А.[2], Рябова І. С.[1]

ДУ «Інститут мікробіології та імунології ім. І. І. Мечникова Національної академії медичних наук України», м. Харків, Україна

[1]Лабораторія біохімії та біотехнології

[2]Лабораторія молекулярної імунофармакології

idandreyeva@gmail.com

Вступ. Одним з перспективних методів боротьби з патогенними бактеріями є використання бактеріоцинів. Поєднання бактеріоцинів з інгібіторами резистентності, які не мають прямої антимікробної дії, може бути перспективним напрямком.

Мета. Визначення стабільності у часі протимікробного ефекту гелю на основі композиції ацильованого нізину з диклофенаком натрія стосовно клінічних штамів мікроорганізмів.

Матеріали і методи. Модифікований нізин отримано за допомогою ацилювання з оцтовим ангідридом. У якості допоміжної речовини застосовано 0,3 % карбоксиметилцеллюлозу. Стабільність протимікробного ефекту гелю на основі композиції 1,0 % ацильованого нізину і діклофенака натрія визначалася шляхом їх витримування у нормальних умовах в реальному часі протягом 60 діб. Для мікробіологічного дослідження досліджуваних речовин використані клінічні штами грампозитивних і грамнегативних бактерій, а саме: *Staphylococcus aureus* 16, *Staphylococcus epidernides* 18, *Streptococcus pneumonia* 14, *Streptococcus pyogenes* 2432, *Enterococcus faecalis* 27, *Escherichia coli* 29, *Proteus vulgaris* 35, *Proteus mirabilis* 49. Культури мікроорганізмів було одержано з колекції лабораторії біохімії та біотехнології ДУ «ІМІ НАМН». Визначення антимікробної активності препаратів проведено дифузійним методом «колодязів» з вимірюванням діаметрів зон затримки росту мікроорганізмів.

Результати. При зберіганні гелю на основі композиції 1,0 % ацильованого нізину і 1,0 % диклофенака натрія в нормальних умовах впродовж 60 діб його протимікробний ефект щодо усіх досліджених клінічних штамів мікроорганізмів зберігався стабільно помірним. Діаметри зон затримки їх росту на початку експерименту коливалися у межах від (21,7±0,5) мм до (18,7±0,5) мм, на 15 добу – від (21,0±0,0) мм до (18,3±0,5) мм, на 30 добу – від (20,0±0,0) мм до (18,0±0,0) мм і на 60 добу – від (19,7±0,5) мм до (18,0±0,0) мм.

Висновки. За результатами проведених досліджень доведено стабільність протимікробного ефекту гелю на основі композиції ацильованого нізину з диклофенаком натрія щодо клінічних штамів мікроорганізмів та перспективність досліджень з метою створення на його основі протимікробних засобів.

ВИЗНАЧЕННЯ СТАБІЛЬНОСТІ ПРОТИМІКРОБНОГО ЕФЕКТУ ГЕЛЮ НА ОСНОВІ КОМПОЗИЦІЇ СУКЦИЛЬОВАНОГО НІЗИНУ З ДИКЛОФЕНАКОМ НАТРІЯ ЩОДО КЛІНІЧНИХ ШТАМІВ МІКРООРГАНІЗМІВ

Осолодченко Т. П.[1], Андреєва І. Д.[1], Мартинов А. В.[2], Завада Н. П.[1]

ДУ «Інститут мікробіології та імунології ім. І. І. Мечникова Національної академії медичних наук України», м. Харків, Україна

[1]Лабораторія біохімії та біотехнології

[2]Лабораторія молекулярної імунофармакології

idandreyeva@gmail.com

Вступ. Одним з перспективних методів боротьби з патогенними бактеріями є використання бактеріоцинів у поєднанні з хелперними речовинами. Пошук хелперних речовин серед вже відомих та гарно вивчених субстанцій вважають вельми перспективним напрямком боротьби з резистентністю мікроорганізмів.

Мета. Оцінка стабільності у часі протимікробного ефекту гелю на основі композиції сукцильованого нізину з диклофенаком натрія стосовно клінічних штамів мікроорганізмів.

Матеріали і методи. Модифікований нізин отримано за допомогою сукцилювання з бурштиновим ангідридом. У якості допоміжної речовини застосовано 0,3 % карбоксиметилцеллюлозу. Досліджено стабільність протимікробного ефекту гелю на основі композиції 1,0 % сукцильованого нізину і діклофенака натрія шляхом їх витримування у нормальних умовах в реальному часі протягом 60 діб. Для мікробіологічного дослідження досліджуваних речовин використані клінічні штами грампозитивних і грамнегативних бактерій, а саме: *S. aureus* 16, *S. epidernides* 18, *S. pneumonia* 14, *S. pyogenes* 2432, *E. faecalis* 27, *E.coli* 29, *P. vulgaris* 35, *P. mirabilis* 49,які було одержано з лабораторії медичної мікробіології з Музеєм мікроорганізмів ДУ «ІМІ НАМН». Визначення антимікробної активності препаратів проведено дифузійним методом «колодязів» з вимірюванням діаметрів зон затримки росту мікроорганізмів. Наявність зони затримки росту до 10 мм розцінювалася як відсутність чутливості, 10–15 мм – як низька, 15–25 мм – як помірна і перевищення 25 мм – як висока.

Результати. Після зберігання впродовж 60 діб протимікробний ефект гелю на основі композиції 1,0 % сукцильованого нізину і 1,0 % диклофенака натрія щодо усіх досліджених зберігався стабільно помірним. Діаметр зон затримки росту клиничних штамів мікроорганізмів на початку експерименту коливався у межах від (20,7±0,5) мм до (18,0±0,0) мм, на 15 добу – від (20,0±0,0) мм до (17,3±0,5) мм, на 30 добу – від (18,7±0,5) мм до (17,0±0,0) мм і на 60 добу – від (17,7±0,5) мм до (16,0±0,0) мм.

Висновки. Результати проведених досліджень свідчать про стабільність в часі протимікробного ефекту гелю на основі композиції сукцильованого нізину з диклофенаком натрія.

ДОСЛІДЖЕННЯ ГОСТРОЇ ТОКСИЧНОСТІ НОВИХ ТІОФЕН ПОХІДНИХ 1,2,4-ТРИАЗОЛІВ

Хільковець А. В., Білай І. М.
Запорізький державний медико-фармацевтичний університет,
м. Запоріжжя, Україна
Кафедра клінічної фармації, фармакотерапії, фармакогнозії
та фармацевтичної хімії
nastia010792@ukr.net

Вступ. Пошук біологічно активних речовин, розробка та створення нових ефективних і безпечних лікарських засобів залишаються однією з основних проблем фармацевтичної науки. Важливими об'єктами для досліджень у цьому контексті є похідні 1,2,4-триазолу. На сьогодні вже розроблено лікарські засоби на основі цих похідних, які використовуються як у медицині, так і в ветеринарії, але наукові дослідження тривають. Ми отримали кілька нових тіофенвмісних 1,2,4-триазолів, будову яких підтверджено за допомогою фізико-хімічного аналізу, і вибрали одну з них для подальшого дослідження. Як відомо, обов'язковою вимогою до новосинтезованих сполук є проведення доклінічних досліджень їх безпечності. Вивчення гострої токсичності необхідне для первинної оцінки токсичності та вивчення симптомів отруєння при одноразовому введенні в організм.

Мета. Метою нашого дослідження було вивчення гострої токсичності 2-((4-феніл-5-(тіофен-3-ілметил)-1,2,4-триазол-3-іл)тіо)ацетату натрію.

Матеріали та методи. Дослідження гострої токсичності проводили за методом *in vivo*, згідно рекомендаціям Стефанова О.В. з подальшим розрахунком параметрів гострої токсичності по методу Кербера в модифікації Лойта А.О і Савченкова М.Ф. Для визначення рівнів доз було проведено комп’ютерне прогнозування гострої токсичності за допомогою сервісу ProTox-II. Комп’ютерний прогноз гострої токсичності був здійснений за структурною формулою у веб-версії цього сервісу

Результати. В експерименті використовували нелінійних щурів (36 осіб), яких було поділено на шість груп (інтакт, рівні доз – 200 мг/кг, 500 мг/кг, 700 мг/кг, 1500 мг/кг та 3000 мг/кг), експеримент тривав 14 діб. Після введення спостерігалося зниження рухової активності, сонливість, міоз зіниць та спрага. У п’ятій групі, де була максимальна доза, усі щури загинули протягом двох годин після введення сполуки, при цьому спостерігалися судоми. У четвертій групі загинуло п’ять щурів, у третій — два, а в перших двох групах усі тварини вижили. Подальше спостереження показало, що тварини з перших двох груп поводилися нормально.

Висновки. Згідно з результатами, досліджувана сполука належить до V класу токсичності (практично нетоксична), а її LD50 становить 1125 мг/кг. Ці дані підтверджують перспективність подальшого вивчення цієї сполуки.

АКТИВНІСТЬ ЛАКТАТДЕГІДРОГЕНАЗИ В ПЕЧІНЦІ ЩУРІВ З АЛОКСАНОВИМ ДІАБЕТОМ ЗА УМОВ УВЕДЕННЯ МЕЛАТОНІНУ

Кушнір О. Ю., Кушнір О. О.

Буковинський державний медичний університет
Кафедра біоорганічної і біологічної хімії та клінічної біохімії
м. Чернівці, Україна
kushnir@bsmu.edu.ua

Вступ. Цукровий діабет – патологія, яка пов'язана з порушенням всіх видів обміну речовин. Відомо, що недостатній вплив інсуліну призводить до зниження активності піруватдегідрогеназного комплексу. Одним із проявів захворювання є лактат ацидоз. Це призводить до напруження буферних систем крові, зокрема бікарбонатної буферної системи. Мелатонін – адаптоген, який володіє антиоксидантними властивостями і має антиглікемізувальну дію. Яким чином впливає мелатонін на активність лактатдегідрогенази (ЛДГ) в печінці щурів з алоксановим діабетом досі мало вивчено.

Мета. Метою роботи було встановити зміни активності ЛДГ в печінці щурів з алоксановим діабетом під впливом мелатоніну.

Матеріали та методи. Досліди проведені згідно із встановленими нормами поводження з експериментальними тваринами. У роботі використано 50 білих безпородних щурів самців масою 150 грам. Діабет викликали одноразовим введення алоксану моногідрату (170 мг на кг маси тіла тварини). Щурів поділили на три групи: інтактні, діабетичні, діабетичні з мелатоніном. Мелатонін уводили щоденно вранці (5 мг на кг маси тіла) впродовж 2 тижнів. Тварин декапітували на 19 добу від початку експерименту. Активність ЛДГ печінки, рівень глюкози натще та pH крові визначали за стандартними методиками.

Результати. Введення алоксану моногідрату призвело до загибелі 50% тварин. Решта сформували групу діабетичних тварин з рівнем глюкози натще на 125 % вищим ніж у інтактних тварин. Активність ЛДГ в діабетичних тварин вдвічі перевищувала показники інтактної групи. В даній групі щурів pH виявилося незначно зміщене в кислий бік у порівнянні з інтактними тваринами. Можливо має місце включення в роботу бікарбонатної буферної системи, що взаємодіє з надлишком лактату, утворюючи натрієву сіль. Уведення мелатоніну призвело до зниження рівня глюкози натще на 42%, нормалізації активності ЛДГ та pH крові алоксандіабетичних щурів порівняно з показниками інтактних тварин. Відомо, що мелатонін активує транспортери глюкози та стимулює процеси її аеробного перетворення.

Висновок. Мелатонін призводить до нормалізування активності ЛДГ в печінці щурів з алоксановим діабетом можливо за рахунок відновлення процесів поглинання та використання глюкози аеробним шляхом, що зменшує рівень лактату та ризик розвитку ацидозу.

ДОСЛІДЖЕННЯ СПАЗМОЛІТИЧНИХ ТА ПРОТИЗАПАЛЬНИХ ВЛАСТИВОСТЕЙ 8-АМІНО-1-БЕНЗИЛ-7-ЕТИЛ-3-МЕТИЛКСАНТИНІВ

Іванченко Д. Г., Крісанова Н. В.

Запорізький державний медико-фармацевтичний університет,
м. Запоріжжя, Україна
Кафедра біологічної хімії
ivanchenko230181@gmail.com

Вступ. Одним із значущих викликів у клінічній практиці є необхідність оперативної та ефективної корекції спастичних розладів, які часто супроводжують різноманітні гастроентерологічні патології. Особливо актуальною ця проблема постає при захворюваннях шлунково-кишкового тракту (ШКТ), а також при патологічних станах жовчного міхура та жовчовивідних шляхів. Спастичні прояви у цих випадках не лише погіршують якість життя пацієнтів, але й можуть ускладнювати діагностику та лікування основного захворювання. Таким чином, виходячи із вищенаведеного, можна зробити висновок, що проблема розробки оригінальних вітчизняних препаратів спазмолітичної та протизапальної дії є перспективною та актуальною.

Мета. Метою даної роботи є пошук перспективних спазмолітичних та протизапальних засобів серед 8-аміно-1-бензил-7-етил-3-метилксантинів.

Матеріали та методи. Вивчення протизапальної (антиексудативної) дії проводили згідно з методичними рекомендаціями ДЕЦ МОЗ України. Дослідження проведене на самках білих безпородних мишей масою тіла 18,0-20,0 г.

Для скринінгових досліджень застосовували модифікацію карагенінового набряку у мишей. Запалення викликали субплантарним введенням 1% розчину карагеніну виробництва Sigma (США) у дозі 0,02 мл/тварину. Контралатеральна лапа служила контролем.

Синтезовані сполуки вводили за 1 годину до введення флогогену. Через 1 год лапи відрізали на рівні надступаковогомілкового суглобу. Лапи зважували на електронних вагах. Визначали індекс реакції.

Ефективність застосування тест-зразка оцінювали за пригніченням розвитку набряку лапи у порівнянні з тваринами групи позитивного контролю та дією референс-зразка (диклофенак натрію).

Результати. Вивчення гострої токсичності *in vivo* показало, що синтезовані сполуки належать до IV класу токсичності. Серед отриманих сполук за показниками спазмолітичної та протизапальної активності виявлені сполуки, які не поступаються, а в деяких випадках активніші за еталон порівняння.

Висновки. Встановлено, що досліджувані сполуки відносяться до IV класу токсичності. Виявлені певні закономірності впливу замісників на протизапальну та спазмолітичну активність. Для остаточних висновків про закономірності в ряді «хімічна структура – біологічна дія» необхідно розширити гомологічні ряди ксантинових похідних.

ПОШУК ГІПОЛІПІДЕМІЧНИХ ЗАСОБІВ СЕРЕД 8-АМІНО-7-(2-ГІДРОКСИ-2-ФЕНІЛЕТИЛ)-3-МЕТИЛКСАНТИНІВ

Іванченко Д. Г., Рудько Н. П.

Запорізький державний медико-фармацевтичний університет,
м. Запоріжжя, Україна
Кафедра біологічної хімії
ivanchenko230181@gmail.com

Вступ. Атеросклероз, згідно з визначенням ВООЗ, характеризується комплексними змінами внутрішньої оболонки артерій, що включають фокальне накопичення ліпідів, складних вуглеводних сполук, елементів крові та циркулюючих речовин, формування сполучної тканини та кальцифікацію. Ця патологія посідає провідне місце серед серцево-судинних захворювань і набула епідемічних масштабів, особливо в урбанізованих регіонах розвинених країн.

Прогресування атеросклерозу призводить до оклюзії артерій, що спричиняє ряд серйозних ускладнень, зокрема, захворювання коронарних і церебральних судин. Етіологія атеросклерозу мультифакторна, з добре встановленими факторами ризику, такими як куріння, гіпертензія, цукровий діабет та дисліпідемія. Незважаючи на досягнення в корекції ліпідного профілю, атеросклероз залишається основною причиною смертності.

Патофізіологія атеросклерозу включає складні механізми: акумуляцію агрегатів ліпопротеїнів у субендотеліальному просторі, запальні реакції ендотеліальних клітин, адгезію та інфільтрацію моноцитів, трансформацію макрофагів у пінисті клітини та формування атеросклеротичних бляшок. Зниження рівня холестерину в плазмі крові за допомогою дієтотерапії та фармакологічних засобів сприяє уповільненню прогресування та регресії атеросклеротичних уражень.

Дослідження показали, що похідні ксантину та аденіну демонструють гіполіпідемічну активність, що відкриває перспективи для розробки нових терапевтичних агентів.

З огляду на вищезазначене, розробка інноваційних вітчизняних препаратів з гіполіпідемічною дією є актуальним та перспективним напрямком фармацевтичних досліджень, спрямованим на вдосконалення профілактики та лікування атеросклерозу.

Мета. Метою даної роботи є пошук малотоксичних та високоефективних гіполіпідемічних засобів серед 8-аміно-7-(2-гідрокси-2-фенілетил)-3-метилксантинів.

Матеріали та методи. Гостра токсичність синтезованих сполук була вивчена за методом Кербера у дослідах на білих мишах.

Гіпохолестеролемічну активність вивчали при експериментальній гіперліпідемії, яку створювали за короткотривалою моделлю Jowsufszai-Siddigi (пероральне зондове введення дорослим щурам-самцям лінії Вістар вагою 220-280 г холестеролу в добовій дозі 40 мг/кг та фактору порушення ліпідного метабо-

лізму й посилення всмоктування стероїдів в кишечнику – 0,125% олійного розчину ергокальциферолу в добовій дозі 8 мл/кг). Водну суспензію ксантинів в дозі 50 мг/кг додавали per os через одну год після введення гіперліпідогенної суміші протягом 5-ти днів. На шостий день після наркотизації етиловим етером здійснювали забір крові з біфуркації аорти. Формували також групи тварин: інтактні (введення дистильованої води 30 мл/кг), контрольні (введення суміші холестерол-ергокальциферол без додавання препаратів) та щури, яким вводили еталонні препарати (аторвастатин, фенофібрат). Кров центрифугували при 1500 об/хв, виділяли сироватку. В сироватці крові щурів визначали вміст загального холестеролу за методом Ілька. В якості еталону порівняння використовували аторвастатин в дозі 10 мг/кг та фенофібрат в дозі 60 мг/кг.

Результати. Експериментальне дослідження гострої токсичності in vivo продемонструвало, що синтезовані сполуки відносяться до IV класу токсичності за загальноприйнятою класифікацією. Це свідчить про їх відносно низьку токсичність та потенційну безпечність для подальшого вивчення та застосування.

Аналіз гіпохолестеролемічної активності отриманих сполук виявив ряд перспективних кандидатів. Зокрема, деякі з синтезованих речовин продемонстрували рівень активності, що не поступається еталонному препарату порівняння, а в окремих випадках навіть перевищує його ефективність. Ці результати вказують на потенціал досліджуваних сполук як основи для розробки нових гіпохолестеролемічних засобів.

Отримані дані створюють підґрунтя для подальших досліджень, спрямованих на детальне вивчення механізмів дії найбільш перспективних сполук, оцінку їх фармакокінетичних властивостей та можливості їх застосування в терапевтичній практиці для корекції порушень ліпідного обміну.

Висновки. Результати токсикологічних досліджень продемонстрували, що синтезовані сполуки належать до IV класу токсичності.

У ході експериментальних досліджень були виявлені певні закономірності щодо впливу структурних модифікацій, зокрема введення різних замісників, на гіпохолестеролемічну активність досліджуваних сполук.

Однак, для формулювання більш ґрунтовних висновків щодо залежності "структура-активність" та встановлення чітких закономірностей у ряді ксантинових похідних, необхідне проведення додаткових досліджень з розширенням гомологічних рядів досліджуваних сполук. Це дозволить отримати більш повну картину впливу структурних модифікацій на фармакологічні властивості та може сприяти ідентифікації оптимальних структур для подальшої розробки потенційних лікарських засобів з покращеними гіполіпідемічними характеристиками.

Таким чином, отримані результати створюють підґрунтя для подальших цілеспрямованих досліджень у напрямку оптимізації структури ксантинових похідних з метою розробки нових ефективних та безпечних фармакологічних агентів.

ВИВЧЕННЯ ПАРАМЕТРІВ ГОСТРОЇ ТОКСИЧНОСТІ РОСЛИННОЇ СУБСТАНЦІЇ КАЛИНИ ЗВИЧАЙНОЇ ПЛОДІВ ЕКСТРАКТУ РІДКОГО

Шпичак О. С.[1], Міщенко О. Я.[2]

Інститут підвищення кваліфікації спеціалістів фармації
Національного фармацевтичного університету, м. Харків, Україна
[1]Кафедра фармацевтичної технології, стандартизації та сертифікації ліків
phtaqa@nuph.edu.ua
[2]Кафедра клінічної фармакології
clinpharmacol2023@gmail.com

Вступ. Обґрунтування та встановлення очікуваних фармакологічних властивостей одержаного нами з калини звичайної плодів рослинної субстанції у вигляді екстракту рідкого – КЗПЕР є передумовою для перспективи щодо подальшого його застосування у ЛЗ, що розробляються на її основі для застосування в медичній практиці. Оскільки причиною кардіоміопатій та ішемічної хвороби серця є фактори, що нерідко призводять до підвищення ризику розвитку серцевих патологій, наявність мембраностабілізувальної та антиоксидантної дії КЗПЕР відкриває реальні можливості щодо розширення спектру використання розробленого на його основі ЛЗ у формі таблеток під умовною назвою «Вібурнікор», а впровадження його в промисловість та терапевтичну практику має значно вплинути на поліпшення якості надання кваліфікованої медичної допомоги хворим на серцево-судинні захворювання.

Мета. Дослідити параметри гострої токсичності рослинної субстанції калини звичайної плодів екстракту рідкого (*Viburni opuli fructi extracti soluti),* що входить до складу таблеток «Вібурнікор» кардіопротекторної та мембраностабілізувальної дії.

Матеріали та методи. В роботі були використані сучасні методи біологічних досліджень. Експериментальні дані оброблено методами варіаційної статистики та непараметричних методів аналізу. Для отримання статистичних висновків використовували стандартний пакет програм STATISTICA (версія 6).

Результати. З метою відтворення клініки гострого отруєння та визначення параметрів токсичного впливу, дослідження гострої токсичності КЗПЕР проводили на статевозрілих білих безпородних щурах обох статей (самцях, самках) масою 160–180 г при внутрішньошлунковому введенні.

Відповідно до методичних рекомендацій максимально допустимий об’єм при внутрішньошлунковому введенні – 20 мл/кг, що перевищує дозу VI класу токсичності, яка складає 15 мл/кг. У зв’язку з цим, визначення гострої токсичної дії субстанції КЗПЕР проводили починаючи з дози 20 мл/кг. У разі не встановлення летальності КЗПЕР у дозі 20 мл/кг, подальші дослідження можна вважати недоцільними.

Як показали проведені дослідження, одноразове введення КЗПЕР у максимальному об’ємі 20 мл/кг (2 мл/100 г) не виявляло випадків летальності тварин. Також не відзначалося значущих порушень основних інтегральних показ-

ників загального стану і поведінки тварин. Щоденні спостереження за щурами експериментальних груп свідчать про те, що вони були активними, охайними, мали задовільний апетит, нормально реагували на звукові і світлові подразники, процеси сечовиділення і дефекацій були у нормі, порушення дихання та судом не спостерігали. Впродовж всього терміну спостереження тварини дослідних груп не відрізнялися від контрольних тварин за апетитом.

Дослідження маси тіла щурів показало, що приріст маси у тварин був позитивний в усіх дослідних групах і не виходив за межі значень групи негативного контролю. По закінченні спостереження (14 доба), тварин піддали евтаназії для наступного макроскопічного огляду внутрішніх органів та визначали абсолютну масу внутрішніх органів для розрахунку їх відносної маси.

При макроскопічному розтині розташування органів середостіння відповідає нормальному: тимус варіює за розміром, конусоподібної форми; серце звичайної конфігурації та розміру, з типовим розташуванням коронарних артерій та вен; легені блідо-рожеві, займають усю плевральну порожнину, еластичні, без спайок між листками плеври. Розташування органів очеревинної порожнини анатомічно правильне: очеревина прозора, гладенька, без стороннього вмісту; печінка червоно-коричневатого кольору, усі частки вільно розділяються, капсула не напружена, краї часток не закруглені, поверхня гладка, без вузликових утворювань. Підшлункова залоза блідо-рожево-жовтуватого кольору, без ознак крововиливів, склерозу та жирових некрозів. Селезінка пружна, повнокровна, червоно-вишневого кольору. Нирки і наднирники нормального вигляду, розміру. Слизова оболонка залозистого відділу шлунку з характерним рельєфом складок, без геморагій, набряку, ерозивних ушкоджень. Яєчки, придатки яєчок, передміхурова залоза та сім’яні пухирці у самців щурів та роги матки і яєчники у самок без видимої патології. Розрахунок та наступний аналіз відносних показників коефіцієнтів маси внутрішніх органів тварин засвідчив, що застосування водного розчину упареного залишку БАР КЗПЕР не призвело до їх зміни порівняно з тваринами групи негативного контролю, як у самців, так і у самок. Показники знаходяться в межах значень контрольної групи.

Висновки. За результатами дослідження, дана схема експерименту не виявила ознак летальності тварин та токсичних ефектів між тваринами дослідних груп. Вплив на поведінку тварин, на органи тварин (масу тварин та відносні коефіцієнти мас внутрішніх органів), на слизову оболонку ШКТ, як у самців, так і у самок, залишався відсутнім. Отже, результати проведеного експерименту свідчать про те, що при внутрішньошлунковому введенні водний розчин упареного залишку БАР КЗПЕР у об’ємі 20 мл/кг не виявляв негативний вплив на щурів (самців і самок), що дозволяє віднести його до VI класу токсичності (відносно нешкідливі речовини).

Дослідження гострої токсичності рослинної субстанції КЗПЕР дозволило встановити безпечність при її використанні у складі таблетованих пероральних ЛЗ для застосування в кардіологічній практиці.

ПЕРСПЕКТИВНІ ВИДИ ВІТЧИЗНЯНОЇ ЛІКАРСЬКОЇ РОСЛИННОЇ СИРОВИНИ, ЩО ЗАСТОСОВУЮТЬСЯ У ТЕРАПІЇ ЗАХВОРЮВАНЬ ОПОРНО-РУХОВОГО АПАРАТУ

Хворост О. П., Опрошанська Т. В., Скребцова К. С., Горда А. О.
Національний фармацевтичний університет, м. Харків, Україна
Кафедра фармакогнозії та нутриціології
Кафедра фармацевтичної технології, стандартизації та сертифікації ліків
khvorost09101960@gmail.com

Вступ. За даними ВООЗ захворювання опорно-рухової системи призводять до обмеження рухливості і посідають друге місце у світі як фактор постійної втрати працездатності та 4-те місце у світі як причина інвалідності та смертності. В Україні близько 3,5 млн осіб мають проблеми з опорно-руховим апаратом та його різними ускладненнями, які вимагають безперервної багаторічної терапії. Альтернативою синтетичним лікарським засобам є фітозасоби, які виявляють широкий спектр фармакологічної активності. Безумовно, вони поступаються своєю протизапальною активністю, але є більш безпечними для здоров'я пацієнта та завдяки різноманітному хімічному складу, виявляють комплексну дію на організм, що значно полегшує перебіг захворювань опорно-рухової системи.

Мета. Метою роботи – проаналізувати асортимент лікарської рослинної сировини наукової та традиційної медицини, що застосовуються у лікуванні захворювань опорно-рухової системи, та обрати найперспективніші.

Матеріали та методи. контент-аналіз доступних джерел інформації з подальшою інтерпретацією результатів.

Результати. Аналіз літературних джерел щодо рослинної сировини, яка використовується в народній медицині для лікування даних захворювань, показав що компонентами зборів є сировина представників родин *Asteraceae, Betulaceae, Boraginacae, Fabaceae, Polygonaceae*, *Rosaceae, Pionaceae, Equisetaceae, Poaceae та Urticaceae,* яка містить поліфенольні сполуки, органічні кислоти, аскорбінову кислоту, полісахариди, леткі сполуки, вітаміни, мінерали та проявляє широкий спектр фармакологічної активності. Нами обрано 29 видів рослинної сировини, джерела отримання яких належать до 10 родин. Із обраних видів для подальшого дослідження на основі скринінгу антиоксидантної активності *in vitro* та кількісного вмісту суми поліфенолів обрано сировину представників родин *Polygonaceae, Rosaceae* та *Asteraceae*: корені л. великого, л. малого і л. павутинистого, корені ш. собачої і ш. травневої, кореневище з коренями р. лікарського, кореневище з коренями щ. кінського.

Висновки: подальше фармакогностичне дослідження цих видів сировини представників родин *Rocaceae, Polygonaceae, Asteraceaeє* є актуальним для створення на їх основі ефективних засобів комплексної терапії.

MODERN ASPECTS OF COMPLEX THERAPY FOR PATIENTS WITH PSORIASIS

Galuzinska L. V., Seniuk I. V.
National University of Pharmacy, Kharkiv, Ukraine
Department of Clinical Laboratory Diagnostics, Microbiology and Biological Chemistry

Introduction. Psoriasis is one of the most common chronic skin diseases, and according to statistics from various authors, it affects about 3-4% of the world's population. Despite the widespread prevalence of psoriasis and numerous studies of this problem, there is currently no unanimous opinion on its pathogenesis and treatment methods.

Aim. The aim is to find new ways to treatfor patients with psoriasis.

Materials and methods. To substantiate the effectiveness of treatment of patients with psoriasis using the amino acid and coenzyme complex Cardonate, three groups of patients were formed: in the first group, general topical therapy with the inclusion of Cardonate was performed; in the second group, a traditional treatment regimen (systemic and topical therapy) was performed; in the third group, treatment was performed according to the traditional methodology supplemented with Cardonate. The effectiveness of the treatment was assessed by the results of the study in the dynamics (before and after treatment) of immunological, biochemical and clinical parameters.

Results. The analysis of cellular and humoral immunity parameters showed that during the treatment of patients, the reactivity of the immune system decreased to some extent, which led to the formation of endothelial dysfunction of the microcirculatory vascular bed. This was especially noticeable in the treatment of the third group of patients, a significant decrease in the level of immunoglobulins E indicates the suppression of the manifestation of autoimmune processes and the effectiveness of metabolite-adapted complex treatment of patients in this group, as evidenced by the length of stay of patients in the hospital.

The study of the effectiveness of treatment of patients was carried out by the dynamics of changes in monitoring metabolic parameters: AST, ALT, γ-GT, ALP, LDH, CK, CK-MB, bilirubin, creatinine, urea, total protein, glucose, albumin, cholesterol, phospholipids, , phosphorus, The effectiveness of treatment was assessed in relation to the traditional treatment regimen.

Conclusions. The results of the study showed that the additional administration of the amino acid coenzyme complex ‘Cardonate’ to the traditional regimen in patients with advanced psoriasis reduced the time to remission and contributed to the normalisation of monitoring metabolic parameters. The efficacy of the proposed therapy for patients with widespread psoriasis is confirmed by the data on shortening the course of the progressive and stationary stages by 5-8 days and the onset of clinical remission.

ВИВЧЕННЯ ПРОТИЗАПАЛЬНОЇ АКТИВНОСТІ КРЕМУ «ПІРИСАЛІКС»

Малоштан Л. М., Підгайна В. В., Яковлєва Л. В.

Національний технічний університет «Харківський політехнічний інститут», м. Харків, Україна

Кафедра загальної фармації

lnm004@gmail.com

Національний фармацевтичний університет, м. Харків, Україна

valentina.pidgaina@gmail.com

Вступ. Запалення є одним з складних процесів, які часто зустрічаються в патології та являються причиною багатьох патологічних процесів, а несвоєчасне лікування може трансформуватися в типовий патологічний процес.

Тому однією з актуальних проблем сучасної медицини є регуляція запального процесу. Використання для цієї цілі синтетичних лікарських препаратів не завжди впливає сприятливо, оскільки може викликати ряд небажаних побічних ефектів.

В зв'язку з цим залишається актуальним пошук ефективних лікарських засобів з протизапальним ефектом і мінімальними побічними діями. Запальна реакція складає початкову фазу ранового процесу та розвивається у відповідь на дію альтеруючого чинника, при цьому її роль полягає в локалізації та ліквідації ушкоджуючого чинника, а також забезпеченні умов для наступного відновлення тканин.

Проте виражений запальний процес сам по собі може обумовлювати вторинне ушкодження тканин та затримувати загоєння рани, тому виключно важливе значення для реалізації лікувальної дії засобів, що застосовуються в терапії ранового процесу, є наявність протизапальних властивостей.

Мета. Дослідження є фрагментом роботи, метою якої було вивчення фармакологічних властивостей крему на основі кори верби білої та цинку під умовною назвою «Пірисалікс» та ефективності його місцевого застосування для лікування уражень шкіри та слизових оболонок різного походження в якості репаративного засобу.

Матеріали та методи. Вивчення протизапальної активності крему «Пірисалікс» проводили на моделі термічного запалення лап у мишей. Досліди було проведено на білих мишах, які були поділені на три експериментальні групи: перша група використовувалась в якості контрольної, тварин ІІ-ої групи лікували кремом «Пірисалікс», ІІІ-ої референт препаратом- кремом «Псорікап».

Термічне запалення лапи у мишей відтворювали шляхом занурення правої задньої кінцівки у гарячу (66,5°С) воду на 4 секунди. Після цього тваринам другої дослідної групи на обпечену кінцівку наносили крем «Пірисалікс», тваринам третьої групи - препарат порівняння крем «Псорікап». У тварин контрольної групи лікування ураженої кінцівки не проводили.

Через 24 години піддослідних тварин виводили з експерименту шляхом дислокації шийних хребців під наркозом. Виразність запального процесу оцінювали за збільшенням ваги ураженої кінцівки. Для цього обидві кінцівки (здорову та ушкоджену) відрізали на рівні гомілковостопного суглобу та зважували на торсійних терезах марки «ВТ-500» і розраховували різницю в масі набряклої та здорової лапи.

Результати, що були отримані при обстеженні тварин дослідних груп, порівнювали зі значенням досліджуваного показника в контрольній групі та розраховували протизапальну активність, яку виражали у відсотках.

Результати. Спостереження показало, що вплив температурного фактора призводить до розвитку запального процесу з вираженим ексудативним компонентом: у піддослідних тварин спостерігалося почервоніння ураженої кінцівки, набряк та біль. Отримані в ході експерименту дані свідчать, що місцеве застосування крему «Пірисалікс» сприяє зменшенню виразності запального процесу, що виявляється зниженням маси набряклої кінцівки, порівняно з контролем, на 35,7%. Встановлено також, що за виразністю протизапального ефекту крем «Пірисалікс» в 1,4 рази перевищує препарат порівняння крем «Псорікап» (протизапальна активність – 25,5%).

Висновки. Таким чином, дослідження проведені на моделі гострого ексудативного запалення, викликаного дією термічного чинника показали, що крем «Пірисалікс» виявляє виразні протизапальні властивості.

Було також встановлено, що за виразністю протизапальної дії на моделі термічного запалення лап у мишей крем «Пірисалікс» виявляє певну перевагу перед препаратом порівняння кремом «Псорікап».

ПЕРСПЕКТИВИ ВИКОРИСТАННЯ ЦИТОСТАТИКА ЦИКЛОФОСФАМІДУ ДЛЯ ЕКСПЕРИМЕНТАЛЬНОГО ВІДТВОРЕННЯ ВТОРИННОГО ІМУНОДЕФІЦИТУ У ЩУРІВ ДИТЯЧОГО ВІКУ

Єрмоленко Т. І., Шаповал О. М.

Харківський національний медичний університет, м. Харків, Україна

Кафедра фармакології та медичної рецептури

kaf.1med.farmakologii@knmu.edu.ua

Вступ. Беззаперечним фактом є те, що здоров’я нації, і української – зокрема, обумовлене якістю здоров’я дітей та підлітків, що відбивається на тривалості та повноцінності подальшого життя кожного майбутнього громадянина й на розвитку усього соціуму. Згідно з даними Глобальної стратегії ВООЗ з охорони здоров'я жінок і дітей визначено, що інвестиції у здоров'я дітей мають високу економічну доцільність і гарантують наявність продуктивної робочої сили у майбутньому. Специфіка перебігу соціально-економічних і демографічних процесів в Україні у перші десятиріччя XXI сторіччя, таких як епідемії та воєн-

ні дії внаслідок російської агресії призвела до збільшення захворюваності та розповсюдженості таких хвороб, що обумовлюють розвиток вторинних імунодефіцитних розладів здоров'я дитячого населення. Вторинні імунодефіцити є порушенням функціонування імунітету, що виникає на тлі нормально сформованої імунної системи в процесі життя і не пов'язане з генетичними дефектами. Вторинний імунодефіцит зустрічається частіше, ніж первинний імунодефіцит. Вторинні імунодефіцити у дітей не є самостійною нозологією, а розвиваються внаслідок компрометації такими факторами, як інфекційні захворювання (вірусні, бактеріальні та грибкові), специфічні дисфункції органів та систем, в тому числі органів дихання, травми та операційні втручання, застосування імуносупресивних лікарських засобів при терапії аутоімунних та онкологічних захворювань та тощо. Тому такі пацієнти потребують фармакокорекції стану імунної системи для скорішого та якісного одужання.

Мета. Зважаючи на вищевикладене та в рамках пошуку та створення нових лікарських засобів з імунотропними властивостями для терапії дітей, метою даної роботи є експериментальне відтворення вторинного імунодефіциту у щурів віком 1 міс., який відповідає людському дитячому віку 4-7 років, компрометованого цитостатиком циклофосфамідом.

Матеріали та методи. Дослідження проводили з дотриманням загальноприйнятих в Україні етичних принципів поводження з лабораторними тваринами й з використанням 20 білих безпорідних статевонезрілих щурів віком 1 міс, який відповідає людському дитячому віку 4-7 років, з початковою масою 90-110 г., які були розподілені на 2 групи: 1) ІК – інтактний контроль та 2) ЦФА – група тварин, яка отримувала циклофосфамід. Вторинний компрометований імунодефіцит (ВКІ) моделювали за допомогою внутрішньом'язового введення щурятам цитостатика циклофосфаміду в дозі 10 мг/кг 1 раз на добу протягом 7 діб. В цей же час групі тварин ІК уводили внутрішньом'язово адекватну кількість води для ін'єкцій. На початку та наприкінці експерименту визначали масу тіла тварин для оцінки трофічних процесів в їхньому організмі. На 8-му добу тварин обох груп виводили з досліду декапітацією під наркозом, збирали кров для проведення клінічного аналізу крові з метою визначення гематологічних показників та показників, що характеризують стан імунної системи: фагоцитарну активність лейкоцитів, активність комплементу, концентрацію циркулюючих імунних комплексів (ЦІК) в сироватці крові. Також з метою оцінки трофічних процесів в організмі тварин та його функціонального стану визначали масу тіла тварин до виводу з досліду, а після знеживлення виймали основні органи та визначали їх масу, необхідну для розрахунку коефіцієнту маси внутрішніх органів.

Результати. Аналіз результатів впливу цитостатика циклофосфаміду на трофічні процеси, гематологічні та імунологічні показники стану організму щурів віком 1 місяць, який відповідає людському дитячому віку 4-7 років, в порівнянні з групою інтактного контролю такого ж віку, свідчать про симптоми розвитку вторинного імунодефіциту. Так, у групі ЦФА спостерігали достовірне в

порівнянні з групою ІК зниження маси та коефіцієнтів маси селезінки на 112% та тимусу – на 152,6%. Як відомо, ці органи беруть безпосередню участь в діяльності імунної системи: у селезінці відбувається утворення імунокомпетентних клітин, а в тимусі – лейкоцитів та лімфоцитів. Ці дані корелюють з гематологічними та імунологічними показниками, визначеними в цьому дослідженні, та що характеризують діяльність імунної системи. Такий вплив ЦФА на ці імунокомпетентні органи відбивається достовірним в порівнянні з групою ІК зниженням кількості лейкоцитів на 70% та моноцитів - на 68,5%. В групі ЦФА спостерігали достовірне в порівнянні з групою ІК зниження кількості ЦІК на 54% та на 353% (з 127 у.о. в групі ІК до 28 у.о. в групі ЦФА) – індекс фагоцитарної активності нейтрофілів, що свідчить про значне пригнічення неспецифічної імунної відповіді організму 1-місячних щурят.

Висновки. Таким чином, отримані результати свідчать про те, що спроба відтворити штучний вторинний імунодефіцит у щурят віком 1 місяць, який відповідає людському дитячому віку 4-7 років, компрометований цитостатиком циклофосфамідом, є вдалою. Отже цю експериментальну модель вторинного компроментованого імунодефіциту можна використовувати для подальших досліджень з метою пошуку та створення потенційних лікарських засобів, які можуть застосовуватися в комплексній терапії дітей віком 4-7 років з захворюваннями, що супроводжуються вторинними імунодефіцитами.

ПАТОГЕНЕТИЧНЕ ОБГРУНТУВАННЯ ЕФЕКТИВОСТІ КАРБАЦЕТАМУ ПРИ МЕТАБОЛІЧНИХ ПОРУШЕННЯХ У ЩУРІВ

Прижбило О. М., Кметь О. Г.

Буковинський державний медичний університет,
м. Чернівці, Україна
Кафедра фармакології
prizbilo0950721@gmail.com

Вступ. На сьогоднішній день метаболічний синдром є актуальною медико-соціальною проблемою, яка привертає увагу спеціалістів різного профілю. Враховуючи зростання показників розповсюдженості патологічного стану, насущою проблемою сучасної науки залишається розуміння причин його розвитку і прогресування, розробка та удосконалення методів діагностики, профілактики та лікування.

Відомо, що багато факторів впливають на патофізіологію метаболічного синдрому. Однак, є наукові дані, які засвідчують, що окислювальний стрес у поєднанні з хронічними запальними станами є основою розвитку метаболічних захворювань. Водночас дисбаланс між оксидантами та антиоксидантами, який часто схиляється на користь оксидантів, є причиною окислювального стресу, який, у свою чергу, спричиняє порушення окисно-відновної передачі сигналів

та регуляції, а також молекулярне та клітинне пошкодження. Відповідно дисбаланс між прооксидантними та антиоксидантними механізмами вважається одним із найважливіших патофізіологічних механізмів хронічних захворювань. За науковими даними окислювальний стрес можна вважати ранньою діагностичною ознакою в механізмах метаболічного синдрому, а не просто його наслідком.

Опираючись на сучасні дані розуміння нейрохімічних та нейроанатомічних основ мозкової діяльності, зокрема зв'язок між різними системами нейротрансмітерів, що є основою більш специфічного та цілеспрямованого лікування, сприяє зменшенню кількістю побічних ефектів. Зокрема, відомо про широке розповсюдження ГАМК-ергічних рецепторів у головному мозку. Тому нас зацікавило з’ясувати їхню роль у процесах пероксидації ліпідів та білків при метаболічному синдромі.

Мета. Обґрунтувати ефективність карбацетаму, як модулятора ГАМК-рецепторів, на основі вивчення стану проксидантної системи гіпокампа щурів з метаболічним синдромом.

Матеріали та методи. Дослідження проводились на лабораторних нелінійних білих щурах чоловічої статті з дотриманням принципів Гельсінської декларації щодо гуманного поводження з тваринами та Європейської конвенції із захисту хребетних тварин, які використовують в експериментах та інших наукових цілях. Для відтворення моделі метаболічного синдрому щурів 60 діб утримували на високожировій дієті збагаченій жирами за рахунок твердого свинячого жиру з вільним доступом до розчину фруктози. Карбацетам вводили внутрішньоочеревинно дозою 5 мг/кг 14 днів. Евтаназію щурів здійснювали під наркозом хлороформом. На холоді виймали головний мозок і за стереотаксичним атласом виділяли гіпокамп. Інтенсивність пероксидного окиснення ліпідів оцінювали за вмістом продуктів, що реагують із 2-тіобарбітуровою кислотою (ТБКАП). Вміст окиснювальної модифікації білків (ОМБ) у гомогенетах визначали методом спектрофотометрії при довжині хвилі 370 і 430 нм: 370 нм – визначали кетодинітрофенілгідразони нейтрального характеру; 430 нм – альдегідодинітрофенілгідразони основного характеру. Для обробки результатів дослідження обраховували середню арифметичну та її похибку, розбіжності у середніх тенденціях перевіряли за допомогою критерію Стьюдента після позитивної перевірки вибірки на нормальність розподілу в ній за критерієм Shapiro-Wilk.

Результати. На основі проведених нами досліджень встановлено, що у гіпокапмі щурів із метаболічним синдромом збільшувався вміст ТБКАП. Так, порівняно з контрольною групою, вміст ТБКАП збільшувався на 85,4 %. Водночас у щурів із метаболічним синдромом виявлено підвищення процесів пероксидного окиснення білків при різних довжинах хвиль. Так, вміст ОМБ 370 та ОМБ 430 перевищували значення контрольної групи на 81,4 % і на 35,8 % відповідно. Такі показники інформують про інтенсифікацію окислювальної деструкції ліпідів і білків, як важливих складових мембран клітин, зокрема нейронів, що порушує їхню функціональну активність.

Після введення карбацетаму вміст ТБКАП знижувався на 28,2 %. Водночас спостерігали зменшення вміст ОМБ. Зокрема, вміст кетодинітрофенілгідразонів нейтрального характеру, знижувався на 16,9 % та альдегідодинітрофенілгідразони основного на – 15,4 %.

Отже, на основі отриманих результатів дослідження, можна припустити, що модуляція ГАМК-рецепторів сприяла зниженню процесів пероксидного окиснення ліпідів та білків нейронів гіпокампа щурів із метаболічним синдромом.

Висновок. Отримані результати свідчать про корегувальний вплив карбацетаму через модуляцію ГАМК-рецепторів нейронів гіпокампа на маркери оксидативного стресу у щурів із метаболічним синдромом.

COMBINED USE OF THE SUBSTANCES OF HERBAL ORIGIN WITH XANTHINE OXIDASE INHIBITORS: EXAMPLE OF *AEGOPODIUM PODAGRARIA* L.

Tovchiga O. V.[2], Koiro O. O.[1], Shtrygol' S. Yu.[1], Stepanova S. I.[1]
[1]National University of Pharmacy, Kharkiv, Ukraine
Department of Pharmacology and Clinical Pharmacy
[2]Medical University of Gdańsk, Gdańsk, the Republic of Poland
Department of Pharmaceutical Pathophysiology
olga.tovchiga@gumed.edu.ua

Introduction. Xanthine oxidase (XOD) is not only a significant target for antigout interventions, but a pathogenetically important enzyme for overcoming of cardiovascular and renal diseases, limiting of oxidative stress, normalization of the CNS function [1, 2]. Search of the new biologically active substances is being actively conducted among plant sources. High massive of evidence has been collected about XOD inhibitors of plant origin as well as herbal substances able to influence renal excretion of uric acid [3, 4]. Less data are available about possibilities of such inhibitors combinations with currently used XOD inhibitors, namely allopurinol and febuxostat. These aspects are important because of the possibility of the reducing of the dose of the latter thus increasing their safety or broadening pharmacodynamics with additional favourable effects.

Goutweed (*Aegopodium podagraria* L., GW) is wide-spread in temperate climates including northern and western Ukraine and as evidenced by Latin, English and Polish names of the plant, has been used in traditional medicine for gout treatment since time immemorial. Our previous studies have confirmed favourable influence of GW aerial part tincture and extract on uric acid metabolism realized through xanthine oxidase (XOD) inhibition, enhancement of the renal uric acid excretion or combination of both mechanisms [5, 6]. Further studies were directed towards verifi-

cation of the results of combined administration of GW preparations with commonly used drugs [7].

The aim of the study. This study aimed to provide a summary of the own results concerning herbal drugs combinations with XOD inhibitor and explore new research areas based on the data found in the literature.

Results. A number of important data on the possibilities of GW combined use with allopurinol (ALL) as the widely used and commonly available hypouricemic drug were obtained [7]. Namely, the effects of such combinations were as follows:

– combination of GW tincture (1 ml/kg) or GW extract (1 g/kg) with a subtherapeutic dose of ALL of 2.5 mg/kg enabled statistically significant XOD inhibition under conditions of potassium oxonate induced hyperuricemia in mice (after single administration);

– complete inhibition of XOD by ALL (10 mg/kg) was maintained when it was combined with the tincture (1 ml/kg) or the extract (1 g/kg) during the prolonged (3-weeks) hyperuricemia induced by potassium oxonate in mice;

– absence of the increase in toxicity of ALL at a high dose (50 mg/kg) combined with the tincture (1 ml/kg) and moderate nephroprotective effects of the latter (decrease in proteinuria, maintenance of sodium reabsorption as well as normokalemia, urea clearance normalization) in rats against the background of an excess of purine derivatives and proteins;

– decrease in the acute toxicity of ALL in mice against the background of the extract and the tincture.

The studied GW preparations main components are hydroxycinnamic acids and flavonoids. In the literature available, there are reports about in vivo studies of herbal extracts and ALL combinations efficacy, still these herbal preparations are of different chemical composition [8, 9]. And as for the mechanism of XOD inhibition by plant polyphenols, it has been studied for the numerous compounds and it is important that both competitive and non-competitive inhibitors were revealed [3, 4]. There are also robust data about chlorogenic acid as XOD inhibitor [10] as well as data about the benefits of coffee as a source of hydroxycinnamic acids for gout prevention [11].

Proceeding from the beforementioned data, the next logical step is the assessment of the possible mechanisms of interaction of these polyphenols and ALL (or other XOD inhibitors) which could be realized at the level of XOD active site or allosteric sites. Besides, there is a lack of data of the changes of prooxidant-antioxidant status under the influence of such combinations, given the role of XOD in generation of free radicals.

Thus, it could be expedient to take the following steps in the research of XOD inhibitors combinations: *in silico* modeling of the interaction with the enzyme; *in vivo* assessment of enzyme activity with individual substances combinations as well as their influence on the prooxidant-antioxidant processes, especially the superoxide generation under the influence of XOD; *in vitro* verification of the most promising combinations activity and safety.

Other aspect that may seem to be unexpected, still possessing sufficient evidence base, is the positive influence of XOD inhibitors on the CNS function [12, 15]. This influence is possibly realized through the mechanism other than hypouricemic action (since uric acid is neuroprotective itself at the certain range of concentrations) and can include (being not limited to) the reducing of the prooxidant role of XOD mentioned above especially that localized within the cerebral vascular bed. In this context, flavonoids are discussed as promising XOD inhibitors with the possible benefits for mental disorders treatment [13]. For chlorogenic acid there are data both on the normalizing influence on uric acid metabolism (also mentioned above) and experimentally established ability to decrease anxiogenic and depressive processes in the laboratory animals [14].

Thus, it is of great interest, how these effects of chlorogenic acid (or those of flavonoids) will be modulated in their combined use with ALL or other XOD inhibitors. Interactions results could proceed from changes of uricemia, from the limiting of oxidative stress during XOD functioning and from the changes of specific mechanisms of these herbal compounds mentioned above.

Partially we addressed these issues in studies of GW combinations with ALL [7]. Importantly, on the model of hyperuricemia, when uric acid level was normalized by ALL combined with GW extract or the tincture, no negative changes of the CNS functional state were registered. Moreover, when combined with ALL, GW preparations counteracted the increase in depressive signs, the tincture also reduced anxiety signs.

Conclusions. The data presented elucidate the prospects of the combined use of the substances of herbal origin with xanthine oxidase inhibitors.

References:

1. Furuhashi M. New insights into purine metabolism in metabolic diseases: role of xanthine oxidoreductase activity. American Journal of Physiology-Endocrinology and Metabolism 2020;319:E827–34. https://doi.org/10.1152/ajpendo.00378.2020.
2. [2] Burrage EN, Coblentz T, Prabhu SS, Childers R, Bryner RW, Lewis SE, et al. Xanthine oxidase mediates chronic stress-induced cerebrovascular dysfunction and cognitive impairment. Journal of Cerebral Blood Flow & Metabolism 2023; 43:905–20.
3. Xue H, Xu M, Gong D, Zhang G. Mechanism of flavonoids inhibiting xanthine oxidase and alleviating hyperuricemia from structure–activity relationship and animal experiments: A review. Food Front 2023; 4:1643–65. https://doi.org/10.1002/fft2.287.
4. Liu L, Zhang L, Ren L, Xie Y. Advances in structures required of polyphenols for xanthine oxidase inhibition. Food Front 2020; 1:152–67. https://doi.org/10.1002/fft2.27.
5. Tovchiga OV. The investigation of the goutweed (Aegopodium podagraria L.) diuretic, nephroprotective and hypouricemic action as the basis for the drug development. PhD thesis in specialty “Pharmacology.” National University of Pharmacy, 2009.
6. Koiro OO. Role of goutweed (Aegopodium podagraria L.) biologically active substances in nephroprotective, hepatoprotective and hypouricemic activity. PhD thesis in specialty “Pharmacology.” National University of Pharmacy, 2014.
7. Tovchiga OV. The antidiabetic and organotropic action of goutweed (Aegopodium podagraria L.) preparations and their combinations with antihyperglycemic, diuretic, and hypouricemic drugs . doctoral thesis in specialty “Pharmacology.” National University of Pharmacy, 2019.
8. Sung Y-Y, Kim D-S. Synergistic Impacts of Alpinia oxyphylla Seed Extract and Allopurinol against Experimental Hyperuricemia. Biomed Res Int 2022; 2022:1–9.

9. Mehmood A, Zhao L, Wang C, Hossen I, Nadeem M. Stevia residue extract alone and combination with allopurinol attenuate hyperuricemia in fructose–PO-induced hyperuricemic mice. J Food Biochem 2020;44. https://doi.org/10.1111/jfbc.13087.
10. Gawlik-Dziki U, Dziki D, Świeca M, Nowak R. Mechanism of action and interactions between xanthine oxidase inhibitors derived from natural sources of chlorogenic and ferulic acids. Food Chem 2017;225:138–45. https://doi.org/10.1016/j.foodchem.2017.01.016.
11. Shirai Y, Nakayama A, Kawamura Y, Toyoda Y, Nakatochi M, Shimizu S, et al. Coffee Consumption Reduces Gout Risk Independently of Serum Uric Acid Levels: Mendelian Randomization Analyses Across Ancestry Populations. ACR Open Rheumatol 2022;4:534–9. https://doi.org/10.1002/acr2.11425.
12. Tovchiga O V., Shtrygol' SYu. Uric acid and central nervous system functioning (a literature review). Biology Bulletin Reviews 2014;4:210–21.
13. Martorell M, Lucas X, Alarcón-Zapata P, Capó X, Quetglas-Llabrés MM, Tejada S, et al. Targeting Xanthine Oxidase by Natural Products as a Therapeutic Approach for Mental Disorders. Curr Pharm Des 2021;27:367–82. https://doi.org/10.2174/1381612826666200621165839.
14. Nguyen V, Taine EG, Meng D, Cui T, Tan W. Chlorogenic Acid: A Systematic Review on the Biological Functions, Mechanistic Actions, and Therapeutic Potentials. Nutrients 2024;16:924. https://doi.org/10.3390/nu16070924.
15. Song Y, Racette BA, Camacho-Soto A, Searles Nielsen S. Biologic targets of prescription medications and risk of neurodegenerative disease in United States Medicare beneficiaries. PLOS ONE. 2023 May 17;18(5):e0285011.

THE TOPICAL ISSUES ASPECTS OF CHARACTERISTICS OF ENDOVASCULAR SURGERY AND PHARMACOLOGICAL AND NON-PHARMACOLOGICAL TREATMENT CHALLENGES FOR GASTROINTESTINAL AND DUODENAL ULCER BLEEDING WITH BRIEF CASE REPORT

Gocha Chankseliani[1], Avtandil Girdaladze[1], Omar Gibradze[2], Paata Meshveliani[2], Kakha Chelidze[3], Mirian Cheishvili[3], Ana Kvernadze[3], Nodar Sulashvili[4]

[1]Tbilisi State University; Faculty of Medicine;
[2]Akaki Tsereteli State University, Faculty of Medicine;
[3]Multiprofile-Clinic «L&J»;
[4]Department of Pharmaceutical Management of Yerevan State Medical University after Mkhitar Heratsi, Yerevan, Armenia
n.sulashvili@ug.edu.ge

Abstract. Acute gastrointestinal bleeding is a common medical emergency that ranges from minor to potentially life-threatening bleeding. Endoscopy is the first-line diagnostic procedure for upper and lower gastrointestinal bleeding. Treatment options for acute GI bleeding include conservative management, therapeutic endoscopy, transcatheter embolization, and surgery. Transcatheter embolization and surgery are both options for recurrent GI bleeding when therapeutic endoscopy fails; However, both options are associated with several complications and risk of bleeding. The choice of management depends on the patient's status. Emergency surgery is usually associated with high rates of morbidity and mortality. Recently, superelective

transcatheter embolization has become a safer procedure and is now widely used to treat acute gastrointestinal bleeding. This review article describes the role of interventional radiology in the management of acute GI bleeding. Improvements in catheter technology, development of more compatible embolization devices, and expansion of embolization techniques have led to angiography and embolization for the treatment of upper and lower gastrointestinal bleeding. Transcatheter embolization therapy for the treatment of acute GI bleeding is a safe procedure with high technical performance and clinical success, but it should be reserved as a treatment option for patients who have failed endoscopic and medical management. MDCT imaging is a useful tool for identifying the site of bleeding and evaluating the anatomical structure of the gastrointestinal tract in stable patients. Close working relationships between interventional radiologists, gastroenterologists, and diagnostic radiologists are essential for the optimal management of patients with GI bleeding. Endovascular embolization dramatically reduces the mortality rate in high-risk patients who require open surgery after failed endoscopy, further studies are needed to fully address these objectives.

Key words: *Endovascular surgery, gastrointestinal duodenal, ulcer bleeding, treatment.*

Introduction. Acute gastrointestinal bleeding is a common medical emergency that ranges from minor, uncontrollable bleeding to potentially life-threatening bleeding. The site of bleeding can be located anywhere in the gastrointestinal tract, which makes it difficult to determine its exact location. Patients with upper gastrointestinal bleeding usually have hematemesis or melena with the bleeding point proximal to the ligament of Treitz, while patients with lower gastrointestinal bleeding usually have melena or hematochezia with the bleeding point distal to the ligament of Treitz. Endoscopy is a first-line diagnostic procedure with 100% sensitivity to detect upper gastrointestinal bleeding; However, it has only 60% sensitivity for diagnosing lower gastrointestinal bleeding. Therapeutic options for treating acute GI bleeding include conservative treatment, therapeutic endoscopy, transcatheter embolization, and surgery. Transcatheter embolization and surgery are both options for recurrent GI bleeding [1-4]. When therapeutic endoscopy fails; However, both options are associated with several complications and risk of bleeding. The choice of management depends on the patient's status (for example, the degree of hemodynamic instability or hypotension, or whether resuscitation is required). Emergency surgery is usually associated with high rates of morbidity and mortality. However, recent technical improvements in superselective transcatheter embolization have increased the safety of the procedure, and it is widely used in the treatment of acute gastrointestinal bleeding. This review article describes the role of interventional radiology in the emergency management of acute GI bleeding [5-8].

Upper gastrointestinal bleeding is defined as bleeding originating from the distal esophagus, stomach, or duodenum (ie, proximal to the ligament of Treitz). The most common cause of upper GI bleeding is peptic ulcer disease, but the differential

diagnosis is varied and includes benign and malignant tumors, ischemia, gastritis, arteriovenous malformations, Mallory-Weiss tears, trauma such as a Dieulafoy injury, and iatrogenic causes [9-12].

Upper gastrointestinal bleeding is a potentially fatal condition, so immediate management and accurate diagnosis of the location and etiology of bleeding is essential. The primary diagnostic procedure for upper gastrointestinal bleeding is endoscopy, which has high sensitivity and specificity for locating bleeding lesions in the upper gastrointestinal tract. Once a bleeding lesion is identified, therapeutic endoscopic techniques such as thermal coagulation or hemoclip placement can be used to achieve acute hemostasis. Endoscopic management achieves hemostasis in most patients, but 10% to 30% of patients experience recurrent bleeding for various reasons [13-16].

When hemostasis is not achieved with endoscopic management, other options are surgery and transarterial embolization. Surgery has long been the standard of care, but with the development of interventional radiology, more and more patients are now being referred for embolotherapy. Transarterial embolization can prevent unnecessary resection of the upper gastrointestinal tract and should be considered as an alternative to surgery [17-19].

Aim of the research was to study the key issue aspects of features of endovascular surgery and treatment for gastrointestinal and duodenal ulcer bleeding and brief case report.

Materials and Methods. The material of the article was the data from scientific publications, which were processed, analyzed, overviewed and reviewed by generalization and systematization. research studies are based on a review/overview assessment of the development of critical visibility and overlook of the modern scientific literature. use the following databases: (for extensive literature searches to identify key issue aspects of features of endovascular surgery and treatment for gastrointestinal and duodenal ulcer bleeding and brief case report). PubMed, Web of Science, Clinical Key, Tomson Reuters, Google Scholar, Cochrane library, and Elsevier foundations. national and international policies and guidelines were also reviewed and as well as grey literature.

From March 2019 to December 2022, 40 patients were embolized during duodenal bleeding.

These patients were divided into the following groups:

– Massive, active bleeding.
– Recurrence of bleeding in clinic and endoscopy was unsuccessful.
– Unstable hemodynamics and solid hemostasis could not be achieved during endoscopy.
– Failed to evacuate stomach contents and failed to see a bleeding ulcer.
– High risk of bleeding recurrence (Forest classification).
– Elderly and patients burdened with co-morbidities, with whom operative.

Result and Discussion. Lower gastrointestinal bleeding is defined as bleeding originating from a source distal to the ligaments of Treitz. About 80% of all lower GI bleeding comes from a colorectal source and 5% to 10% from a small intestinal

source; 10% to 15% are classified as blood of upper gastrointestinal tract origin. A small bowel bleeding source is more likely than a colorectal bleeding source to be obscure or occult.

A common cause of lower GI bleeding is a colonic diverticulum. Differential diagnoses include colitis or enteritis, anorectal abnormalities (hemorrhoids, proctitis), tumors, arteriovenous malformations or angiodysplasia, and postpolyectomy bleeding.

Therapeutic colonoscopy is currently the first-line intervention for colonic bleeding. Colonoscopy is the diagnostic method of choice in patients with lower gastrointestinal bleeding, but therapeutic endoscopy can also be successful in a limited number of patients. Therapeutic colonoscopy fails in approximately 32% of cases due to the presence of stool or blood clots, or technical difficulties such as time required to prepare patients. Further disadvantages include the fact that small bowel bleeding cannot be accessed through colonoscopy and that colonoscopy is relatively ineffective when performed in patients with significant bleeding without bowel preparation. If bleeding cannot be stopped by therapeutic colonoscopy, transarterial embolization is the next line of therapy to control hemostasis. As with upper gastrointestinal bleeding, transarterial embolization is the first-line therapy for patients with lower gastrointestinal bleeding. The efficacy of transarterial embolization in the treatment of acute gastrointestinal bleeding when medical or endoscopic techniques are inadequate has been demonstrated in several large studies. For both upper and lower gastrointestinal bleeding, surgery is usually reserved as a last-line treatment for patients whose bleeding has failed to respond to previous treatments.

Lower gastrointestinal bleeding is defined as bleeding originating from a source distal to the ligaments of Treitz. About 80% of all lower GI bleeding comes from a colorectal source and 5% to 10% from a small intestinal source; 10% to 15% are classified as blood of upper gastrointestinal tract origin. A small bowel bleeding source is more likely than a colorectal bleeding source to be obscure or occult. A common cause of upper GI bleeding is colonic diverticula. Differential diagnoses include colitis or enteritis, anorectal abnormalities (hemorrhoids, proctitis), tumors, arteriovenous malformations or angiodysplasia, and postpolyectomy bleeding [20-24].

Therapeutic colonoscopy is currently the first-line intervention for colonic bleeding. Colonoscopy is the diagnostic method of choice in patients with lower gastrointestinal bleeding, but therapeutic endoscopy can also be successful in a limited number of patients. Therapeutic colonoscopy fails in approximately 32% of cases due to the presence of stool or blood clots. Technical difficulties, such as the time required to prepare patients. Further disadvantages include the fact that small bowel bleeding cannot be accessed through colonoscopy and that colonoscopy is relatively ineffective when performed in patients with significant bleeding without bowel preparation. If bleeding cannot be stopped by therapeutic colonoscopy, transarterial embolization is the next line of therapy to control hemostasis. As with upper gastrointestinal bleeding, transarterial embolization is the first-line therapy for patients with lower gastrointestinal bleeding. The efficacy of transarterial embolization in the treatment of acute gastrointestinal bleeding when medical or endoscopic techniques

are inadequate has been demonstrated in several large studies. For both upper and lower gastrointestinal bleeding, surgery is usually reserved as a last-line treatment for patients whose bleeding has failed to respond to previous treatments [25-28].

In angiography, vascular access is usually obtained by transfemoral catheterization with a 4- or 5-Fr catheter and sheath. Diagnostic visceral arteriography, which includes angiography of the celiac trunk, superior mesenteric artery, and inferior mesenteric artery, is performed to examine the suspected vascular region, and then a microcatheter is inserted coaxially for superselective cannulation of the bleeding artery. In upper gastrointestinal bleeding, the source of the bleeding is usually identified by endoscopy. Therefore, angiography is most often performed only as a precursor to transcatheter embolotherapy. Angiograms are considered positive when they show direct angiographic evidence of active GI bleeding (eg, contrast medium extravasation) or indirect angiographic evidence of bleeding (eg, pseudoaneurysm). For embolization, the appropriate position of the catheter is selected, and transcatheter vessel occlusion is performed with embolized material. Embolization is carried out as selectively as possible, the catheter is technically possible near the site of bleeding. The goal is to achieve proximal and distal control of the bleeding lesion (embolization of both inflow and outflow vessels) to reduce the risk of recurrent bleeding via collateral circulation. Embolization will continue to the occlusive angiographic endpoint without antegrade arterial blood flow in the embolized artery. Post-embolization arteriography is performed to confirm completion of the procedure. The most common embolic material used to treat upper gastrointestinal bleeding is a fibrous platinum microcoil, which is usually placed in the bleeding artery distally proximal to the angiographic position [29-33].

Extravasation of the contrast medium is stopped and complete occlusion of the bleeding vessel occurs. Coiling of the gastroduodenal artery from the celiac axis may be inadequate because the gastroduodenal artery may be fed by collateral branches of the superior mesenteric artery. A "sandwich" technique has been proposed, in which the gastroduodenal artery is looped distally proximally. Sandwich occlusion can be used at the level of the gastroduodenal artery with the catheter directed toward the origin of the right gastroepiploic artery, and when the catheter is removed the coils are inserted into the proximal gastroduodenal artery. Complete embolization of the gastroduodenal artery, including proximal and distal embolization and exclusion of its two side branches, is the technical end point [34-35].

Selective superior mesenteric arteriography is performed after embolization to ensure that the bleeding site is not secured. If extravasation is detected, superselective catheterization of the lower pancreatic duodenal artery and side branch is performed with a microcatheter. Over the past 10 years, significant improvements in this technique have made superelective embolization a safer procedure by minimizing the risk of intestinal ischemia.

The development of sophisticated rotating wires and coaxial microcatheters, along with advances in digital fluoroscopic imaging, now allow for more precise vascular interventions. In a recent report, transarterial embolization for upper gastroin-

testinal bleeding was associated with a high technical success rate (93%) and a minimal complication rate (9%). In addition, a recently published international consensus recommendation considers transarterial embolization as an alternative therapy for the treatment of upper gastrointestinal bleeding in patients who have failed an endoscopic hemostatic procedure or who have recurrent bleeding [36-38].

Transarterial embolization for the treatment of lower GI bleeding was first introduced in 1974 and involved the non-elective injection of an autologous clot. In 1977, Gelfoam and Oxycel injection were described for embolization of diverticular bleeding. Although injection of autogal clot or gelatin sponge has been shown to achieve hemostasis, these early embolization techniques were characterized by high rates of intestinal infarction. The development of coaxial microcatheters has increased interest in the use of embolization to control lower GI bleeding. The use of a microcatheter, delivered via a 4- or 5-wire guide catheter, to a specific margin near the bleeding site in the arteries or vas rectus can be obtained to remove embolic material. Because this technique is superselective, the risk of intestinal infarction is significantly lower than nonselective embolic techniques or vasopressin infusion, and there are no vasopressin-related systemic side effects. In addition, the risk of bleeding from collateral vessels is reduced as embolic material is delivered to the site of bleeding [39-40].

The most common embolic materials for lower GI bleeding, used alone or in combination, are microcoil, polyvinyl alcohol (PVA) particles, and gelatin sponge. Microcoils are persistent embolic agents that can be superselectively placed near the bleeding site and are easily identified under fluoroscopy. However, due to the small caliber of the target vessels, positioning these coils correctly can be difficult. Coils can back out of small vessels and provoke ischemia if they enter a large feeding vessel. PVA is a permanent embolic agent that is less selective than microcol. The basis for PVA embolization with respect to flow is that PVA particles preferentially flow to the area of least resistance (ie, the bleeding site). Defrin et al demonstrated in 10 patients that lesions inaccessible by superelective catheterization could be safely embolized by flow-directed PVA embolization. However, a consensus on the optimal PVA particle size for embolization in lower GI bleeding has not yet been reached. Previous reports have recommended a PVA particle size of 300 to 500 μm, as early animal studies have shown that smaller particles may be associated with a higher risk of intestinal ischemia. The choice of embolic agent in relation to the characteristics of the bleeding vessel is important, but which embolic agent is best among coils, cyanoacrylate glue, gelatin sponge, and calibrated particles remains a matter of debate. In our department, microcoil, 1000 μm gelatin sponge particles, and cyanoacrylate glue are used to treat acute GI bleeding. Unlike PVA, the gelatin sponge is a temporary embolic agent that allows recanalization of the vessel from a few days to a few weeks. If the angiogram is negative for active bleeding, empiric embolization is performed based on discussion with the gastroenterologist or surgeon. For empiric embolization, if endoscopy showed that the bleeding source was located in the proximal stomach, the left gastric artery was selectively embolized. If endoscopy shows that the source

of bleeding is in the distal stomach or duodenum, the gastroduodenal artery, the right gastroepiploic artery, the pancreaticoduodenum, or all three are selectively embolized. If endoscopic intervention fails to control bleeding, radioplaque clips are positioned as guides to the bleeding site via colonoscopy and transarterial embolization is performed [41-42].

Bleeding, deposition of coils is guided by endoscopy with pre-placed hemoclips. As a result, angiography and embolization of vessels causing GI bleeding have been gradually adopted and have revolutionized the management of lower GI bleeding.

Complications associated with embolization include angiography itself (eg, hematoma, arterial thrombosis, dissection, embolism, pseudoaneurysm) as well as intestinal infarction. Early transcatheter intervention involved vasopressin infusion, but the high rate of rebleeding and high complication rates led to its reduced use. Higher rates of complications and bleeding have been described in patients treated with vasopressin. Although the first embolic techniques improved hemostasis, their use was limited by the high incidence of intestinal infarction. Until the advent and development of microcatheter technology, transarterial embolization became a safer, more effective method for managing gastrointestinal bleeding. Improvements in microcatheter systems have enabled more selective delivery of embolic material near bleeding sites; This overcame the systemic side effects of vasopressin and resulted in a reduced risk of intestinal infarction and vascular bleeding. Recently, many reports have suggested that superelective embolization for the management of gastrointestinal bleeding rapidly stops bleeding with minimal risk of ischemia. However, the risk of ischemia after embolization is increased in patients with a history of surgical intervention in the same area or when the therapeutic intervention involves embolic agents that may advance into the vascular bed. Such agents include liquids (for example, fabric adhesives such as cyanoacrylate) or very small particles [43-44].

In stable patients, multidetector computed tomography (MDCT) imaging is a useful tool to identify bleeding sites and assess the anatomical structure of the gastrointestinal tract, thereby allowing for more targeted intervention. Scintigraphy of red blood cells has a sensitivity and specificity of more than 90%; However, its simulated resolution does not provide an accurate diagnosis. Computed tomography angiography (CTA) is also used (sensitivity up to 86%) in the diagnosis of acute GI bleeding and can be used to pinpoint the location and etiology of bleeding and thus direct further management. A positive MDCT angiogram may be useful in selecting patients suitable for rapid targeted embolization. Visualization of active extravasation of contrast medium in the gastrointestinal tract requires careful attention to technique, including the use of fine collimation, rapid administration of contrast medium, and appropriate scan times. Additionally, multimodal reconstruction and three-dimensional imaging are useful in determining the exact source of bleeding. Although further studies are needed to determine which course of action is best when bowel preparation is not possible, CTA may be useful in this situation to identify the site of bleeding [18, 34, 45].

The idea of embolization of duodenal bleeding as an alternative use of surgery belongs to Roche (1972). Since then, arterial embolization has been considered as an effective diagnostic and surgical method

Common causes of duodenal bleeding:

– arteriovenous malformation;
– visceral aneurysms/pseudoaneurysm;
– angiodysplasia;
– aortoenteric fistula;
– hemophilia;
– intestinal diverticula;
– inflammatory bowel disease (ulcer disease);
– benign anorectal lesions.

The main causes of duodenal bleeding (figure1).

Frequency of duodenal bleeding: 375 cases per 100,000 population. Acute case of bleeding-75%. 70% of patients are >65 years old. Recurrent bleeding-25%. Mortality is 19-40%. Ratio male 2:1 female.




Fig. 1 Transcatheter arterial embolization for acute nonvariceal upper gastrointestinal bleeding: Indications, techniques and outcomes.

Clinical manifestations of duodenal bleeding: hematemesis, color brownish vomit, melena, anemia, tachycardia (if blood loss > 500 ml), hypotension (if blood loss > 500 ml), systemic shock (if blood loss is more than 15% of circulating blood volume).

Diagnosis by fibrogastroscopy (figure 2).




Fig. 2 Fibrogastroscopy technics.

Angiography as a method of determining bleeding and result. (figure 3).



Fig. 3 Angiography as a method of determining bleeding and result.

The task:

– To determine indications for endovascular occlusion in ulcer patients complicated by gastroduodenal bleeding.

– The technique of endovascular interventions should be perfected.

– To evaluate the effectiveness of the endovascular occlusion method compared to the surgical method of treatment in high-risk patients.

– The tactics of surgical treatment of the mentioned patients should be developed based on the use of the endovascular occlusion method.

Forrest's classification during endoscopy: IA; IB; IIA; IIB; IIC; III.

Risk of rebleeding according to the Forrest classification (figure 4).




Fig. 4 Risk of rebleeding according to the Forrest classification.

Testimonials. Low-risk patients - surgical treatment. And high-risk patients - endovascular surgery. Recurrent bleeding after surgery - endovascular surgery.

The diagnosis and management of gastrointestinal (GI) bleeding are complicated and requires a multidisciplinary approach involving gastroenterologists, interventional radiologists, and surgeons.

Testimonials. With active bleeding (requiring 4 units of blood transfusion in 24 hours). With hemodynamic instability (low systolic pressure, pulse 100 or more, hypovolemic shock) - who did not have satisfactory results during endoscopic hemostasis, in this case it will be important to calculate the Rokall score. A high risk of rebleeding should be used in the Forrest classification. Forrest 1A, Forrest 1B Forrest 2A and Forrest 2B. Co-morbidities that aggravate the patient's medical history.

Graphical comparison of mean age of TAE and surgical groups in included studies (figure 5).




Fig. 5 Graphical comparison of mean age of TAE and surgical groups.

Material and methods**:** From March 2019 to December 2024, 40 patients were embolized during duodenal bleeding.

These patients were divided into the following groups:

– massive, active bleeding;
– recurrence of bleeding in clinic and endoscopy was unsuccessful;
– unstable hemodynamics and solid hemostasis could not be achieved during endoscopy;
– failed to evacuate stomach contents and failed to see a bleeding ulcer;
– high risk of bleeding recurrence (Forest classification);
– elderly and patients burdened with co-morbidities, with whom operative intervention represents a high risk of lethality.

Embolization performed according to Forrest's classification in 40 patients (figure 6).




Fig. 6 Embolization performed according to Forrest's classification

Embolization methods: blind- embolization; empirical embolization.

Technical aspects of embolization: radial approach, femoral artery approach.

Factors influencing the choice of embolic agent: angiographic conclusion, vascular anatomy, vascular size, desired level of vascular occlusion, temporary or permanent occlusion is preferred, catheter position, operator experience.

Embolization techniques (figure 7).

Case description. The patient is 78 years old**.** Pain in the chest**.** Marked respiratory failure**.** Troponin 2.6 ug/L**.** ST-elevation on the cardiogram.

Embolization Sendvich technique (Figure 8).

Study results: Technical success in 39 patients (97.5%). Technical failure in 1 patient (2.5%).

Embolization results obtained on 40 patients (Figure 9).

Angiography results (Figure 10).




Fig. 7 Embolization techniques.




Fig. 8 Embolization Sendvich technique.




Fig. 9 Embolization results.

The following criteria are used to evaluate the effectiveness of endovascular hemostasis. Technical success - interruption of blood flow in the embolization zone. Clinical success - correction of bleeding recurrence and stabilization of hemodynamics. Unsuccessful embolization.

Studies show that coagulopathy is significantly associated with mortality after failed embolization.




Fig. 10 Angiography results.

The risk of rebleeding after successful embolization is three times greater in patients with coagulation problems, and for the same reason there is a 10-fold greater risk of death compared to patients with normal coagulation.

Conclusion. Improvements in catheter technology, development of more compatible embolization devices, and expansion of embolization techniques have led to

angiography and embolization for the treatment of upper and lower gastrointestinal bleeding. Transcatheter embolization therapy for the treatment of acute GI bleeding is a safe procedure with high technical performance and clinical success, but it should be reserved as a treatment option for patients who have failed endoscopic and medical management. MDCT imaging is a useful tool for identifying the site of bleeding and evaluating the anatomical structure of the gastrointestinal tract in stable patients. Close working relationships between interventional radiologists, gastroenterologists, and diagnostic radiologists are essential for the optimal management of patients with GI bleeding. Endovascular embolization dramatically reduces the mortality rate in high-risk patients who require open surgery after failed endoscopy, further studies are needed to fully address these objectives.

Acknowledgments. Declaration of Interest Statement. We confirm that the manuscript has been read and approved by all named authors and that there are no other persons who satisfied the criteria for authorship but are not listed. We further confirm that the order of authors listed in the manuscript has been approved by all of us. Conflict of interest-None.

References:

1. Cherian M.P., Mehta P., Kalyanpur T.M. et al. Arterial interventions in gastrointestinal bleeding. Semin Intervent Radiol, 2009; 26: 184-195.
2. Sheth R., Someshwar V., Warawdekar G. Treatment of acute lower gastrointestinal hemorrhage by superselective transcatheter embolization. Indian J Gastroenterol, 2006; 25: 290-294.
3. Rozycki G.S., Tremblay L., Feliciano D.V. et al. Three hundred consecutive emergent celiotomies in general surgery patients. Ann Surg, 2002; 235: 681-689.
4. Pennoyer W.P., Vignati P.V., Cohen J.L. Management of angiogram positive lower gastrointestinal hemorrhage: long term follow-up of non-operative treatments. Int J Colorectal Dis, 1996; 11: 279-282.
5. Schuetz A., Jauch K.W. Lower gastrointestinal bleeding: therapeutic strategies, surgical techniques and results. Langenbecks Arch Surg, 2001; 386: 17-25.
6. Mirsadraee S., Tirukonda P., Nicholson A. et al. Embolization for non-variceal upper gastrointestinal tract haemorrhage: a systematic review. Clin Radiol, 2011; 66: 500-509.
7. Bjorkman D.J., Zaman A., Fennerty M.B. et al. Urgent vs. electiveendoscopy for acute non-variceal upper GI-bleeding: an effectiveness study. Gastrointest Endosc, 2004; 60: 1-8.
8. Church N.I., Palmer K.R. Diagnostic and therapeutic endoscopy. Curr Opin Gastroenterol, 1999; 15: 504.
9. Yap F.Y., Omene B.O., Patel M.N. et al. Transcatheter embolotherapy for gastrointestinal bleeding: a single center review of safety, efficacy, and clinical outcomes. Dig Dis Sci, 2013; 58: 1976-1984.
10. DeBarros J., Rosas L., Cohen J. et al. The changing paradigm for the treatment of colonic hemorrhage: superselective angiographic embolization. Dis Colon Rectum, 2002; 45: 802-808.
11. Lewis B.S. Small intestinal bleeding. Gastroenterol Clin North Am, 1994; 23: 67-91.
12. Jenson D.M., Machicado G.A., Jutabha R., Kovacs T.O. Urgentcolonoscopy for the diagnosis and treatment of severe diverticular hemorrhage. N Engl J Med, 2000; 342: 78-82.
13. Zuccaro G. Management of the adult patient with acute lower gastrointestinal bleeding. Am J Gastroenterol, 1998; 93: 1202-1208.

14. Angtuaco T.L., Reddy S.K., Drapkin S., Harrell L.E., Howden C.W. The utility of urgent colonoscopy in the evaluation of acute lower gastrointestinal tract bleeding: a 2-year experience from a single center. Am J Gastroenterol, 2001; 96: 1782-1785.
15. Bloomfield R.D., Rockey D.C., Shetzline M.A. Endoscopic therapy of acute diverticular hemorrhage. Am J Gastroenterol, 2001; 96: 2367-2372.
16. Eisen G.M., Dominitz J.A., Faigel D. et al. American Society for Gastrointestinal Endoscopy. Standards of Practice Committee: an annotated algorithmic approach to acute lower gastrointestinal bleeding. Gastrointest Endosc, 2001; 53: 859-863.
17. Zuckerman G.R., Prakash C. Acute lower intestinal bleeding. Part I: clinical presentation and diagnosis. Gastrointest Endosc, 1998; 48: 606-617.
18. Fisher L., Krinsky M.L., Anderson M.A. et al. The role of endoscopy in the management of obscure GI bleeding. Gastrointestinal Endoscopy, 2010; 72: 471-479.
19. Funaki B., Kostelic J.K., Lorenz J. et al. Superselective microcoil embolization of colonic hemorrhage. Am J Roentgenol, 2001; 177: 829-836.
20. Funaki B. Superselective embolization of lower gastrointestinal hemorrhage: a new paradigm. Abdom Imaging, 2004; 29: 434-438.
21. Barkun A.N., Bardou M., Kuipers E.J. et al. International consensus recommendations on the management of patients with nonvariceal upper gastrointestinal bleeding. Ann Intern Med, 2010; 152: 101-113.
22. Weldon D.T., Burke S.J., Sun S., Mimura H., Golzarian J.: Interventional management of lower gastrointestinal bleeding. Eur Radiol, 2008; 18: 857-867.
23. Mirsadraee S., Tirukonda P., Nicholson A. et al. Embolization for non-variceal upper gastrointestinal tract haemorrhage: a systematic review. Clin Radiol. 2011; 66(6): 500-9.
24. Charbonnet P., Toman J., Buhler L. et al: Treatment of gastrointestinal hemorrhage. Abdom Imaging, 2005; 30: 719-726.
25. Gocha Chankseliani, Avtandil Girdaladze, Omar Gibradze et al. The features of endovascular surgery for duodenal ulcer bleeding. RS Global - World Science [Internet]. 2023. Dec.11;(4(82). Available from: https://rsglobal.pl/index.php/ws/article/view/2690
26. Gocha Chankseliani, Avtandil Girdaladze, Omar Gibradze et al. The manifestation of some key issue aspects of peculiarities of endovascular surgery based on endovascular embolization for bleeding from gastro-duodenal ulcer and features of some essential pharmacotherapy. Conference proceedings: The First International Scientific-Practical Conference In Health Innovations & Research: Prognosis, Achievement and Challenges (december 15-16, 2023). Tallinn, Estonia-2023. P:14-15.
27. G. Chankseliani, A. Girdaladze, O. Gibradze et al. The manifestation of some key issue aspects of peculiarities of endovascular surgery based on endovascular embolization for bleeding from gastro-duodenal ulcer and features of some essential pharmacotherapy. Southern caucasus scientific journals; Ambiance in life; International scientific journal in medicine; Journal of behavioral medicine, refereed and reviewed journal; VOL. 15, ISSUE 01, 2023; 15.01.2023. Tbilisi-Georgia-2023. P: 20-38.
28. Gocha Chankseliani, Avtandil Girdaladze, Omar Gibradze et al. The scientific discussion singularities of endovascular surgery and endovascular embolization for bleeding from gastro-duodenal ulcer and treatment of complications. Proceedings book; Proceedings book of 5 baskent international conference on multidisciplinary studies; December 21-22, 2023, Ankara, Turkey-2023. P:1041-1059.
29. Talley N.J., Vakil N., for the Practice Parameters Committee of the American College of Gastroenterology. Guidelines for the management of dyspepsia. Am. J. Gastroenterol. 2005 Oct. 100(10):2324-37.
30. Tajima A., Koizumi K., Suzuki K. et al. Proton pump inhibitors and recurrent bleeding in peptic ulcer disease. J Gastroenterol Hepatol. 2008 Dec. 23 suppl 2: S237-41.

31. McConnell D.B., Baba G.C., Deveney C.W. Changes in surgical treatment of peptic ulcer disease within a veteran's hospital in the 1970s and the 1980s. Arch Surg. 1989 Oct. 124(10):1164-7.
32. Gisbert J.P., Calvet X., Cosme A. et al, for the H. pylori Study Group of the Asociacion Espanola de Gastroenterologia (Spanish Gastroenterology Association). Long-term follow-up of 1,000 patients cured of Helicobacter pylori infection following an episode of peptic ulcer bleeding. Am J Gastroenterol. 2012 Aug. 107(8):1197-204.
33. Tarasconi A., Coccolini F., Biffl W.L. et al. Perforated and bleeding peptic ulcer: WSES guidelines. World J Emerg Surg. 2020. 15:3.
34. Berne T.V., Donovan A.J. Nonoperative treatment of perforated duodenal ulcer. Arch Surg. 1989 Jul. 124(7):830-2.
35. Donovan A.J., Berne T.V., Donovan J.A. Perforated duodenal ulcer: an alternative therapeutic plan. Arch Surg. 1998 Nov. 133(11):1166-71.
36. Wangensteen O.H. Non-operative treatment of localized perforations of the duodenum. Proc Minn Acad Med. 1935. 18:477-80.
37. Strand D.S., Kim D., Peura D.A. 25 years of proton pump inhibitors: a comprehensive review. Gut Liver. 2017 Jan 15. 11(1):27-37. .
38. Kajihara Y., Shimoyama T., Mizuki I. Analysis of the cost effectiveness of using vonoprazan-amoxicillin-clarithromycin triple therapy for first-line Helicobacter pylori eradication. Scand J Gastroenterol. 2017 Feb. 52(2):238-41.
39. Ng J.C., Yeomans N.D. Helicobacter pylori infection and the risk of upper gastrointestinal bleeding in low dose aspirin users: systematic review and meta-analysis. Med J Aust. 2018 Sep 1. 209(7):306-11.
40. Mirabella A., Fiorentini T., Tutino R. et al. Laparoscopy is an available alternative to open surgery in the treatment of perforated peptic ulcers: a retrospective multicenter study. BMC Surg. 2018 Sep 25. 18(1):78.
41. Havens J.M., Castillo-Angeles M., Nitzschke S.L., Salim A. Disparities in peptic ulcer disease: A nationwide study. Am J Surg. 2018 Dec. 216(6):1127-8.
42. Satoh K., Yoshino J., Akamatsu T. et al. Evidence-based clinical practice guidelines for peptic ulcer disease 2015. J Gastroenterol. 2016 Mar. 51(3):177-94.
43. Lau J.Y., Sung J., Hill C. et al. Systematic review of the epidemiology of complicated peptic ulcer disease: incidence, recurrence, risk factors and mortality. Digestion. 2011. 84(2):102-13.
44. Sonnenberg A. Temporal trends and geographical variations of peptic ulcer disease. Aliment Pharmacol Ther. 1995. 9 suppl 2:3-12.
45. Mohamed W.A., Schaalan M.F., Ramadan B. The expression profiling of circulating miR-204, miR-182, and lncRNA H19 as novel potential biomarkers for the progression of peptic ulcer to gastric cancer. J Cell Biochem. 2019 Aug. 120(8):13464-77.
46. Sanaii A., Shirzad H., Haghighian M. et al. Role of Th22 cells in Helicobacter pylori-related gastritis and peptic ulcer diseases. Mol Biol Rep. 2019 Dec. 46(6):5703-12.

Розділ 10

МЕТОДИЧНІ АСПЕКТИ ВИКЛАДАННЯ ФАРМАКОЛОГІЇ, КЛІНІЧНОЇ ФАРМАКОЛОГІЇ ТА КЛІНІЧНОЇ ФАРМАЦІЇ В УКРАЇНІ ТА СВІТІ

FORMATION OF THERAPEUTIC THINKING OF PHARMACIST STUDENTS IN THE TEACHING OF PHARMACOLOGY AND CLINICAL PHARMACY

Filipets N. D.[1] , Gerush O. V.[2] , Filipets O. O.[3]
Bukovinian State Medical University, Chernivtsi, Ukraine
[1] Chair of Pharmacology, [2] Chair of Pharmacy, [3] Chair of Nervous Diseases, Psychiatry and Medical Psychology named after S.M. Savenko

The pharmaceutical industry today is characterized by growth in most aspects. At the same time, the volume of information about drugs is increasing, and, accordingly, the issue of rational pharmacotherapy is relevant. The possibilities of pharmacists today are not only to influence the choice of drugs from the position of marketing in the pharmaceutical market, which is extremely complex compared to most other industries. According to the modern strategy for increasing the role of the pharmacist in the health care system, the main concept of the clinical pharmacists professional activity is the assessment of the effectiveness and safety ofpharmacotherapy for a specific patient.

The purpose of the work is to highlight methodological approaches to the formation of the therapeutic thinking of a pharmaceutical worker when teaching pharmacology and clinical pharmacy for the professional performance of the main tasks in future activities related to the prevention of diseases, preservation and strengthening of human health, ensuring the rational use of medicinal products.

The professional activity of a clinical pharmacist, in addition to providing the population with pharmaceutical preparations and medical goods, involves participation in the development of local drug formularies in health care institutions; determination of advantages and disadvantages of drugs of various pharmacological groups taking into account their chemical, physicochemical, biopharmaceutical, pharmacokinetic, pharmacodynamic and pharmacoeconomic features; participation in clinical research and study of bioequivalence of drugs; monitoring of side effects or ineffectiveness of drug therapy; solving problems of polypharmacy. The performance of such professional tasks is ensured by clinical and pharmaceutical knowledge and competences acquired, in particular, during the study of pharmacology and clinical pharmacology.

At the same time, the basic knowledge and skills should be learned and used as the student pharmacist, thanks to interdisciplinary integration, develops skills in the practical application of rational therapy. Here, an important component of competence is the students ability to find useful and reliable information. The student should be involved in the search, collection and analysis of information in the main databases of evidence-based medicine. Implementation of evidence-based medicine is a necessary component of effective pharmacotherapy. Such a process forms a critical attitude to the flow of medical and pharmaceutical information, the ability to identify false information and find new information in reliable scientific sources, which, in the future, is important for making professional decisions. The function of a

pharmacist is to inform the population and doctors about new drugs in comparison with known domestic and foreign drugs. The accumulation, systematization, and dissemination of pharmaceutical and pharmacology information requires the ability to use evidence of the most effective and efficient methods or interventions to support decision-making related to patient health.

Basic pharmacological knowledge of pharmacokinetics and pharmacodynamics is necessary for the development of rational and optimal therapeutic plans. Such competencies ensure the clinical pharmacists participation in providing highly qualified medical care by providing recommendations to doctors and patients about the most effective drugs. Therefore, situational tasks with the main characteristics of the pathological process for determining the treatment drug remain an effective means of learning and knowledge control. It is worth noting that here the teacher should provide pharmacist students with an explanation of the mechanisms of disease development and directions of pathogenetic correction in order to choose the most effective drugs.

Therefore, the clinical pharmacist, together with the doctor, selects drugs and determines their individual doses for patients depending on the main disease and accompanying pathology. However, teaching in class and practicing medication administration in clinical settings is easier in small groups of students. Certain problematic issues when working in small groups can be the choice of the most effective drugs of a certain pharmacological group, their combinations, in particular in fixed doses, analogues, etc. At the same time, the selection of the least toxic drugs or combinations of drugs is a mandatory component of the pharmacists activity. Side effects can be illustrated by tables, diagrams, videos illustrating pharmaceutical / pharmacodynamic incompatibility of drugs or mechanisms of organotoxicity as a result of incorrect selection of drugs.

The professional competence of a pharmacist is the prediction of possible complications with the simultaneous use of several drugs - polypharmacy. Raising awareness of the consequences of polypharmacy is an integral part of the learning process. When covering issues of polypharmacy, it is advisable to use concepts with examples of drugs in a specific clinical case. Feedback and summary summaries allow effective understanding of the consequences of polypharmacy and measures to prevent them.

Finally, in order to acquire the competence of effective and safe selection of drugs, future pharmacists must acquire knowledge of pharmacology and clinical pharmacology of basic drugs and general knowledge about diseases, must master the skills of prescribing drugs taking into account rational routes and modes of administration, incompatibility of drugs, side effects of drugs. Pharmacy is a branch of the health care sector that plays an important role in the practical implementation of tasks aimed at improving the health of the population and providing pharmaceutical care to patients. Modern strategies, such as collaboration between pharmacists and doctors and nurses, as well as the provision of patient-centered healthcare based on teamwork, ensure timely, safe and effective treatment.

ДОСВІД ЗАСТОСУВАННЯ ДИСТАНЦІЙНИХ ФОРМ НАВЧАННЯ З АКЦЕНТОМ НА ФОРМУВАННЯ КЛІНІЧНОГО МИСЛЕННЯ СУЧАСНОГО МАГІСТРА ФАРМАЦІЇ

Крайдашенко О. В., Свинтозельський О. О., Михайлик О. А.
Запорізький державний медико-фармацевтичний університет, м. Запоріжжя, Україна
Кафедра клінічної фармакології, фармації, фармакотерапії і косметології
krayd@ukr.net

Вступ. Жорсткі заходи та умови військового стану призвели до глобальних змін в освітньому процесі та примусили навчальні заклади терміново впроваджувати дистанційні форми навчання, як єдино можливі.

Мета. Визначити можливості використання дистанційних форм навчання при викладанні дисциплін «Клінічна фармація та фармацевтична опіка» при формуванні клінічного мислення у студентів 4 та 5 курсів фармацевтичних факультетів та розробити можливі шляхи їх удосконалення.

Матеріали та методи. Матеріал для дослідження – досвід організації та проведення навчального процесу з використанням дистанційних та змішаних форм навчання за останні 5 років, звітна й інша облікова документація, анкетування студентів. Під час роботи використали методи контент-аналізу, синтезу, групування, дедукції, логіки та узагальнення.

Результати. За час вимушеного карантину та військового стану співробітниками кафедри була налагоджена належна співпраця зі студентами. Подання теоретичного матеріалу, лекцій, проведення практичних занять, підсумкового контролю втілене при розробці та впровадженні в педагогічний процес on-line курсів, але відсутність можливості активного спілкування студентів з тематичними хворими, що є основою у формуванні клінічного мислення, потребувало втілення нових форм дистанційної освіти з максимальним використанням доступних ресурсів.

Вирішенням проблеми недостатнього спілкування студентів з хворими при викладанні дисциплін «Клінічна фармація та фармацевтична опіка» є використання фармацевтичного симулятора (віртуальна аптека) «Pharmacy Simulator™», який дозволяє опрацьовувати навички збору скарг пацієнта, аналіз його захворювання, лікарський анамнез, виключення загрозливих станів при наданні фармацевтичної опіки, вибору раціональної лікарської форми препарату, навички ділового спілкування відповідно до затверджених протоколів. Це дозволить значно покращити якість підготовки клінічного фармацевта. Загалом кафедра має 26 акаунтів, які дозволяють одночасно отримати доступ до платформи декільком групам студентів.

Висновки. В умовах військового стану використання в безперервному процесі освіти дистанційних форм навчання за допомогою комп'ютерних навчальних on-line сервісів та використання симулятора «Pharmacy Simulator™», повинно отримати свою роль в удосконаленні навчального процесу.

ФАРМАЦЕВТИЧНА БЕЗПЕКА З ПОГЛЯДУ ФАРМАЦЕВТИЧНОЇ ТЕРМІНОСИСТЕМИ

Панькевич О. Б., Громовик Б. П.

Львівський національний медичний університет імені Данила Галицького, м. Львів, Україна

Кафедра організації і економіки фармації

pankevych.lviv@gmail.com, hromovyk@gmail.com

Вступ. Повномасштабна російська агресія проти України загострила питання фармацевтичної безпеки як запоруки національної безпеки. Відтак, поняття фармацевтичної безпеки потребує поняттєвого висвітлення.

Мета. Метою цієї роботи є окреслення термінологічного поля фармацевтичної безпеки як складової частини фармацевтичної терміносистеми.

Матеріали та методи. Вихідними матеріалами були статті з електронних баз даних (Google Scholar & PubMed) та посилання, що містилися в цих статтях. Використано два взаємодоповнювальні методи дослідження в термінології: статистичний метод, який полягає в означенні наявності тієї чи іншої мовної реалії безвідносно до її правильності, а також аналітичний метод, який уможливлює з'ясування наукової обґрунтованості та доцільності конкретного терміну (http://dspace.nbuv.gov.ua/handle/123456789/51281).

Результати. З'ясовано, що дефініція фармацевтичної безпеки у вітчизняній літературі слабо опрацьована, а наявні підходи до цього поняття [https://www.lute.lviv.ua/fileadmin/www.lac.lviv.ua/data/pidrozdily/Aspirantura/Rady/Spec_vchena_rada/Dysertacii/2021_04/Datsko_Disert.pdf; https://apir.org.ua/wp-content/uploads/2016/12/Voronina7.pdf; https://doi.org/10.13165/PSPO-20-25-01] важкі для сприйняття внаслідок багатослівності. Зважаючи на зазначене вище, ми пропонуємо під фармацевтичною безпекою розуміти невіддільну частину національної безпеки, що характеризується таким станом розвитку фармації, при якому забезпечується задоволення потреб населення у рівному його доступі до якісної фармацевтичної допомоги. При цьому для кращого розуміння змісту досліджуваного терміну опрацьовано ще дві дефініції:

– політика фармацевтичної безпеки як загальні наміри й спрямованість розвитку фармацевтичної безпеки, офіційно сформульовані урядом держави у вигляді стратегії, концепції, плану чи доктрини на підставі розуміння реальних та потенційних загроз середовища фармацевтичної безпеки;

– культура фармацевтичної безпеки як складова частина її політики, яка базується на системі матеріальних та духовних індивідуальних і групових цінностей, що визначають прихильність і моделі поведінки на рівні особи, суспільства та держави щодо потенційної можливості прояву загрози для якості життя, пов'язаного зі здоров'ям в частині обігу лікарських засобів.

Висновки. Внаслідок проведеного дослідження уточнено дефініцію «фармацевтична безпека» та здійснено трактування двох термінів: «політика фармацевтичної безпеки» та «культура фармацевтичної безпеки».

ПРОПОЗИЦІЇ ТА ОБҐРУНТУВАННЯ ВВЕДЕННЯ ПЕВНИХ ВИБІРКОВИХ КОМПОНЕНТІВ ДО ОСНОВНИХ В ОСВІТНЬО-ПРОФЕСІЙНІЙ ПРОГРАМІ СПЕЦІАЛЬНОСТІ 226 «ФАРМАЦІЯ, ПРОМИСЛОВА ФАРМАЦІЯ»

Слєсарчук В. Ю., Кайдаш С. П., Соколова К. В.
Дніпровський державний медичний університет, м. Дніпро, Україна
Кафедра фармакології, загальної та клінічної фармації
Vladlenaslesarchuk@gmail.com

Вступ. Всестороннє формування знань, вмінь та навичок у студентів спеціальності «Фармація, промислова фармація» має важливе значення в освітньому процесі. Викладання вибіркових компонентів освітньої програми, таких як «Побічна дія ліків» та «Лабораторна діагностика» поглиблюють знання здобувача освіти та розширюють кругозір освітянина у майбутній трудовій діяльності. На кафедрі фармакології, загальної та клінічної фармації Дніпровського Державного медичного університету дисципліна «Клінічна фармація з фармацевтичною опікою» викладається на 4-5 курсах здобувачам другого (магістерського) рівня вищої освіти спеціальності 226 «Фармація, промислова фармація». Дисципліна є профільною, як складова входить до інтегрованого іспиту «Крок-2» (17 % питань від загального обсягу іспиту). Метою викладання дисципліни «Клінічна фармація з фармацевтичною опікою» є підготовка спеціалістів-фармацевтів, які б володіли достатнім обсягом теоретичних знань та практичних навичок для проведення разом з лікарем роботи по забезпеченню максимально раціональної фармакотерапії хворого, а також для здійснення фармацевтичної опіки пацієнтів в аптеці.

Мета. Метою роботи стала демонстрація значимості вибіркових компонентів «Побічна дія ліків» та «Лабораторна діагностика» при вивчені основного освітнього компонента «Клінічна фармація з фармацевтичною опікою».

Результати. Дисципліна «Клінічна фармація з фармацевтичною опікою» інтегрується й базується на вивченні здобувачам вищої освіти блоку таких дисциплін як загальна фармакологія, медична біологія, фізіологія та патологічна фізіологія з основами патологічної анатомії, мікробіологія, фармакотерапія, лікарська токсикологія, лабораторна діагностика, побічна дія ліків та інші, які передують вивченню дисципліни або вивчаються одночасно.

Одними із основних завдань вивчення освітнього компоненту «Клінічної фармації з фармацевтичною опікою» є: вміння вибирати оптимальний лікарський засіб з високими клінічними ефектами та максимальною безпекою; вміння враховувати фактори ризику та ступінь клінічних проявів побічної дії лікарських засобів, їх комбінацій та взаємодію лікарських засобів при одночасному прийомі; вміння складати та реалізовувати план заходів щодо зменшення або попередження виникнення побічних ефектів лікарських засобів при призначенні їх хворим; засвоєння принципів оцінки клінічної ефективності та профілю безпеки застосування конкретних фармакологічних груп та лікарських препара-

тів; вміти інтерпретувати та контролювати показники лабораторного обстеження хворих у динаміці, які є результатом фармакотерапії та здійснювати таким чином фармакологічну опіку хворого. По суті, це є основними завданнями таких дисциплін як «Побічна дія ліків» та «Лабораторна діагностика», що є вибірковими освітніми компонентами в поточному навчальному плані.

Керуючись змінами до ЗУ «Про вищу освіту», що набрав чинності 16 серпня 2024 року, запроваджено додаткове регулювання – обсяг вибіркових компонентів освітньої програми - не менш як 10% загальної кількості кредитів ЄКТС. З огляду на можливість внесення змін до навчального плану та важливість таких вибіркових освітніх компонент як «Лабораторна діагностика» та «Побічна дія ліків» для базового вивчення основної профілюючої дисципліни «Клінічна фармація з фармацевтичною опікою», пропонується ці вибіркові освітні компоненти внести до переліку основних компонентів, що дасть змогу поглибити та розширити знання освітян.

Результатом таких змін мають стати поглиблення знань не лише клініко-фармакологічної характеристики окремих груп лікарських препаратів, принципів їх взаємодії, але й методів та критеріїв оцінки ефективності та безпеки фармакотерапії, зокрема впливу лікарських засобів на показники лабораторного обстеження хворих та запобігання хибного тлумачення змін лабораторних показників хворих під впливом ліків.

Висновки. Введення до основних компонентів таких дисциплін як «Побічна дія ліків» та «Лабораторна діагностика» має покращити показники результатів складання ліцензійних іспитів серед здобувачів вищої освіти спеціальності «Фармація, промислова фармація» та сприятиме поглибленню знань та формуванню умінь застосовувати отримані знання в своїй професійній діяльності майбутнім фахівцям.

МЕТОДИЧНІ ТА ТЕХНОЛОГІЧНІ АСПЕКТИ ВИКЛАДАННЯ ФАРМАКОЛОГІЇ

Горчакова Н. О., Шумейко О. В., Клименко О. В.
Національний медичний університет імені О.О. Богомольця
Кафедра фармакології
pharmacology@nmu.ua

Вступ. Колектив кафедри фармакології опанував класичні форми викладання предмету та почав активно впроваджувати тестові завдання, а також кейсові методики, що допомогає студентам більш глибоко зрозуміти класифікації, фармакокінетику і фармакодинаміку ліків. На кафедрі була видана монографія: «програмно-математичні технології в розробці та технології лікарських засобів», що спонукало колектив шукати нові підходи із застосуванням новітніх методичних технологічних підходів у викладанні фармакології.

Мета. Встановити нові методики і технологічні аспекти викладання фармакології.

Матеріали і методи. Результати аналізу вітчизняної і зарубіжної літератури щодо встановлення нових аспектів методичної роботи та технологічних і наочних прийомів викладання фармакології.

Результати. Значні досягнення в галузі фармакології і фармації дозволили не тільки розширити арсенал лікарських засобів, необхідних для опанування знань з фармакології на третьому курсі. В перелік необхідних препаратів включені засоби, узгоджені з клінічними та фармацевтичними кафедрами університету. Для проведення практичних занять були створені робочі зошити з лікарськими фармпрепаратами для кожного заняття, в які, крім необхідних теоретичних знань, включено завдання та медичну компетентність. Викладачі проводять заняття з максимальним залученням технічної бази, для кращого засвоєння та подачі матеріалу. На кафедрі організовані тематичні навчальні аудиторії: *«університетська аптека»,* інтерактивна фармакологічна студія з поглибленим вивченням нейрофармакології та лікарських засобів, що впливають на центральну нервову систему людини, *«фітофармакологія», «кардіофармакологія».* Кожен квадратний метр новоствореного освітнього простору надає корисну інформацію про повний цикл лікування захворювань: від технологічних етапів створення відповідних пігулок до визначення коефіцієнту корисної дії після їх застосування у конкретних клінічних випадках. Інтерактивні стенди дають можливість здобувачам освіти самостійно обрати монокомпонентний препарат або комбінацію лікарських засобів та відразу побачити ефект, який він спричинить. Подвійний коефіцієнт корисної дії таких стендів полягає в тому, що розсувні панелі дозволяють удвічі збільшити площу інформаційного поля, відображаючи анатомо-терапевтично-хімічну класифікаціюких засобів і механізми впливу ліків на різні ланки патогенезу захворювань. Візуальне відображення механізмів дії ліків дозволяє реалізувати сучасні освітні технології та наочно продемонструвати ефективність синергістичної дії препаратів, що полегшує запам’ятовування здобувачами освіти сфер їх клінічного застосування.

Висновки. Колектив кафедри вважає, що тільки нові завдання, методики, рекомендації, учбові колекції поліпшать опанування основ фармакології. Цьому буде сприяти застосування технологічних форм наочності матеріалу. Технології і методики удосконалення викладання фармакології продовжуються.

ПЕРСПЕКТИВИ РОЗРОБКИ ПРОФЕСІЙНОГО СТАНДАРТУ «ПРОФЕСІОНАЛ З КЛІНІЧНИХ ДОСЛІДЖЕНЬ»

Кузнєцова В. Ю., Федосов А. І., Файзуллін О. В.
Національний фармацевтичний університет,
м. Харків, Україна
Кафедра клінічної фармакології,
clinpharmacol_ipksph@nuph.edu.ua

Вступ. Наказом Міністерства охорони здоров'я України від 29.11.2023 р. № 2034 «Про внесення змін до наказу Міністерства охорони здоров'я України від 28 жовтня 2002 року № 385» у закладах охорони здоров'я запроваджено нову посаду – професіонал з клінічних досліджень.

Відповідно до змін законодавчої бази, що відбулися протягом останніх років, наявність необхідної професійної кваліфікації є однією з найважливіших умов працевлаштування. Створення в Україні Національної системи кваліфікацій є важливою умовою реалізації євроінтеграційних зусиль і передбачено Угодою про асоціацію з Європейським Союзом. Згідно статті 4 Кодексу законів про працю України та статті 34 Закону України «Про освіту» вимоги до професійної кваліфікації працівників визначаються професійним стандартом.

Галузевою радою з розроблення професійних стандартів у сфері охорони здоров'я було прийнято рішення про створення робочої групи з розробки професійного стандарту 2212.2 «Професіонал з клінічних досліджень», до складу якої увійшли представників від різних установ, і робота якої триває.

Мета. Метою цієї роботи є аналіз професійної діяльності спеціалістів, що залучені до клінічних досліджень, та визначення основних трудових функцій, професійних компетентностей, відповідних знань та вмінь, необхідних для здійснення трудової діяльності професіонала з клінічних досліджень.

Матеріали та методи. При виконанні даної роботи був використаний метод функціонального аналізу трудової діяльності.

Результати. Під час розробки професійного стандарту «Професіонал з клінічних досліджень» учасники робочої групи спираються на Методичні рекомендації щодо розроблення професійних стандартів, схвалені рішенням Національного агентства кваліфікацій №1 від 25.01.2023 р., та Практичні рекомендації для розробників професійних стандартів, які базуються на вимогах чинного законодавства.

Загальний порядок розробки, впровадження та використання професійних стандартів в Україні визначається Кодексом законів про працю України (ст. 1, 39, 39^{-1}), Законом України «Про освіту» (ст. 4^1, 4^2, 96) та Порядком розроблення, введення в дію та перегляду професійних стандартів, затвердженим постановою Кабінету Міністрів України від 31.05.2017 р. № 373 «Про затвердження Порядку розроблення, введення в дію та перегляду професійних стандартів» зі змінами, внесеними постановою Кабінету Міністрів України № 1453 від

27.12.2022 р. «Про внесення змін до постановою Кабінету Міністрів України від 31 травня 2017 р. № 373».

Згідно з «Порядком розроблення, введення в дію та перегляду професійних стандартів» розроблення професійних стандартів здійснюється розробником професійного стандарту разом із членами робочої групи, до якої обов'язково включаються представники роботодавців, їх організацій та об'єднань, профспілок, їх організацій та об'єднань, представники сфери освіти. До складу групи також можуть входити за згодою представники центральних і місцевих органів виконавчої влади, інші фахівці за видом економічної діяльності, для якого розробляється професійний стандарт.

Головними завданнями, що мають бути вирішені під час розробки професійного стандарту «Професіонал з клінічних досліджень» є формулювання основної мети професійної діяльності, правильне визначення місця та функціоналу нової професії, розмежування трудових функцій професіоналу з клінічних досліджень та інших спеціалістів в суміжних галузях, що залучені до проведення клінічних досліджень (лікар, фармацевт, лаборант тощо). З метою ефективного вирішення цих завдань, було досліджене та проаналізоване поле професійної діяльності спеціалістів, що залучені на різних етапах клінічних досліджень.

Висновки проведеного дослідження були сформульовані у вигляді трудових дій, трудових функцій, професійних компетентностей, відповідних знань та вмінь. Відповідно до Методичних рекомендації щодо розроблення професійних стандартів ці дані були внесені до опитувальних анкет, оскільки результати аналізу, що був проведений, обов'язково мають бути підтверджені або відкориговані в ході опитування досвідчених працівників сфери клінічних досліджень.

Іншим важливим завданням, що вирішується в ході розробки професійного стандарту «Професіонал з клінічних досліджень», є визначення порядку та форм підвищення кваліфікації за професією, що створює умови для забезпечення безперервного професійного розвитку.

Висновки. В ході розробки професійного стандарту «Професіонал з клінічних досліджень», сумісно з роботодавцями, проведено аналіз професійної діяльності спеціалістів, що залучені до проведення клінічних досліджень, та визначені основні трудові дії, трудові функції та компетентності, а також відповідні вимоги до знань, умінь і навичок, необхідних для вирішення широкого кола практичних завдань, що пов'язані з плануванням, організацією та проведенням клінічних випробувань лікарських засобів.

Наразі розробка перебуває на стадії підготовки робочою групою проєкта професійного стандарту «Професіонал з клінічних досліджень».

ДОСВІД КАФЕДРИ ОРГАНІЗАЦІЇ, ЕКОНОМІКИ ТА УПРАВЛІННЯ ФАРМАЦІЄЮ ІПКСФ НФАУ З ПРОВЕДЕННЯ РЕФЛЕКСІЇ ФАРМАЦЕВТІВ-ІНТЕРНІВ ПІД ЧАС ДИСТАНЦІЙНОГО НАВЧАННЯ

Артюх Т. О.

Інститут підвищення кваліфікації спеціалістів фармації Національного фармацевтичного університету, Кафедра організації, економіки та управління фармацією м. Харків, Україна

artiukh.ipksf@nuph.edu.ua

Вступ. Первинна спеціалізація шляхом навчання в інтернатурі є обов'язковою формою післядипломної підготовки випускників закладів вищої медичної (фармацевтичної) освіти, після закінчення якої присвоюється кваліфікація зі спеціальності «Фармація». Карантинні обмеження та воєнний стан в Україні обумовили необхідність реалізації освітньої складової підготовки фармацевтів-інтернів з використанням технологій дистанційного навчання, зокрема й для проведення рефлексії під час проведення навчальних занять.

Мета. Аналіз ефективності реалізації використання науково педагогічними працівниками технологій дистанційного навчання для здійснення рефлексії під час лекційних, семінарських та практичних занять для фармацевтів-інтернів в рамках курсу «Управління та економіка фармації» під час проходження ними первинної спеціалізації на етапі післядипломної підготовки зі спеціальності «Фармація» на кафедрі організації, економіки та управління фармацією Інституту підвищення кваліфікації спеціалістів фармації Національного фармацевтичного університету (ОЕУФ ІПКСФ НФаУ).

Матеріали та методи. Аналіз наукової літератури та нормативної документації, інформаційно-логічний аналіз, порівняння, систематизація та групування, метод польового дослідження та спостереження.

Результати дослідження. Дослідження реалізації використання науково педагогічними працівниками технологій дистанційного навчання для здійснення рефлексії під час лекційних, семінарських та практичних занять для фармацевтів-інтернів в рамках курсу «Управління та економіка фармації» під час проходження ними первинної спеціалізації на етапі післядипломної підготовки зі спеціальності «Фармація» на кафедрі ОЕУФ ІПКСФ НФаУ показало, що використання різних методів та інструментів для проведення рефлексії широко застосовується для самоаналізу знань і вчинків, їх значень та меж під час кожного виду занять з фармацевтами-інтернами, а також під час здійснення ними самостійної роботи для критичної оцінки власних здобутків самими фармацевтами-інтернами та своєчасної корекції досягнення ними програмних результатів навчання на шляху здобуття як професійних, так і загальних компетентностей. Крім того, науково-педагогічними працівниками кафедри ОЕУФ ІПКСФ НФаУ широко застосовуються цифрові інструменти та різні додатки для безпо-

середнього проведення рефлексії та організації її процесу, які легко вбудовуються в інтерактивні платформи, такі як Miro та платформи Zoom, та модульне об'єктно-орієнтоване динамічне навчальне середовище, яке називають також системою управління навчанням (LMS) Moodle (Modular Object-Oriented Dynamic Learning Environment), запроваджене та широко використоване в ІПКСФ НФаУ для підготовки фармацевтів-інтернів, зокрема в умовах широкомасштабного вторгнення в Україну агресора та неможливість проведення очних занять в аудиторіях через близкість розташування ІПКСФ НФаУ в місті Харків до території бойових дій та фронту.

Так, для аналізу участі у співпраці та динаміки емоційного стану окремих фармацевтів-інтернів у процесі дистанційного навчання, здатності учасників віртуальної спільноти до емпатії, а також подальшого генерування коригувальних впливів з боку викладача, на кафедрі ОЕУФ ІПКСФ НФаУ використовується наступний перелік педагогічних інструментів: тест особистості Айзенка; тест «Індекс міжособистісної реактивності» IMP Девіса; тест цільових орієнтацій Елліота та самозвіт емоційного стану.

Крім того, здійснення рефлексії настрою та емоційного стану здобувачів освіти уможливлюється комплексом прийомів та засобів навчання. Зокрема, можна зауважити, що вивчення та аналіз передового педагогічного досвіду свідчить про дієвість рефлексивної діяльності наприкінці заняття, організованої на основі моделі «BOFF»: Behaviour (поведінка) – «Що відбувалося?» «Деталізуйте, що ви робили» (факти, опис подій). Outcome (результат) – «Який результат отримали?» «Чого ви навчилися?» «Що ви зрозуміли?». Feelings (почуття) – «Що відчували в процесі заняття?». Future (майбутнє) – «Як цей досвід можна використати в майбутньому?». Для здійснення рефлексії здобувачами освіти педагогічної діяльності та взаємодії наприкінці онлайн заняття використані прийоми формування критичного мислення, як-от: «Фішбоун», «Рефлексивна мішень», «Синквейн», «Незакінчене речення», «Стратегія 3-2-1» (задайте 3 питання, дайте 2 зауваження, залиште 1 пропозицію) тощо.

Також, викладачі під час проведення занять мають змогу провести рефлексію за допомогою виведення на екран «Дерева очікувань», яке учасники розробили на початку заняття, і поруч зобразити «Дерево плодів». Рефлексійна вправа передбачає перетягнення на це дерево стікерів із написаними очікуваннями, що виправдалися, коментування своїх результатів. Підбиття підсумків дозволяє дати відповіді на наступні запитання: «Чи справдилися очікування в учасників?», «Чи досягнуті цілі онлайн заняття?», «Що варто врахувати і на що звернути увагу під час планування наступного онлайн заняття?».

Висновки. Таким чином, аналіз використання технологій дистанційного навчання для здійснення рефлексії під час лекційних, семінарських та практичних занять для фармацевтів-інтернів в рамках курсу «Управління та економіка фармації» на кафедрі ОЕУФ ІПКСФ НФаУ показав широкий асортимент використаних педагогічних методів, прийомів та інструментів для проведення рефлексії, що значно впливає на ефективність засвоєння матеріалу фармацевтами -інтернами.

УДК 615.1:37.018.43:378.046-021.68

ДОСВІД ВИКОРИСТАННЯ ДИСТАНЦІЙНОЇ СКЛАДОВОЇ ОЧНО-ДИСТАНЦІЙНОГО НАВЧАННЯ В УМОВАХ ВОЄННОГО СТАНУ НА КАФЕДРІ КЛІНІЧНОЇ ФАРМАКОЛОГІЇ

Бондарєв Є. В., Бутко Я. О., Міщенко О. Я., Файзуллін О. В., Тимченко Ю. В., Подольский І. М.

Національний фармацевтичний університет, м. Харків, Україна
Інститут підвищення кваліфікації спеціалістів фармації
Кафедра клінічної фармакології
clinpharmacol_ipksph@nuph.edu.ua

Анотація. Мета цієї роботи полягає в аналізі та оцінюванні системи очно-дистанційного навчання у закладі післядипломної освіти ІПКСФ НФаУ, виявленні переваг та вирішення проблем, що впливають на якість післядипломної освіти. Результати дослідження показали, що в умовах воєнного стану дистанційне навчання дозволило якісно закінчити курси із спеціалізації «Клінічна фармація» працівникам з повною вищою фармацевтичною освітою та первинною спеціалізацією «Клінічна фармація» / «Фармація» та опанувати нові знання щодо сучасних досягнень фармацевтичних та медичних наук; знання щодо клінічних аспектів діяльності клінічного фармацевта, чинного законодавства про охорону здоров'я та нормативні документи, що регламентують організацію фармацевтичного забезпечення населення, закладів охорони здоров'я лікарськими засобами та іншими товарами аптечного асортименту; навичок застосовування на практиці вмінь та формування нових навичок щодо клінічної діяльності, зокрема адміністрування антимікробних препаратів з метою протидії антибіотикорезистентності, та можливостей впровадження й використання сучасних методів, технологій, технік, практик, форм у діяльність спеціалістів фармації. Платформи дистанційного навчання: Moodle, Zoom, Youtube та Google, що ефективно використовують для освіти та можуть бути використанні у якості корисного доповнення для очного навчання.

Ключові слова: *дистанційне навчання, післядипломна освіта, підвищення кваліфікації спеціалістів фармації та медицини, Moodle.*

UDC 615.1:37.018.43:378.046-021.68

EXPERIENCE OF USING THE DISTANCE EDUCATION IN THE CONDITIONS OF MARTIAL STATE AT THE DEPARTMENT OF CLINICAL PHARMACOLOGY

Bondarev E. V., Butko Y. O., Mishchenko O. Ya., Faizullin O. V., Tymchenko Yu. V., Podolsky I. M.

Abstract. The purpose of this work is to evaluate the distance learning system in the postgraduate education Institution of the Advanced Training of Pharmacy Specialists of the National Pharmaceutical Univercity, to identify the advantages and

solve the problems that affect the quality of postgraduate education. The results of the study showed that in the conditions of martial law, distance learning allowed employees with a complete higher pharmaceutical education and primary specialization “Clinical Pharmacy” / “Pharmacy” to complete courses in the cycle of specialization “Clinical Pharmacy” and acquire new knowledge of modern achievements in pharmaceutical and medical sciences; deepening knowledge of the clinical aspects of the clinical pharmacist's activities, including the administration of antimicrobial drugs to combat antibiotic resistance, current health care legislation and regulations governing the organization of pharmaceutical Distance learning platforms: Moodle, Zoom, Youtube and Google, which are effectively used for education and can be used as a useful supplement to face-to-face training.

Keywords: *distance learning, postgraduate education, advanced training of pharmacy and medicine specialists, Moodle.*

Вступ. Останні події, а саме пандемія COVID-19 та військовий стан в Україні, змусили систему освіти шукати нові методи викладання та навчання (дистанційна освіта). Таке навчання дозволяє об’єднати використання декількох форм очного навчання та платформ дистанційного навчання та закласти основи змішаного навчання [1-3, 7]. В Національному фармацевтичному університеті (НФаУ) в Інституті підвищення кваліфікації спеціалістів фармації та медицини (ІПКСФ) в освітньому процесі все більше йде акцент на очно-дистанційне навчання, віддалену працю педагогічних працівників та продуктивну роботу з інтернами та слухачами циклів підвищення кваліфікацій на етапі післядипломної освіти.

Матеріали та методи дослідження. На сучасному етапі розвитку суспільства характерні зміни, які викликані глобалізацією, створенням інформаційного суспільства, розвитком технологій. Все це приводить до пошуку нових форм здобуття освіти, однією з яких є дистанційна (очно-дистанційна) освіта. Дистанційна освіта призначена надати всім категоріям громадян рівні можливості в реалізації права людини на освіту незалежно від статі, раси, національності, соціального та майнового стану, характеру занять, світоглядних переконань, ставлення до релігії, віросповідання, стану здоров’я та місця проживання.

В статті проведено аналіз досвіду організації освітнього процесу в очно-дистанційному режимі на післядипломному етапі навчання в ІПКСФ НФаУ. Проведено огляд сучасних навчальних платформ навчання, технологій, комунікацій, методів та форм навчання.

Очно-дистанційна форма навчання оптимально поєднує в єдиному навчальному процесі принципи, методи та засоби очного та дистанційного навчання.

Метою очно-дистанційного навчання є надання освітніх послуг шляхом застосування сучасних інформаційно-комунікаційних технологій відповідно до державних стандартів освіти. Завданням очно-дистанційного навчання є забезпечення громадянам можливості реалізації конституційного права на навчання незалежно від статі, раси, національності, соціального і майнового стану, роду

та характеру занять, світоглядних переконань, належності до партій, ставлення до релігії, віросповідання, стану здоров'я, місця проживання відповідно до їх здібностей.

Очно-дистанційна форма післядипломної освіти дає можливість: підвищити рівень та якість самостійної роботи слухачів; розвивати інформаційну культуру слухачів на базі широкого застосування у навчанні сучасних інформаційних, комп'ютерних та телекомунікаційних технологій.

Дистанційна – це форма навчання, за якою викладачі, студенти, слухачі циклів фізично розділені, але взаємодіють за допомогою інформаційно-комунікативних технологій (конференцій, відео-чатів, веб-конференцій, вебінарів, он-лайн курсів, тощо). Розвиток такої технології в освіті дозволяє взаємодіяти між викладачем та слухачем за допомогою цифрових інструментів: смартфон, ноутбук, веб-камера, планшет, інтернет, таким чином дозволяють використовувати такі технології та створювати віртуальні класи, віртуальні лабораторії. Саме таке навчання дозволяє використовувати віртуальне середовище та використання навчальних платформ [2, 4].

Он-лайн навчання в закладах вищої фармацевтичної та медичної освіти не замінює очного навчання, воно його доповнює та є корисним. Так, курси підвищення кваліфікацій слухачам медико-фармацевтичних спеціальностей можна проводити синхронно за допомогою аудіо-, відео-, веб-конференцій, вебінарів або асинхронно за допомогою он-лайн форумів, дискусій, обміну повідомленнями, електронної пошти, прикріплення даних до платформи Moodle, тестування.

Перехід до очно-дистанційного навчання в закладах вищої освіти та на етапі післядипломного навчання стає перспективним напрямком. Система викладання матеріалів включає: створення програми навчання, цілі, ресурси, компетентності та заплановані результати навчання, структура освітньої компоненти, зміст програми, теми лекцій та семінарських та практичних занять, завдання для самостійної роботи, критерії оцінювання та ін.

В умовах воєнного стану в Україні в закладах вищої фармацевтичної та медичної освіти використовують платформу Moodle, використовують програми для спілкування за технологією Zoom, Youtube та Google.

Незважаючи на переваги навчальних платформ, дистанційне навчання для майбутніх спеціалістів медико-фармацевтичних спеціальностей явище тимчасове і застосовується під час катаклізмів, воєнного стану та ін. За допомогою дистанційного навчання студенти, слухачі курсів підвищення опрацьовують практичні та семінарські заняття, вебінари та певну кількість годин індивідуальної роботи. Тому, саме дистанційне навчання сприяє індивідуальне навчання, за якою студент (слухач циклу) розробляє для себе власний навчальний розклад занять залежно від мотивації, власних потреб та інтересів.

Індивідуальна робота в рамках самостійної підготовки передбачає формування відповідних навичок, використання підручників, додаткової літератури на курсі, методичних розробок, використання платформи Moodle при пошуку

необхідної інформації. Використання пошукових систем Google, Bing, Copilot та ін. передбачає не тільки перегляд а і розуміння отриманої інформації, збереження її для подальшого використання [5].

Застосування сучасних технологій та комунікацій – комп'ютерів, інтернет мереж, програмного ліцензійного забезпечення, платформи Moodle дає змогу широко використовувати дистанційне навчання в освіті [6, 8-10].

Мета цього дослідження полягає в оцінці методів дистанційного навчання шляхом вивчення платформи Moodle, визначити переваги, зручність використання платформи Moodle в процесі проведення дистанційного навчання на етапі післядипломної освіти.

Результати та їх обговорення. В ІПКСФ НФаУ заняття для спеціалістів фармації та медицини проводять в очно-дистанційному режимі з використанням технології Zoom та на платформі Moodle. Ця платформа дозволяє розробляти онлайн курси, які віртуально об'єднують велику кількість слухачів, інтернів, спеціалістів фармації та медицини. Кожен слухач може проходити курси вдома, на роботі чи в безпечному місці, зберігаючи при цьому взаємодію з іншими слухачами. Ця концепція дистанційного навчання виникла в американських університетах та країнах ЄС [8].

В рамках організації навчального процесу в закладах вищої освіти України використовують саме платформу Moodle, яка передбачає синхронний, асинхронний тип навчання.

Синхронна форма навчання - ця форма навчання передбачає електронне навчання, в якій слухач перебуває одночасно з іншими слухачами (учасниками) на заняттях. Викладання та навчання здійснюється на відстані з використанням інформаційних технологій та платформ.

Асинхронна форма навчання – он-лайн навчання, де слухач перебуває одночасно з іншими учасниками заняття, використовуючи підключення до інтернет мережі та пристроїв зв'язку (ноутбук з камерою, ПК з камерою, колонки та ін.).

Впровадження дистанційного навчання передбачає декілька прийомів: тільки он-лайн навчання; тільки очне навчання; або комбіноване навчання. Саме комбіноване навчання є найефективнішим.

Платформа (Moodle) – це платформа для дистанційного навчання та базується на педагогічних принципах, дозволяє через мережу налагоджувати взаємодію між студентами, викладачами, навчальними програмами та ресурсами. Управління платформою в університетах та в закладах вищої освіти вимагає адміністрування (управління обліковими записами, реєстрація, дозволи та безпеки, резервне копіювання, зовнішній вигляд, тощо.).

В ІПКСФ НФаУ навчальні матеріали усіх освітніх компонент доступні на платформі (Moodle) та закріплені за кожною з кафедр інституту. На платформі (Moodle) на кожному з курсів завантажено всю необхідну інформацію, матеріали для навчання, презентації лекцій, відео-лекції тощо.

Найбільшим недоліком використання платформи є нестабільність мережі інтернет, і у разі знеструмлення серверів робота платформи буде недоступною для всіх користувачів.

Moodle – це зручна система управління дистанційним навчанням, яке використовують у всьому світі для викладання он-лайн.

Дистанційне навчання є успішним у всьому світі, де викладачі визначають завдання, методи та форми надання матеріалів, терміни виконання, систему контролю знань, порядок звітності.

Робота на платформі Moodle дозволяє слухачам диференціювати завдання, обирати зручний час роботи на платформі, здійснювати самостійно пошук відповідей на практичних заняттях та семінарах. Навчаючись самостійно слухачі можуть навчатися за своїм графіком, вибирати час навчання, організує своє навчання на власний розсуд.

Однак кожна з платформ має свої як переваги так і недоліки. До недоліків можна віднести: нестабільність інтернет, віруси в мережі, певні вразливості. Слабким місцем у Zoom є обмеження тривалості заняття (необхідність отримання ліцензії), необхідно створювати окремі акаунти (електронна пошта) для входу до конференції (проведення занять), яка має бути відокремлена від інших акаунтів та доступом до соціальних мереж.

Впродовж 2023 року слухачі спеціалізації «Клінічна фармація» опанували знання та підвищили професійний рівень знань з основ клінічної медицини, клінічної фармакології основних груп лікарських засобів, питань клінічної фармації та фармацевтичної опіки, лабораторної діагностики, основ медичної стандартизації, організації охорони здоров'я, страхової медицини, фармакоекономіки та формулярної системи, інформаційних технологій, необхідних для здійснення професійної діяльності у закладах охорони здоров'я для забезпечення раціонального використання лікарських засобів. Впродовж циклу слухачі (понад 130 осіб) отримали за допомогою очно-дистанційного навчання на платформах Zoom та Moodle нові знання, розібрали клінічні аспекти діяльності клінічного фармацевта, поглибили навички застосовування на практиці вмінь та формування нових навичок щодо клінічної діяльності та можливостей впровадження та використання сучасних методів, технологій, технік, практик, форм у діяльність спеціалістів фармації.

Наприкінці курсів підвищення кваліфікації слухачі проходили анкетування щодо оцінювання курсів. Результати анкетування показали, що на курсі навчалися фармацевти, клінічні фармацевти, завідувачі аптек, доценти закладів вищої освіти (ЗВО), директори аптек, тощо. Стаж роботи слухачів від 1до 36 років. Завдяки очно-дистанційній освіті на курсах були зареєстровані слухачі з різних регіонів України таких як: Харків, Полтава, Суми, Запоріжжя, Київ, Житомир, Чернігів, Біла Церква, Кропивницький, Івано-Франківськ, Рівно, Дубно, Мукачево, Тернопіль, Луцьк та ін. місць.

Клінічні фармацевти після завершення курсів залишили свої відгуки та побажання. Так, більшість слухачів написали, що вони довго чекали саме на ці

курси, і що вони впродовж шести місяців навчання опанували знання щодо клінічної фармакології, отримали безцінний досвід та знання за спеціальністю, здобули навички практичного застосування отриманих знань. Більшість слухачів отримали для себе цінну інформацію, лекційний матеріал, можливість долучатися до дискусій та обміну думками, зручний формат навчання (очно-дистанційна форма навчання).

Висновки. Отримані результати дослідження дозволяють зробити висновок, що в ІПКСФ НФаУ система очно-дистанційної освіти забезпечує належний рівень знань слухачів в умовах воєнного стану в Україні. Встановлено, що впровадження системи очно-дистанційного навчання в ІПКСФ НФаУ в освітній процес післядипломної освіти є перспективним та дієвим для спеціалістів фармації та медицини, є можливість охоплювати спеціалістів з медичною та фармацевтичною освітою з різних регіонів нашої України.

Список використаних джерел інформації:

1. Інформаційні технології у фармації: підручник / О. А. Рижов , Ю. М. Пенкін, Н. А Іванькова. Львів: Магнолія, 2021. 260с.
2. Дистанційне навчання, переваги, недоліки, питання організації : Рекомендаційний список літератури / уклад. І. А. Каліниченко ; ЗДМУ, наукова бібліотека. – Запоріжжя, 2021. – 12 с.
3. Бачурін Г. В. Дистанційне навчання – як виклик часу під час пандемії COVID–19 / Г. В. Бачурін, С. С. Ломака, Ю. С. Коломоєць // Урологія. – 2020. – Т. 24, N 3. – С. 257–258.
4. Купрієвич В. Дистанційне навчання в системі підвищення кваліфікації як засіб професійного самовдосконалення керівників професійно- технічних навчальних закладів / В. Купрієвич // Післядипломна освіта в Україні. – 2017. – N 1. – С. 94–96.
5. Тимощук О.В. Особливості впливу дистанційного навчання на якість життя студентів сучасних закладів різних типів / О. В. Тимощук // Art of Medicine. – 2020. – N 4. – С. 124–128.
6. Швирка В. Потенціал дистанційного навчання у вищих навчальних закладах Сходу України (сучасні реалії) / В. Швирка // Вища школа. –2015. – N 7/8. – С. 66–72.
7. Дистанційна вища технічна освіта в умовах карантину / Г. Козлакова [та ін.] // Вища освіта України. – 2020. – N 2. – С. 67–74.
8. Авдєєв, О. В. Використання дистанційної системи MOODLE для оптимізації навчального процесу у вищій школі / О. В. Авдєєв // Мед. освіта. – 2015. – N 1. – С. 6–8.
9. Дистанційне навчання як інформаційно– комунікативна технологія додипломної підготовки лікарів з педіатрії в умовах сучасних викликів/ О. П. Волосовець [та ін.] // Мед. освіта. – 2020. – N 3. – С. 9–12.
10. Досвід впровадження онлайн–технологій у систему підготовки фахівців галузі знань 22 «Охорона здоров'я» / Ю. М. Колесник [та ін.] // Мед. освіта. – 2018. – N 2. – С. 69–73.

ПИТАННЯ БЕЗПЕКИ ЛІКІВ ТА БЕЗПЕЧНОЇ ФАРМАКОТЕРАПІЇ ПРИ ВИВЧЕННІ ПОБІЧНОЇ ДІЇ ЛІКІВ СТУДЕНТАМИ СПЕЦІАЛЬНОСТІ «ФАРМАЦІЯ, ПРОМИСЛОВА ФАРМАЦІЯ»

Кишкан І. Г.

Буковинський державний медичний університет, м. Чернівці, Україна

Кафедра фармакології

kishkaninna@bsmu.edu.ua

Вступ. На сьогодні проблема безпеки ліків та безпечної фармакотерапії стає все більш актуальною як у клінічній практиці, так і в освітній сфері на етапі підготовки майбутніх фармацевтів. Адже кожен лікарський засіб (ЛЗ) крім фармакотерапевтичної дії може викликати небажані побічні ефекти. Тому при медичному застосуванні ЛЗ одним із важливих питань залишається профілактика їх побічної дії (ПД).

Мета. Висвітлити роль курсу за вибором студента «Побічна дія ліків» у засвоєнні майбутніми фармацевтами клініко-фармакологічних аспектів ПД ліків, його важливість у формуванні умінь щодо оцінки співвідношення «користь/ризик» при застосуванні ЛЗ та розробці обґрунтованих заходів попередження ПД при проведенні фармакотерапії та фармацевтичної опіки.

Результати. Під час вивчення курсу за вибором «Побічна дія ліків» студенти ІІІ курсу спеціальності «Фармація, промислова фармація» засвоюють основні терміни та загальні положення стосовно ПД ліків, фактори, що сприяють її виникненню, вивчають форми, методи отримання та аналізу інформації про ПД ЛЗ, закріплюють клініко-фармакологічну класифікацію ПД ліків та заходи запобігання ускладнень фармакотерапії. Майбутні фармацевти на окремих заняттях детально обговорюють питання стосовно особливостей ПД ЛЗ у дітей та людей літнього віку, ролі генетичних факторів, впливу шкідливих звичок на фармакодинаміку та фармакокінетику ліків, проблему лікарської алергії, тахіфілаксії, поліпрагмазії, несумісності ЛЗ, можливої ембріотоксичної, тератогенної дії, лікарської залежності та «синдрому відміни». Курс «Побічна дія ліків» допомагає студентам сформувати уміння визначати причетність до ПД чинників, що впливають на процеси всмоктування, розподілу, депонування, метаболізму та виведення ЛЗ, встановлювати фактори, обумовлені індивідуальними особливостями організму людини та фізико-хімічними властивостями ЛЗ. Також у процесі вивчення цього курсу студенти закріплюють уміння визначати переваги, недоліки, ПД ліків різних фармакологічних груп з урахуванням їхніх біофармацевтичних, фармакокінетичних та фармакодинамічних особливостей, уміння рекомендувати препарати «прикриття» для зменшення та попередження ПД. Опановують навички стосовно раціонального й безпечного застосування ЛЗ із наданням консультативної допомоги пацієнтам.

Висновок. Вивчення курсу «Побічна дія ліків» сприяє формуванню у майбутніх фармацевтів фахових компетентностей, необхідних для проведення безпечної фармакотерапії та грамотної фармацевтичної опіки.

УДК 615.03

ПОРІВНЯЛЬНИЙ АНАЛІЗ УКРАЇНСЬКОГО ТА МІЖНАРОДНОГО ДОСВІДУ ПІДГОТОВКИ КЛІНІЧНОГО ФАРМАЦЕВТА

Штриголь С. Ю.[1], Жаботинська Н. В.[1], Кіреєв І. В.[2]

[1]Національний фармацевтичний університет

[1]Кафедра фармакології та клінічної фармації

[2]Інститут підвищення кваліфікації спеціалістів фармації

Кафедра клінічної фармакології[2]

м. Харків, Україна

bronkevih@gmail.com

Анотація. Спеціальність «Клінічна фармація» набуває великої популярності на ринку праці України, що вимагає оновити підходи до підготовки клінічних фармацевтів з урахуванням сучасних тенденцій щодо практичної реалізації їх діяльності.

Мета. Провести порівняльний аналіз сучасного міжнародного досвіду в підготовці клінічних фармацевтів і порівняти його з аналогічною практикою в Україні.

Матеріали та методи. При підготовці статті були вивчені інформаційні матеріали та освітні програми підготовки клінічних фармацевтів закордоном. Використовувався аналітичний метод.

Результати та їх обговорення. В усьому світі клінічні фармацевти є членами мультидисциплінаної команди та несуть відповідальність за медикаментозну терапію та результати лікування пацієнтів разом із лікарями. Підготовка клінічного фармацевта починається з бакалаврського рівня, далі магістратура, яку можуть закінчити особи, що мають ступінь бакалавра наук в іншій галузі медичної освіти. Далі клінічні фармацевти можуть отримати додаткову підготовку в програмах післядипломної ординатури першого (PGY-1) та другого (PGY-2) року навчання. Ліцензію клінічні фармацевти отримують на 6 років.

Висновки. Високі міжнародні вимоги до отримання освіти клінічним фармацевтом спонукають поглиблювати освітні програми для клінічного фармацевта та розширювати в Україні можливості вузькоспеціалізованого безперервного професійного розвитку клінічних фармацевтів.

Ключові слова: *клінічний фармацевт, освіта, міжнародний досвід.*

UDC 615.03

COMPARATIVE ANALYSIS OF UKRAINIAN AND INTERNATIONAL TRAINING EXPERIENCE OF CLINICAL PHARMACISTS

Shtrygol' S. Yu., Zhabotynska N. V., Kireyev I. V.

Abstract. The specialty "Clinical Pharmacy" is gaining great popularity in the Ukraine, which requires updating approaches to the training of clinical pharmacists, taking into account modern trends in the practical implementation of their activities.

Aim. Conduct a comparative analysis of modern international experience in the training of clinical pharmacists and compare it with similar practice in Ukraine.

Materials and methods. When preparing the article, informational materials and educational programs for training clinical pharmacists abroad were studied. An analytical method was used. Results and discussion. Around the world, clinical pharmacists are members of a multidisciplinary team who are responsible for drug therapy and patient outcomes alongside physicians. The training of a clinical pharmacist begins with a bachelor's degree, followed by a master's degree, which can also be completed by persons who have a bachelor's degree in another field of medical education. Further, clinical pharmacists can receive additional training in first (PGY-1) and second (PGY-2) year postgraduate residency programs. Clinical pharmacists receive a license for 6 years.

Conclusions. High international requirements for the education of a clinical pharmacist encourage the deepening of educational programs for clinical pharmacists and the expansion of highly specialized continuous professional development of clinical pharmacists in Ukraine.

Key words: *clinical pharmacist, education, international experience.*

Вступ. Відомо, що в Україні певний час була недостатньо реалізованою роль клінічного фармацевта в системі охорони здоров'я. Тривалий час виникали труднощі при працевлаштуванні клінічних фармацевтів. Але Наказ МОЗ від 03.08.2021 року № 1614 зі змінами від 21.02.2023 року не лише розв'язав цю проблему, а й став своєрідним визнанням внеску таких спеціалістів у медичне забезпечення пацієнтів та гарантією працевлаштування клінічних фармацевтів у закладах охорони здоров'я [1]. Нині спеціальність «Клінічна фармація» набуває великої популярності на ринку праці України, що вимагає оновити підходи до підготовки клінічних фармацевтів з урахуванням сучасних тенденцій щодо практичної реалізації їх діяльності.

Мета. Провести порівняльний аналіз сучасного міжнародного досвіду підготовці клінічних фармацевтів і порівняти його з аналогічною практикою в Україні.

Матеріали та методи. Інформаційні матеріали щодо можливостей навчання на спеціальності «Клінічна фармація» в різних країнах, освітні програми підготовки клінічних фармацевтів. Використовувався аналітичний метод.

Результати та їх обговорення. У багатьох країнах працюють клінічні фармацевти, що поліпшують надання медичної допомоги пацієнтами будь-якого віку в усіх сферах невідкладної та амбулаторної допомоги. Але можна вважати найкращою модель їх підготовки та подальшої практичної діяльності в Сполучених Штатах Америки. Американський коледж клінічної фармації є закладом, що зосереджений на розвитку, просуванні та позиціонуванні клінічних фармацевтів серед інших постачальників, громадськості та професійних товариств [2]. Тому будемо спиратися саме на їх визначення клінічного фармацевта

як спеціаліста, що пройшов спеціальну підготовку та надає безпосередню медичну допомогу пацієнтам у комплексному застосуванні лікарських засобів.

Де ж можуть працювати клінічні фармацевти? Згідно із світовою практикою, клінічні фармацевти практикують у всіх закладах охорони здоров'я та використовують глибокі знання про лікарські засоби та захворювання, щоб керувати медикаментозною терапією як частина мультидисциплінарної команди. Клінічні фармацевти несуть відповідальність за фармакотерапію та результати лікування пацієнтів разом із лікарями. Вони є основним джерелом науково обґрунтованої інформації щодо безпечного, відповідного клінічному випадку та економічно ефективного використання ліків.

В Україні посада клінічного фармацевта згідно з вищезгаданим Наказом МОЗ № 1614 входить до штатного розпису відділів інфекційного контролю, що мають бути створені в усіх без винятку закладах охорони здоров'я та соціального захисту незалежно від форми власності та відомчого підпорядкування (за винятком аптек та аптечних підрозділів). Насамперед такі фахівці займаються впровадженням програми адміністрування антимікробних препаратів (антибіотиків, противірусних, протипаразитарних та протигрибкових лікарських засобів) на рівні закладу охорони здоров'я.

Крім того, клінічні фармацевти в США мають приватну практику та забезпечують медичну допомогу пацієнтам, яких скерували лікарі різного профілю. На жаль, це досвід в Україні поки що не реалізований.

Для реалізації свої повноважень клінічні фармацевти в США практикують уніфікований алгоритм діяльності Pharmacist Process of Care – процес фармацевтичної допомоги. З урахуванням цього алгоритму діяльності клінічного фармацевта сформульовано його узагальнені посадові обов'язки [3, 4]:

– Допомога іншим медичним працівникам у створенні планів лікування пацієнтів.

– Створення аптечних програм, які допомагають навчати інших фармацевтів і пацієнтів.

– Навчання пацієнтів щодо застосування призначених їм лікарських препаратів, зокрема дозування, будь-яку потенційну взаємодію ліків та побічні ефекти.

– Робота з пацієнтами з метою виявлення негативних побічних ефектів.

– Клінічний фармацевт повинен переконатися, що пацієнти дотримуються інструкцій щодо застосування рецептурних лікарських засобів.

– Контроль використання пацієнтами рецептів, щоб переконатися, що вони не зловживають ліками.

– Контроль надходження ліків та замовлення додаткових ліків.

– Рекомендація найбільш оптимальних шляхів введення лікарських засобів на основі потреб пацієнта.

– Надання консультацій пацієнтам щодо їх ліків та медичних проблем.

– Відстеження листів призначення лікарських засобів, щоб переконатися, що медичні сестри дають пацієнтам правильні ліки та точно записують інформацію.

– Клінічний фармацевт повинен бути в курсі всіх аспектів медицини, щоб стежити за новими розробками лікарських засобів, відкликанням ліків або іншою важливою інформацією про ліки, що відпускаються за рецептом.

Як бачимо, перелік функціональних обов'язків клінічного фармацевта вимагає ґрунтовної освіти. Підготовка клінічного фармацевта відрізняється в різних країнах світу. У США навчання починається з бакалаврського рівня, далі магістратура, яку можуть закінчити також особи, що мають ступінь бакалавра наук в іншій галузі медичної освіти. Для безпосередньої участі в медичній допомозі пацієнтами клінічні фармацевти можуть отримати додаткову підготовку в програмах післядипломної ординатури першого року (PGY-1) з невідкладної медичної допомоги або з медичної допомоги в амбулаторних умовах. Це широкомасштабний акредитований рівень підготовки у сфері клінічної допомоги, інформації про ліки, адміністрування, методів навчання та проектів/досліджень протягом 12 місяців. Ті, хто зацікавлений у спеціалізації, можуть пройти другий рік післядипломної ординатури (PGY-2) у таких різноманітних сферах, як будь-яка медична спеціальність (амбулаторна допомога, реанімаційна допомога, інфекційні захворювання, внутрішня медицина, онкологія та багато інших). Клінічний фармацевт має право на отримання ліцензії після 6 років навчання та отримання ступеня доктора фармації. Після навчання багато клінічних фармацевтів прагнуть отримати сертифікат Ради зі спеціальності фармація – Board of Pharmacy Specialties (BPS) з однієї або декількох з восьми фармацевтичних спеціальностей: амбулаторна допомога, реанімація, базова медична допомога, підтримка харчування, онкологія, педіатрія, фармакотерапія та психіатрична фармація. В майбутньому спеціальні сертифікати можуть включати інфекційні захворювання та кардіологію. Сертифіковані фармацевти повинні пройти суворий іспит, а потім підтримувати акредитацію шляхом продовження навчання або додаткового тестування кожні 7 років. Сертифікація Ради є типовою вимогою для отримання привілеїв для приватної або спільної практики в мультидисциплінарній команді.

В Україні нині освіту клінічного фармацевта другого (магістерського) рівня у галузі знань 22 «Охорона здоров'я» за спеціальністю 226 «Фармація, промислова фармація», спеціалізації 226.01 «Фармація» за освітньою програмою «Клінічна фармація» можна здобути тільки в Національному фармацевтичному університеті [5]. Далі на базі Інституту підвищення кваліфікації спеціалістів фармації ці спеціалісти мають пройти інтернатуру за спеціальністю «Фармація» з наступною спеціалізацією «Клінічна фармація» та отримати сертифікат зі спеціальності «Клінічна фармація». В подальшому клінічні фармацевти можуть навчатися на курсах підвищення кваліфікації в рамках безперервного розвитку, але в Україні вони не мають такого вузькоспеціалізованого характеру, як у

США. Ліцензування діяльності клінічного фармацевта в Україні не передбачено.

Висновки. Таким чином, існують високі міжнародні вимоги до отримання освіти клінічного фармацевта з подальшою вузькою спеціалізацією та підвищенням кваліфікації для отримання ліцензії кожні 7 років. Це зумовлено тим, що професійна діяльність клінічного фармацевта підкреслює його роль як медичного працівника, що виходить за рамки традиційного відпуску ліків та активно сприяє ефективній медичній допомозі пацієнтові та забезпеченню громадського здоров'я. В Україні потрібно розширювати можливості вузькоспеціалізованого безперервного професійного розвитку клінічних фармацевтів.

Перелік використаних джерел інформації:

1. Про організацію профілактики інфекцій та інфекційного контролю в закладах охорони здоров'я та установах/ закладах надання соціальних послуг/ соціального захисту населення. Наказ МОЗ № 1614 від 03.08.2021 року URL: https://zakononline.com.ua/documents/show/502234___760886 (дата звернення: 16.10.2024).
2. American College of Clinical Pharmacy URL: https://www.accp.com/ (дата звернення: 16.10.2024).
3. Спеціальність фармакологія в університетах США та Великобританії/ URL: https://unimind.com.ua/ua/holovna/blog1/spetsialnist-farmakologiya-u-universitetakh-ssha-ta-velikobritanii (дата звернення: 16.10.2024).
4. Massachusetts College of Pharmacy and Health Sciences URL: https://edusteps.com.ua/programs/universities/university-usa/1993-massachusetts-college-of-pharmacy-and-health-sciences.html (дата звернення: 16.10.2024).
5. Освітня програма клінічна фармація. НФаУ. URL: https://nuph.edu.ua/klinichna-farmaciya-osvitnya-programa-2/ (дата звернення: 16.10.2024).

SCIENTIFIC DISCOURSE OF ELEVATING THE ROLE OF PHARMACISTS: A GLOBAL AND GEORGIAN PERSPECTIVES ON CLINICAL RESPONSIBILITIES, PROFESSIONAL ENHANCEMENT, JOB CONTENTMENT AND HEALTHCARE IMPACT

Nodar Sulashvili[1], Margarita Beglaryan[2], Nana Gorgaslidze[3], Luiza Gabunia[4], Natia Kvizhinadze [5], Nato Alavidze [6], Nino Abuladze[7], Ketevani Gabunia[8], Shorena Khetsuriani[9], Natia Gamkrelidze[10], Lia Gumbaridze[11], Londa Rukhadze[12], Giorgi Pkhakadze[13], Diego Rada Fernandez de Jauregui[14], Marika Sulashvili[15], Marina Giorgobiani[16], Tamar Okropiridze[17], Igor Seniuk[18], Lela Grigolia[19], Kakhaber Robakidze[20], Irine Zarnadze[21], Shalva (Davit) Zarnadze[22]

[1]Tbilisi State Medical University-Scientific Research-Skills Center; Georgian National University SEU- Faculty of Medicine; Sulkhan-Saba Orbeliani University - Faculty of Medicine; Alte University International - School of Medicine; Shota Meskhia Zugdidi State University - Pharmacy Program; David Aghmashenebeli University of Georgia - School of Medicine; University of Georgia - School of Health Sciences, Biochemistry and Pharmacology Direction; East European University - Faculty of Medicine; Tbilisi Humanitarian Teaching University - Faculty of Dentistry and Pharmacy; Tbilisi, Georgia; Yerevan State Medical University after Mkhitar Heratsi - Department of Pharmaceutical Management, Yerevan, Armenia.

[2]Yerevan State Medical University After Mkhitar Heratsi - Department of Pharmaceutical Management, Yerevan, Armenia.

[3]Tbilisi State Medical University, Department of Social and Clinical Pharmacy, Tbilisi, Georgia.

[4]Tbilisi State Medical University - Department of Medical Pharmacology, Tbilisi State Medical University- Scientific Research-Skills Center, Tbilisi, Georgia.

[5]Tbilisi State Medical University, Department of Social and Clinical Pharmacy, Tbilisi, Georgia.

[6]Akaki Tsereteli State University, Faculty of Medicine, Department of Pharmacy, Kutaisi, Georgia. East European University, Tbilisi, Georgia.

[7]Akaki Tsereteli State University, Faculty of Medicine, Department of Pharmacy, Kutaisi, Georgia.

[8]Akaki Tsereteli State University, Faculty of Medicine, Department of Pharmacy, Kutaisi, Georgia.

[9]Tbilisi State Medical University-Scientific Research Skills Center; Tbilisi State Medical University Microbiology Department, Tbilisi, Georgia.

[10]Tbilisi State Medical University - Scientific Research Skills Center, Tbilisi State Medical University - Pathophysiology Department, Tbilisi, Georgia.

[11]Tbilisi State Medical University Scientific Research-Skills, Tbilisi State Medical University-Department of Public Health Management, Policy and Economics, Tbilisi, Georgia.

[12]Tbilisi State Medical University Scientific Research Skills Centre, Tbilisi, Georgia. The University of Porto, Faculty of Healthcare, Porto, Portugal.

[13]David Tvildiani Medical University -School of Public Health, Tbilisi, Georgia; United Nations Secretary General's Independent Accountability Panel, Geneva, Switzerland; Accreditation San Frontières, Paris, France, Lviv Ukraine.

[14]University of the Basque Country, Pharmacy Faculty, Preventive Medicine and Public Health Department, Spain.

[15]Tbilisi State Medical University, Lecturer of Department of Molecular and Medical Genetics, Tbilisi, Georgia.

[16]*Tbilisi State Medical University, Department of Hygiene and Medical Ecology, Tbilisi, Georgia.*
[17]*Alte University, Division of Dentistry of International School of Medicine; University Geomedi, Department of Dentistry; The University of Georgia, Dentistry Department - The School of Health Sciences, Tbilisi, Georgia.*
[18]*National University of Pharmacy of Ukraine, Faculty of Pharmacy, Biological Chemistry Department, Kharkiv, Ukraine.*
[19]*Caucasus International University; Tbilisi, Georgia.*
[20]*Caucasus International University, Tbilisi, Georgia.*
[21]*Tbilisi State Medical University, Department of Public Health, Health Care Management, Policy and Economy, Tbilisi, Georgia.*
[22]*Tbilisi State Medical University, Department of Nutrition, Aging Medicine, Environmental and Occupational Health, Tbilisi, Georgia.*
n.sulashvili@ug.edu.ge

Abstract. The primary aim of this study was to examine the role of pharmacists and their perspectives on clinical responsibilities in Georgia and globally, while analyzing the factors influencing job satisfaction. Using a quantitative approach, the study involved comprehensive surveys and in-depth interviews across seven respondent groups, including chief pharmacists, pharmacy students, healthcare specialists, and patients. Data were gathered through approved questionnaires and analyzed using systematic sociological, statistical, and graphical methods, with SPSS version 11.0 for descriptive statistics and regression analyses. A Chi-square test was used to determine the statistical significance of associations ($p<0.05$). Findings highlight the need for pharmacists to be recognized as regulated healthcare professionals, with certification, licensing, and accreditation mandated by government bodies. Notably, around 25% of respondents viewed pharmacists as pivotal in treatment roles, akin to physicians, with well-trained pharmacists able to reduce prescription errors, thus enhancing patient safety and healthcare quality. The study underscored key areas of pharmaceutical practice, such as knowledge of new drugs, communication skills, pharmacotherapy, pharmacodynamics, pharmacokinetics, and pharmaceutical care. It emphasized the necessity for higher education institutions to prioritize subjects like pharmacotherapy, clinical pharmacy, and drug toxicity. Furthermore, a significant portion of public health specialists acknowledged the importance of pharmacists in dispensing medications, advising on drug efficacy, and assisting in drug selection, reinforcing the need for their greater authority in the healthcare system. To effectively educate patients on proper drug usage, pharmacists require extensive knowledge in medicine, pharmacology, clinical pharmacy, and related disciplines. Properly educated pharmacists play a critical role in enhancing healthcare services, pharmaceutical care, and ensuring patient safety. Additionally, the establishment and expansion of residency programs for clinical pharmacists are essential to cultivate a highly qualified workforce capable of providing advanced pharmaceutical care in clinical settings in Georgia. These measures are vital for elevating healthcare quality and ensuring patient safety.

Key words: *Pharmacists', role, perspectives, clinical, Georgia, worldwide, factors, pharmacist, job gratification.*

Introduction. The term "clinical pharmacy" was created to describe the role of pharmacists who primarily engage with other healthcare professionals. Their responsibilities include conducting meetings, interviews, and patient evaluations, providing specific pharmacotherapeutic advice, monitoring patients' responses to medications, and ensuring accurate information about pharmaceuticals. Clinical pharmacists typically work in clinics, hospitals, health insurance companies, and acute care settings, emphasizing patient-centered care over a production-focused approach. To effectively practice rational pharmacotherapy, clinical pharmacists must possess a comprehensive understanding of pharmaceutical sciences, medicine, pharmacology, pharmacotherapy, clinical pharmacology, and pharmaceutical care. This knowledge enables them to deliver care that maximizes therapeutic outcomes while minimizing costs, ultimately ensuring the health and safety of patients [1-4].

Responsible administering of drugs involves that healthcare network mediator capabilities and activities are balanced to assure that patients get the right drug, on the proper time, using properly and patient have profited from them. Delivering the right drugs into patients' demands commitment of all representatives, inclusive Government and a desire on how to consolidate private and public interests and mobilize sources. That is significant for the public to be guaranteed that expenses on pharmaceuticals productions are an equivalent cost of cash. On the viewpoints of the pharmacists' comprehensive academically field and their traditionary function in composing, qualifying, delivering and ensuring drugs. A pharmacist is informing customers, consumers and patients on the drug using; they are greatly positioned to suppose professional liability for the monitoring of pharmacotherapy. They are members of the healthcare team immediately engaged in patients' health care services. Their responsibility is to assistance patients in using their drugs, which is impossible to do alone. Thus, in terms pharmacists' profession have been progressed. New type pharmacists have done the work in more efficient way. Pharmacists holding the higher, university-level education. They understand the biochemical mechanisms of metabolism, mechanisms actions of drugs, medicines pharmacotherapeutic characteristic, side effects of drugs, potential interactions of drug and the argumentations monitoring. It is conjugated of specialized knowledge of biochemistry, anatomy, therapy, physiology, pathology, pharmacology and other pharmacy subjects. The pharmacists explain this particularized knowing when communicating with physicians, patients and another health care providers [5-7].

The modern system of pharmaceutical care is to improve and enhance the life quality of patients, which is promoted by highly skilled professionals in pharmacies, whose competence has been growing along the process of professional development. Pharmaceutical specialists should not only be to be capable to use their knowledge and skills gained at the educational institutions, but also should be ready and motivated for the professional self-development, because without qualified pharmaceutical care there is no qualified health care system. Since the scope of drug treatment, pharmacy is one of the most socially significant areas of the state regulation, the se-

quence of carrying out the reform measures, accumulation of experience, and also smooth introduction and application of new methods become crucial nowadays. Development of an organizational and functional model of licensing of pharmaceutical activities, pharmacists work as one of the mechanisms can improve the efficiency of public pharmaceutical administration, which has the great relevance, scientific and practical value [8-10].

The literature analysis showed that a study of individual aspects of the pharmaceutical specialists' professional development is directed to elaborating of the necessary requirements to ensure effective pharmaceutical care, postgraduate education, finding strategies for the better management of pharmaceutical personnel, pharmacists' job satisfaction and issues of their psycho-social adaptation to the emerging market conditions. However, weighable studies aimed at understanding the perspectives of the pharmaceutical specialist in the career aspect and the ways to provide high-quality pharmaceutical care have not been carried out yet [11-12].

The aim of the research was to provide a complex study, analysis and evaluation the pharmacists' role and perspectives from the view of clinical issue aspects in Georgia and worldwide and the factors effect to pharmacist job gratification.

Material and methods: Methodology of the study was sociological research: The study was quantitative investigation by using survey (Questionnaire). Research objectives are materials of sociological research: the study was quantitative investigation by using survey (Questionnaire). The in-depth interview method of the respondents was used in the study. The 7 types of approved questionnaires were used (Respondents were randomly selected): Questionnaire for chief pharmacists: 410 chief pharmacists participated in the study. Questionnaire for patients: 1506 patients (customers of drug-stores) participated in the study. Questionnaire for the employed pharmacy faculty-student: 222 employed pharmacy faculty students participated in the study. Questionnaire for healthcare specialists: 307 public health specialists participated in the study. Questionnaire for pharmacist specialist, 810 pharmacist specialists participated in the study. Totally 3888 respondents were interviewed in Georgia. We used methods of systematic, sociological (surveying, questioning), comparative, segmentation, mathematical-statistical, graphical analysis. The data was processed and analyzed with the SPSS program. Results and discussion: The survey was conducted through the questionnaires.1506 patients were interviewed in Georgia. Questions and answers are given in the tables. On each question are attached diagrams or table. Questionnaire and diagrams are numbered. The data was processed and analyzed with the SPSS program. We conducted descriptive statistics and regression analyses to detect an association between variables. Statistical analysis was done in SPSS version 11.0. A Chi-square test was applied to estimate the statistical significance and differences. Study of the data was processed and analyzed with the SPSS program. We conducted descriptive statistics and regression analyses to detect an association between variables. Statistical analysis was done in SPSS version 11.0. A Chi-square test was applied to estimate the statistical significance and differences. We defined $p<0.05$ as significant for all analyses. The study's ethical items. In order to provide the study's ethical character each participant of it was informed about the study's goal and

suggested of willingness of the work to be done. So, the respondents' written or oral compliance was got on that issue. All the studies were carried out by the selected organizations administrations' previous compliance. Were used Informed consent form for each respondent to participate in an anonymous survey. During the whole period of research, the participants incognita was also provided. For the international rules and criteria' conformity this human subject comprising given study was discussed and confirmed on the Bioethics Committee sessions of the YSMU. In order to meet the objectives, set in the research we also used the results obtained through analysis of available official information, studies and opinions about pharmacists, as well as the methods of quantitative studies. The research implementation required the following sub studies: the pharmacists' role and perspectives from the view of clinical issue aspects in Georgia and worldwide and the factors effect to pharmacist job gratification.

Results and discussion. On the basis of performed study results the following have been founded: The developed countries and many from the developing countries in the field of pharmacy have also worked out as family medicine. A pharmacist, as a family doctor, should not have just the higher, post-graduate and consistent education in pharmacy, but also needs the pharmacist license and periodic accreditation by the board of pharmacy (BOP). In the western countries' pharmacies, just the specialists with the higher pharmaceutical education who have graduated from the state-recognized and accredited colleges and universities are allowed to work. A pharmacy opening permit is issued only to a person of higher pharmaceutical education with the diploma [13-14]. In this concern a study of the distinctive specificities of pharmacists' higher educational perspectives from the viewpoints of clinical outlooks in Georgia, the pharmacist specialist's professional features, the factors influence on distinctive specificities of pharmacists' higher educational perspectives from the viewpoints of clinical outlooks in Georgia, the pharmacists' profession (occupational) choice and career satisfaction, pharmacists' work satisfaction by the duration of job and a balance between the pharmacists' workload and the personal life, professional peculiarities of student pharmacists and the employed pharmacist students; professional peculiarities of the pharmacists with the perspectives of chief (head), peculiarities of pharmacist' professional features according to the health-care specialists and patients, and professional peculiarities of young pharmacist specialists became of high actuality and the essence of our study. Therefore, should be done analyses of the current situation of distinctive specificities of pharmacists' higher educational perspectives from the viewpoints of clinical outlooks in Georgia and of the official statistical data, a study of the awareness level regarding the distinctive specificities of pharmacists' higher educational perspectives from the viewpoints of clinical outlooks in Georgia [15-16].

Based on the current situation analysis and identifying gaps in the pharmacy system, as well as taking into consideration distinctive specificities of pharmacists from the viewpoints of clinical outlooks in Georgia, compare international regulations in Europe and elaboration of the recommendations aimed at solving the existing problems have been encountered. A comprehension of the existing problems solving urgency moved us to perform the given research with the selected appropriate design,

which defined such objectives, as assessment and analyzing of the current situation of the distinctive specificities of pharmacists' higher educational perspectives from the viewpoints of clinical outlooks in Georgia and detection of the gaps and outlines of the Georgian clinical pharmacist system's perspectives. Development and justifying the practical recommendations for improving the professional development of clinical pharmacists and the quality of pharmaceutical care in whole [17-18].

Clinical pharmacists often apply their knowledge of drugs to a patient-specific treatment plan and evaluate dosage suitability, side effects, efficacy, and drug interactions. If necessary, the clinical pharmacist can discuss any issue and advise the physician, who is primarily responsible for prescribing drugs to patients, to ensure optimal use of the drugs. To practice, clinical pharmacists must graduate in a recognized area of qualification. The specific requirements for these degrees may differ depending on the country of operation. Subjects that are commonly found in the university's clinical pharmacist program include biology, chemistry, pathology, pharmacology, and socio-behavioral sciences. Most clinical pharmacists in the United States hold a Ph.D. in Pharmacy (Phar.D.) in addition to several years of postgraduate education such as a pharmaceutical residency. They can be certified as a clinical pharmacist through the Pharmaceutical Specialties Council, which is independent of the American Pharmacists Association. Education and certification requirements in other countries may differ depending on the guidelines set by the regulatory authorities. Clinical pharmacists are responsible for providing safe, effective, and timely drug therapy. Through various tasks in the department, they provide support for centralized and decentralized drug use systems, as well as optimal drug therapy for patients with a wide range of medical conditions. Clinical specialist pharmacists are competent in delivering direct patient-centered medical care and integrated operational pharmacy services in a decentralized practice with the participation of doctors, nurses and other hospital staff. These physicians are aligned with targeted multidisciplinary programs and specialized services to ensure drug therapy management within specialized patient care services and to ensure that pharmaceutical care programs are properly integrated across the facility. In these clinical roles, clinical pharmacists are involved in all necessary aspects of the drug use system, while providing comprehensive and personalized pharmaceutical care to patients in their assigned areas [19-20].

There are both ethical and practical imperatives to addressing health inequalities associated with chronic disease management for people with social difficulties, and existing programs often do not adequately meet the needs of these people. This leads to low participation rates, suboptimal chronic disease management and higher utilization of health services. Unlike acute conditions, chronic conditions require ongoing care and treatment outside of health care settings, in the community or in primary health care settings in terms of medication use, lifestyle management and behavior change in health. Typically, this is a multi-pronged intervention that includes a review of drug therapy, patient education for treatment, monitoring of medication, immunization, self-care, and support, disease, and / or prescribing authority. Patients who take a lot of medications due to chronic disease are at high risk for drug duplica-

tion, interactions, or ADRS, which can lead to longer hospital stays and higher costs. To improve the safety and efficacy of therapeutic agents, these patients must meet special needs for appropriate drug use. Research has shown that integrating pharmacists into outpatient clinics can improve chronic disease management and optimal medication use. Additionally, involving a pharmacist in patient care can help reduce inappropriate medication use, especially in the elderly. The study shows that the proportion of patients receiving the wrong drug drops considerably after review and optimization of the drug by a team including a pharmacist. Compared to conventional treatment, pharmacist-directed care was associated with a similar frequency or frequency of office visits, emergency room or emergency department visits, as well as hospitalizations and adherence, increases in the amount or dose of drugs received and improvements in study glycemic choices, blood pressure, and lipid target achievement. Another recent study shows that a telemedicine-based chronic disease management program involving clinical pharmacists resulted in statistically significant improvements in diabetes and hypertension outcomes as well as clinically significant improvements in lipid control and withdrawal. smoking [21-22].

The aging of the population has increased the burden of chronic disease around the world. There are both ethical and reasonable goals for addressing health inequalities identified in chronic disease management for people of multiple social origins, and existing programs routinely fail to meet the needs of these people. This translates into poor program support, poor management of chronic disease, and more frequent seeking of health care. Unlike acute conditions, chronic conditions require ongoing care and treatment outside of health care settings, in the community or in primary health care settings in terms of medication use, lifestyle management and behavior change in health. Typically, this is a multi-pronged intervention that includes a review of drug therapy, patient education for treatment, monitoring of medication, immunization, self-care, and support. disease, and / or prescribing authority. Patients who take multiple medications due to chronic disease are at high risk of drug duplication, interactions, or side effects, which can lead to longer hospital stays and higher costs. To improve the safety and effectiveness of the treatment, these patients must have specific needs for the correct use of the drugs encountered. Research has shown that integrating pharmacists into outpatient clinics can improve chronic disease management and optimal medication use. Additionally, involving a pharmacist in patient care can reduce the use of unwanted medications, especially in the past. A study in Canada found that the number of patients taking the wrong drug has decreased, mostly after screening tests and improvement by a group including a pharmacist. Unlike regular nursing, pharmacist-directed medical care was associated with a comparable frequency or pace of office work, major medical care or emergency room visits, and hospitalization and adherence, increased the rate of quantity or quantity of drugs received and improved study choices. indicators, blood circulation and blood circulation. achieve a lipid goal. Another recent study shows that a telemedicine-based chronic disease management program involving clinical pharmacists resulted in sta-

tistically significant improvements in diabetes and hypertension outcomes, as well as clinically significant improvements in lipid control. and smoking cessation [23-26].

The vast majority of the respondents considered that the professional activity of pharmacist is of great importance for the society (tabl.1).

Table 1

Opinion of the respondents (public health specialists) about importance of the pharmacist professional activity

Q-6. Assurance of the respondents in importance of the pharmacist professional activity for the society	Frequency	Percent (%)	Cumulative percent (%)
1. Yes	291	94.8	94.8
2. No	9	2.9	97.7
3. To a small extent	7	2.3	100.0
Total	307	100.0	

More than a half of respondents considered that the sense of the work of pharmacist is optimization of quality of life for people, related to health by providing the pharmaceutical care. Less than a half of respondents considered that work of pharmacist is timely, proper and qualitative supply with pharmaceutical products to the population and hospitals/clinics (table. 2).

Table 2

The respondents' opinion of the meaning of pharmacist work

Q-7. The sense of the pharmacist work	Frequency	Percent (%)	Cumulative percent (%)
1. Timely, proper and qualitative provision of the population and hospitals/clinics with pharmaceutical products	120	39.1	39.1
2. Optimization of quality of life for people related to health by providing the pharmaceutical care	187	60.9	100.0
Total	307	100.0	

The respondents' majority considered that the pharmacists' functions in a pharmacy consisted in realization of drugs and instruments of medical purpose and providing information about drugs to the population. Less than half part of the respondents considered it to be in ultimate care about the patients' health and wellness, the drugs dosage and dispensing. About one third part of the public health specialists considered it to be in creation, development, production and sale of drugs, medical devices, instruments for medical purposes and healthcare products. About one third of the health specialists considered the pharmacists to be experts of drugs; about one third of them – to be in formators of patients in cost-effectiveness and cost-benefits of drugs, the rest part of them considered that pharmacists help in selection of analogue of drugs (tabl. 3).

Table 3

The public health specialists' opinion about the pharmacist's functions in pharmacy

Q-8. Functions performed by pharmacists in pharmacy (no more than 5 answers)	Count	Percent %
1. Realization of drugs and tools (instruments) of medical purpose	164	53.4
2. Creation, development, production and sale of drugs, medical devices, instruments for medical purposes and healthcare products	110	35.8
3. Providing information about drugs to the population	165	53.7
4. Pharmaceutical care	77	25.1
5. Experts of drugs	102	33.2
6. Ultimate care about the patients' health and wellness	131	42.7
7. Dosage and dispensing of drugs	124	40.4
8. Informing the patients in pharmacotherapy direction	107	34.9
9. Informing the patients in cost-effectiveness and cost-benefits of drugs	88	28.7
10. Helping patients in offering or selection of OTC drugs	77	25.1
11. Informing the patients about drug design and drug forms	37	12.1
12. Informing the patients about drugs' generic, chemical and brand name	39	12.7
13. Informing the patients about drugs' effectiveness, safety and toxic effects	66	21.5
14. Informing patients about routes of drug administration	30	9.8
15. Informing patients about rules of drug administration	35	11.4
16. Helping in selection of analogue of drugs	97	31.6

Table 4

The respondents' (public health specialists) opinions about the issues for pharmacists necessary for the further regular studies or trainings

Q-11. The issues for pharmacists necessary for the further regular studies or trainings (several answers were possible)	Count	Percent %
1. New drugs	187	60.9
2. Psychology of communication with customers	103	33.6
3. Issues of pharmacotherapy of certain diseases	197	64.2
4. Safety and effectiveness of drugs	154	50.2
5. Pharmacology and pharmacotherapy	224	73.0
6. Normative legal regulation of pharmaceutical activity	94	30.6
7. Drugs toxicity	164	53.4
8. Drugs dosage	112	36.5
9. Routes of drug administration	110	35.8
10. Drug forms	61	19.9
11. Drug design	43	14.0
12. Rules of drug administration	123	40.1
13. Drugs generic, chemical and brand names	57	18.6
14. Selection of OTC drugs	108	35.2
15. Cost-effectiveness and cost-benefits of drugs	96	31.3

According to that pharmacist job should become regulated and more authorized in health care system [27-28].

The respondents' vast majority considered that the issues to for pharmacists were in need of the further regular studies or trainings in the following fields (tabl. 4): new medications, issues of pharmacotherapy of certain diseases, pharmacology and pharmacotherapy, drugs toxicity.

From the study results it is obvious that in the higher pharmaceutical institutions' pharmaceutical educational programs and curriculum need upgrade, renewal, modernization and adaptation to the new modern medical challenges. Therefore, continuous pharmaceutical educational programs should be created. These programs should be more focused on new medications, pharmacotherapy, drugs toxicity and dosage, routes of drug administration, selection of OTC drugs, cost-effectiveness and cost-benefits of drugs [29-30].

Estimate the impact factors, which have influence on your work satisfaction under 5-points scale (system) (estimate each factor).

On the question – Estimate the impact factors, which have influence on your work satisfaction under 5-points scale (system) (estimate each factor): the content of work (labor)-pharmacists' 3.3% estimate by 1 point, pharmacists' 6.9% estimate by 2 points, pharmacists' 14.2 % estimate by 3 points, pharmacists' 34.8 % estimate by 4 points, pharmacists' 40.7% estimate by 5 points.

On the question – Estimate the impact factors, which have influence on your work satisfaction under 5-points scale (system) (estimate each factor): position held-pharmacists' 3.5% estimate by 1 point, pharmacists' 5.1% estimate by 2 points, pharmacists' 21% estimate by 3points, pharmacists' 37.2% estimate by 4 points, pharmacists' 33.3% estimate by 5 points. (fig. 1)




Fig. 1 Position of respondents, have influenced on work satisfaction, estimated under 5-points scale (system).

On the question – Estimate the impact factors, which have influence on your work satisfaction under 5-points scale (system) (estimate each factor): correspondence of qualification to work-pharmacists' 3.2% estimate by 1 point, pharmacists' 4% estimate by 2 points, pharmacists' 15.7% estimate by 3 points, pharmacists' 34.9% estimate by 4 points, pharmacists' 42.2% estimate by 5 points [31-33] (table 5).

Table 5

Correspondence of qualification to work of respondents, which have influenced on work satisfaction, estimated under 5- points scale (system)

Estimate the impact factors, which have influence on your work satisfaction under 5- points scale (system) (estimate each factor) - Correspondence of qualification to work		Frequency	Percent	Valid percent	Cumulative percent
Valid	1	26	3.2	3.2	3.2
	2	32	4.0	4.0	7.2
	3	127	15.7	15.7	22.8
	4	283	34.9	34.9	57.8
	5	342	42.2	42.2	100.0
	Total	810	100.0	100.0	

On the question – Estimate the impact factors, which have influence on your work satisfaction under 5- points scale (system) (estimate each factor): correspondence of nature of work to my capabilities, aspirations, and inclinations (affections)-pharmacists' 2.2% estimate by 1 point, pharmacists' 5.1% estimate by 2 points, pharmacists' 16.2 % estimate by 3 points, pharmacists' 34.1% estimate by 4 points, pharmacists' 42.5 % estimate by 5 points.




Fig. 2 Existence of perspective for professional promotion (enhancement) of respondents, which have influenced on work satisfaction, estimated under 5- points scale (system).

On the question – Estimate the impact factors, which have influence on your work satisfaction under 5-points scale (system) (estimate each factor): existence of perspective for professional promotion(enhancement)-pharmacists' 5.9% estimate by 1 point, pharmacists' 8.1% estimate by 2 points, pharmacists' 16.9%estimate by 3 points, pharmacists' 32.7% estimate by 4 points, pharmacists' 36.3% estimate by 5 points. (fig. 2).

On the question – Estimate the impact factors, which have influence on your work satisfaction under 5-points scale (system) (estimate each factor): existence of perspective for career promotion (enhancement)–pharmacists' 7.4 % estimate by point 1, pharmacists, 6.7% estimate by points 2, pharmacists' 19.4 % estimate by points 3, pharmacists' 30.9% estimate by points 4, pharmacists' 35.7% estimate by points 5 (table 6).

Table 6

Existence of perspective for career promotion (enhancement) of respondents, which have influenced on work satisfaction, estimated under 5- points scale (system)

Estimate the impact factors, which have influence on your work satisfaction under 5- points scale (system) (estimate each factor) -Existence of perspective for career promotion(enhancement)		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1	60	7.4	7.4	7.4
	2	54	6.7	6.7	14.1
	3	157	19.4	19.4	33.5
	4	250	30.9	30.9	64.3
	5	289	35.7	35.7	100.0
	Total	810	100.0	100.0	

On the question – Estimate the impact factors, which have influence on your work satisfaction under 5-points scale (system) (estimate each factor): the possibility to enhance (improve) qualifications-pharmacists' 4.6% estimate by 1 point, pharmacists' 5.2% estimate by 2 points, pharmacists' 14.3% estimate by 3 points, pharmacists' 34.3% estimate by 4 points, pharmacists' 41.6% estimate by 5 points. (fig. 3).

On the question – Estimate the impact factors, which have influence on your work satisfaction under 5-points scale (system) (estimate each factor): existence of a high degree of responsibility for the result of work- pharmacists' 5.4% estimate by 1 point, pharmacists' 5.2% estimate by 2 points, pharmacists' 13.7% estimate by 3 points, pharmacists' 32.8% estimate by 4 points, pharmacists' 42.8% estimate by 5 points. (fig. 4).

On the question – Estimate the impact factors, which have influence on your work satisfaction under 5-points scale (system) (estimate each factor): regime (schedule) of work-pharmacists' 5.4% estimate by 1 point, pharmacists' 10.1% esti-

mate by 2 points, pharmacists' 25.1% estimate by 3 points, pharmacists' 31.5% estimate by 4 points, pharmacists' 27.9 % estimate by 5 points.




Fig. 3 The possibility to enhance (improve) qualifications of respondents, which have influenced on work satisfaction, estimated under 5- points scale (system). Source – study results.




Fig. 4 Existence of a high degree of responsibility for the result of work of respondents, which have influenced on work satisfaction, estimated under 5- points scale (system).

On the question – Estimate the impact factors, which have influence on your work satisfaction under 5-points scale (system) (estimate each factor): labor salary (compensation)- pharmacists' 11.6% estimate by 1 point, pharmacists' 21.6% esti-

mate by 2 points, pharmacists' 26% estimate by 3 points, pharmacists' 31% estimate by 4 points, pharmacists' 9.8% estimate by 5 points [35-37].

On the question – Estimate the impact factors, which have influence on your work satisfaction under 5-points scale (system) (estimate each factor): existence the system of benefits (incentives) scheme for employees -pharmacists' 9.8% estimate by 1 point, pharmacists' 9.4% estimate by 2 points, pharmacists' 25.7 % estimate by 3 points, pharmacists' 29.1% estimate by 4 points, pharmacists' 26% estimate by 5 points.On the question – Estimate the impact factors, which have influence on your work satisfaction under 5-points scale (system) (estimate each factor): support and assistance of a manager (chief)-pharmacists' 3.8% estimate by 1 point, pharmacists' 5.3% estimate by 2 points, pharmacists' 13.3% estimate by 3 points, pharmacists' 25.1% estimate by 4 points, pharmacists' 52.5% estimate by 5 points (fig. 5).




Fig. 5 Support and assistance of a manager (chief) of respondents, which have influenced on work satisfaction, estimated under 5- points scale (system).




Fig. 6 Direct relations with manager (chief) of respondents, which have influenced on work satisfaction, estimated under 5- points scale (system).

On the question – Estimate the impact factors, which have influence on your work satisfaction under 5-points scale (system) (estimate each factor): direct relations with manager (chief)-pharmacists' 4.2% estimate on point 1, pharmacists' 3.7% estimate on points 2, pharmacists' 10.9 % estimate on points 3, pharmacists' 26% estimate on points 4, pharmacists' 55.2% estimate on points 5 (fig. 6).

On the question – Estimate the impact factors, which have influence on your work satisfaction under 5- points scale (system) (estimate each factor): relations with (to) colleagues-pharmacists' 1.4% estimate by 1 point, pharmacists' 2.1% estimate by 2 points, pharmacists' 6.4% estimate by 3 points, pharmacists 18.6 % estimate by 4 points, pharmacists' 71.5% estimate by 5 points (fig. 7).




Fig. 7 Relations with colleagues of respondents, which have influenced on work satisfaction, estimated under 5- points scale (system).

Report on the question - Estimate the impact factors, which have influence on your work satisfaction under 5-points scale (system) (estimate each factor) (fig. 8).

About half part of the respondents considered that pharmacist is not responsible for registration of adverse effects of the drugs, while less than a third part of them considered pharmacist to be responsible for that (fig. 9). By legislation one of the functions of pharmacist is to register the side effects of drugs, what is very essential for patients' safety. It should increase the awareness of pharmacist as the health professional [38-39].

The respondents' vast majority considered that the government should make the certification of pharmacists (tabl.7). It is very essential and important that higher pharmaceutical educated pharmacist specialists to have pharmacist license issued by the government. The vast majority of the public health specialists considered that the professional activity of pharmacist is very important for the society [40-41].




Fig. 8 Report of impact factors, which have influenced on respondents work satisfaction, estimated under 5- points scale (system).




Fig. 9 The public health specialists' opinion about the pharmacist's responsible being for registration of side effects of drugs.

Thus, the pharmacists' specialty should become a regulated health care profession. According to that government should make certification, licensing and accreditation of pharmacist professionals. About a quarter of the respondents considered that pharmacist is in charge of the treatment as a physician. Properly educated pharmacist

can minimize and reduce the mistakes made by the doctor in the recipe. That has a great importance and value for provision of higher quality health care services for patients' safety [42-43].

Table 7

The respondents' (public health specialists) about the issues of certification of pharmacists by the government

Q-17. Assurance in necessity of the government certification of pharmacists	Frequency	Percent (%)	Cumulative percent (%)
1. I agree	291	94.8	94.8
2. I partially agree	14	4.6	99.4
3. I do not agree	2	0.7	100.0
Total	307	100.0	

The vast majority of the pharmacists considered that for full pharmaceutical activity it is necessary to provide continuous professional education in clinical pharmacy; therefore, higher pharmaceutical education pharmacists consider that professional education should not be ceased. The vast majority of pharmacy faculty students consider that education should not be ceased. Pharmacy faculty students' more than a third was working by specialty. The huge part of the pharmacists considers the continuous professional education is essential for the career growth and professional development, which enables getting information of new drugs and updated knowledge of some diseases' pharmacotherapy, pharmacology, clinical pharmacy and the pharmaceutical care. At the same time, the minority of respondent pharmacists had not used knowledge obtained from the professional publications and literature in their practice, while less than half of them had partially used. Competent pharmacist specialist who is capable of providing qualified pharmaceutical care (assistance) is formed in the professional training process in clinical pharmacy direction.

Mostly essential pharmaceutical activity issues for the respondents' (pharmacists) majority were: new drugs, generic drugs, chemical and brand names of them; psychology of communication (relationships) with customers; issues of pharmacotherapy of certain diseases, pharmacology, pharmacodynamics, pharmacokinetics and pharmaceutical care. It is apparent, that in the higher pharmaceutical education universities programs should be emphasized on the following subjects: pharmacotherapy, pharmacology, pharmaceutical care, clinical pharmacy and drugs toxicity.

A large majority of respondents' (pharmacists) consider that the Government should make the certification of pharmacists. As revealed, it is very important that the occupation of pharmacist should become regulated health profession. To raise pharmacists' sepecialists' professionalism, Government should make the certification of higher pharmaceutical education pharmacists. That is very essential for pharmacist's professional perfection, for successful higher pharmaceutical education, for pharmacist self-realization, for pharmacist's career advancement, for to exist pharmaceutical continuous professional education, for pharmacist professional growth, for pharma-

cist job gratification, for pharmacist career satisfaction, for pharmacists much higher status between health care specialists. Pharmacist certification is essential for pharmacists economic (material) welfare , for allows pharmacists to realize fully the received knowledge from higher education institution in work by the full extent, for to have private pharmaceutical activity, for pharmacists vocational development , for correspondence of pharmacist qualification to work, for further improvement perspective for pharmacists' professional promotion, for possibility to career enhancement strategy, for to realized by the full extent pharmacist professional capabilities, skills and habits, for occupational growth, for pharmacists professional satisfaction, for career enhancement perspective, for satisfaction of income (salary). Therefore, pharmacists' certification should start immediately and pharmacist vocation should become regulated health profession like family doctors [44-46].

The majority of the respondents (chief pharmacists) considered that main qualities, capabilities and skills required for pharmacists were ability to make decision fast and love towards their profession.

Less than half part of chief pharmacists considered that main qualities, capabilities and skills required for pharmacists were flexibility while changing the labor functions, ability to build up relations with people and high level of culture [47-48].

Therefore, the role of pharmacist is underlined in healthcare system. For the higher quality healthcare and pharmaceutical services education level is of great matter. The study provided showed that the health of patients was directly related to the professional education level of pharmacist. Therefore, pharmacist should have appropriate higher pharmaceutical education, higher professional knowledge in pharmacology, pharmaceutical care, pharmacotherapy, clinical pharmacy and other professional subjects.

For the majority of respondents' patients mostly asked the pharmacists about the rules of drugs intake and prices of drugs. For the less than half part of the respondents mostly asked about the drugs' adverse effects and quality. For about the one third of them mostly asked about help in selection of analogue of drugs, indication/contraindication of drugs, the terms and conditions of their storage (conditions and shelf-life), the drugs dosage, rules of drug administration and selection of OTC drugs.

Therefore, pharmacist should possess deep and steady knowledge in pharmacology, pharmacotherapy, toxicology, pharmaceutical care, clinical pharmacy, pharmacokinetics, pharmacodynamics, basics of medicine and other pre-clinical and clinical subjects. Such knowledge can be obtained only from higher pharmaceutical education institutions. Therefore, pharmacist working on pharmacist position must have exclusively the higher pharmaceutical education [49-50].

The respondents' public health specialists' majority considered that the pharmacists' functions in a pharmacy consisted in realization of drugs and instruments of medical purpose and providing information about drugs to the population. Less than half part of the respondents considered it to be in ultimate care about the patients' health and wellness, the drugs dosage and dispensing. About one third part of the

public health specialists considered it to be in creation, development, production and sale of drugs, medical devices, instruments for medical purposes and healthcare products. About one third of the health specialists considered the pharmacists to be experts of drugs; about one third of them – to be inform of customers in cost-effectiveness and cost-benefits of drugs, the rest part of them considered that pharmacists help in selection of analogue of drugs. According to that pharmacist job should become regulated and more authorized in health care system [51-52].

The respondents' public health specialists' majority considered that the pharmacists' functions in a pharmacy consisted in realization of drugs and instruments of medical purpose and providing information about drugs to the population. Less than half part of the respondents considered it to be in ultimate care about the patients' health and wellness, the drugs dosage and dispensing. About one third part of the public health specialists considered it to be in creation, development, production and sale of drugs, medical devices, instruments for medical purposes and healthcare products. About one third of the health specialists considered the pharmacists to be experts of drugs; about one third of them – to be inform of customers in cost-effectiveness and cost-benefits of drugs, the rest part of them considered that pharmacists help in selection of analogue of drugs. According to that pharmacist job should become regulated and more authorized in health care system [53-54].

The respondents' public health specialists' majority considered that importance in work of pharmacist was in personal realization as a specialist, receiving remuneration and provision of necessities of life. The respondents' minority considered it to be in relief of pain in suffering of people.

Less than half part of the respondents' public health specialists considered that the level of basic training of pharmacists was not corresponding to the contemporary requirements. According to the sociological study results of the public care specialists it is obviously, that all pharmacists should have higher pharmaceutical education from the state recognized and accredited higher education institutions and universities. Pharmacists' specialty should become a regulated health care profession. According to that Government should make certification, licensing and accreditation of pharmacist professionals.

The respondents' public health specialists' vast majority considered that the issues to for pharmacists were in need of the further regular studies or trainings in the following fields: new medications, issues of pharmacotherapy of certain diseases, pharmacology and pharmacotherapy, drugs toxicity. From the study results it is obvious that in the higher pharmaceutical institutions' pharmaceutical educational programs and curriculum need upgrade, renewal, modernization and adaptation to the new modern medical challenges. Therefore, continuous pharmaceutical educational programs should be created. These programs should be more focused on new medications, pharmacotherapy, drugs toxicity and dosage, routes of drug administration, selection of OTC drugs, cost-effectiveness and cost-benefits of drugs [55-56].

Approximately half part of the respondents' public health specialists was not familiar to the concept of pharmaceutical care; while more than a quarter of the public health specialists were well familiar to the concept of pharmaceutical care.

The respondents' public health specialists' large majority considered necessity of provision of cooperation between pharmacists and physicians on the issues of pharmacotherapy. The pharmacist must provide information to doctor about new drugs pharmacotherapy, the generic replacement drugs, the cost-effectiveness and cost-benefits of drugs, drugs' generic, chemical and brand names. In our opinion and vision cooperation between pharmacists and physicians on the issues of pharmacotherapy is positively reflected on patients' health and has great importance for provision higher quality health care service for patients' safety [57-58].

More than half part of the respondents' public health specialists considered that pharmacist is not in charge of treatment as a physician, meanwhile about a quarter of the public health specialists considered a pharmacist to be in charge of that. Properly educated pharmacist can minimize and reduce the mistakes made by a doctor in the recipe. That has a great importance and value for provision higher quality health care service for patients' safety.

The respondents' public health specialists' vast majority considered that pharmacist should provide assistance in teaching patients to understand the prescribed drugs intake rules. According to that higher quality pharmaceutical service could be only provided by the pharmacists of higher pharmaceutical education, graduated from the authorized, accredited and licensed by the state higher education institutes and universities.

To provide contribution and assistance in teaching of patients to understand the prescribed drugs intake rules, pharmacists need in deep knowledge in basics of medicine, pharmacology, pharmacotherapy, pharmaceutical chemistry, pharmaceutical care, clinical pharmacy and other pharmaceutical disciplines. Properly educated pharmacists have great importance and value for the provision higher quality health care services, for the provision higher quality pharmaceutical care and very essential for patient's safety.

About half part of the respondents' public health specialists considered that pharmacist is not responsible for registration of adverse effects of the drugs, while less than a third part of them considered pharmacist to be responsible for that. By legislation one of the functions of pharmacist is to register the side effects of drugs, what is very essential for patients' safety. It should increase the awareness of pharmacist as the health professional.

The respondents' public health specialists' vast majority considered that the Government should make the certification of pharmacists. It is very essential and important that higher pharmaceutical educated pharmacists to have pharmacist license issued by the Government. The vast majority of the public health specialists considered that the professional activity of pharmacist is very important for the society.

For the majority of respondents mostly significant factors while choosing a pharmacy were: service culture, wide range of products and reasonable prices. For

less than half part of respondents mostly significant factors while choosing a pharmacy were: possibility to receive consultation about medications with a physician/ a pharmacist, convenient location of the pharmacy, high qualification of pharmacist personnel.

During research the factors, influencing on the pharmacy faculty students' professional development were found and evaluated. These factors included interesting and valuable work, the favorable psychological climate within the colleagues' team, possibility of career development, professional training, social importance of profession and independence in the work.

The employed pharmacy faculty students' vast majority considered that the Government should make the certification of pharmacists to raise professional standards licensing and certification of pharmacists. The certification of higher pharmaceutical education pharmacists is very essential for the pharmacist's professional perfection, for pharmacists' career enhancement, for vocational advancement and it is main determine detector factor for pharmacist professionalism level. Pharmacist position should become regulated health profession as the member of other health profession team (but now unfortunately pharmacists are not member of regulated health teams). Pharmacists' periodic licensing, certification and accreditation should increase pharmacists' professionalism level and is guarantee upper quality pharmaceutical care. All above mentioned is indicator factor of the health care system service quality [5, 18, 29].

First time were complex studied professional peculiarities of the pharmacists per vision by pharmacists specialists, professional peculiarities of the employed pharmacist-student, professional peculiarities of the pharmacists by vision of the chief-pharmacist, peculiarities of professional for pharmacists via per vision of the health-care specialist, pharmacists' professional features as per view of the patients, professional peculiarities of the young pharmacist- specialists, professional peculiarities of the pharmacist-student. To reveals influencing factors for the specificites of the role, achievements, innovations, professional and enhancement prospects of pharmacists in health care sector. In result of the study and evaluation of the pharmacist's professional peculiarities news, objectively reasoned comprehension of the problems in this field has been adopted, which became a base for developing recommendations. In particular, for the first time the following have been studied and established: the peculiarities of professional and career improvement strategy for pharmacists, pharmacist specialist's professional features, specificites of the role, achievements, innovations, professional and enhancement prospects of pharmacists in health care sector globally. First time the process of professional formation of pharmacists in the scope and context of pharmaceutical care, including the stages of professional development was studied and scientifically established. First time the most influencive factors for the pharmacist's professional formation were identified. Deepen defined the role of pharmacist and the specific features for the pharmaceutical specialists' formation at various stages were studied and identified. On the bases of comprehensive studied was revealed, that pharmacist specialists in contradistinction

to other medical specialists like physicians, dentists etc do not have continuous education, periodic certification and licensing. Pharmacists' profession removed from the regulated and certified health professional members' team [31-34].

The results of our study have been shown and substantiated, that the pharmacists, as well as doctors and stomatologists, who are obliged to take part in the mandatory certification by the Government, in order to improve the responsibility on their own professional specialization for motivate and improve their vocational knowledge and skills with the help of continuous education.It would be promoted, that pharmacist to become more responsible, accountable and liable on for enhance their professional knowledge, skills and competencies. All the above mentioned first time we conducted a comprehensive and deep study of the scientific research for specificites of the role, achievements, innovations, professional and enhancement prospects of pharmacists in health care sector globally.

A little less than a fifth of higher pharmaceutical education pharmacists have realized professional capabilities, skills and habits to the full extent, A little bit less than half of higher pharmaceutical education pharmacists have realized professional capabilities, skills and habits partially, more than 50% of own potential, about a quarter of higher pharmaceutical education pharmacists have realized professional capabilities, skills and habits - partially, less than 50% of own potential . At the same time the vast majority of the pharmacists and health care specialists noted that pharmacists' knowledge in disciplines, such as the pharmacology, pharmacotherapy, pharmaceutical care and clinical pharmacy were a lack or insufficient for the successful work. Health care specialists' vast majority think that pharmacists are in need of additional- further regular study in the above-mentioned directions. Approximately half of the respondents considered that just 50% of their own potential was realized at the work position. Anyway, the overwhelming majority of the young pharmacists would not like to leave their profession. The vast majority of young pharmacist specialists consider that in pharmacology, in pharmacotherapy, in pharmaceutical care, in clinical pharmacy their knowledge is a lack or is not enough for successful work [22-23].

The vast majority of the pharmacists (84.4%) considered that for full pharmaceutical activity it is necessary to provide continuous professional education; therefore, higher pharmaceutical education pharmacists consider that professional education should not be ceased. The vast majority of pharmacy faculty students consider that education should not be ceased. Pharmacy faculty students' more than a third was working by specialty. The huge part of the pharmacists (55.6%) considers the continuous professional education is essential for the career growth and professional development, which enables getting information of new drugs and updated knowledge of some diseases' pharmacotherapy, pharmacology and the pharmaceutical care. At the same time, the minority of respondent pharmacists (8%) had not used knowledge obtained from the professional publications and literature in their practice, while less than half of them (41%) had partially used. Competent pharmacist specialist who is capable of providing qualified pharmaceutical care (assistance) is formed in the professional training process [16, 24, 29].

A large majority of chief pharmacists (76.6%), vast majority of patients (82.6%), of the vast majority of the employed pharmacy students (95.9%), the large majority of the healthcare specialists (94.8%) and a big majority of pharmacists (71.9%) considered that the Government should imply the pharmacists' certification in the way acting for other medical specialists. That is very essential for pharmacists' professional perfection, for self-realization, for career advancement, for continuous professional education, for professional growth [15-17].

The necessity of pharmacists' certification was stated more often by employed students than by pharmacists (Chi-square = 57.3, $p<0.001$). Statistically significant was association between patients' educational level and their opinion about the necessity of pharmacists' certification ($p<0.04$): patients with higher education considered certification of pharmacists as mandatory more often than did patients with secondary education. Chi-square test of independence revealed that pharmacists more often than students mentioned mission (the desire to obtain a profession in compliance of own trends, aspirations and inclinations, personal desire, specialty love from childhood) as the main motive of their professional choice (65.5% versus 55.8%). Difference was statistically significant with Chi-square=9.9, $p<0.002$. The difference between pharmacists and young specialists and young specialists and students wasn't statistically significant. Chi-square test of independent has been performed in order to compare the attitude of different sides to the necessity of pharmacists' certification regulation by Government. Opinion that certification of pharmacists should be mandatory was more common among health care specialists than among chiefs (Chi-square = 45.2, $p<0.001$) and among pharmacists (Chi-square = 68.9, $p<0.001$), but the there was no statistically significant difference between chiefs and pharmacists. It was more common also among patients /patients than in pharmacists (Chi-square = 44.2, $p<0.001$) [11, 16, 25, 38, 45].

Conclusions. Based on the findings of this study, several key conclusions have been drawn. The overwhelming majority of respondents emphasized the urgent need for pharmacists to receive ongoing education and training in areas such as new medications, disease-specific pharmacotherapy, pharmacology, and drug toxicity. It is evident that pharmaceutical education programs, particularly in higher institutions, require significant updates and modernization to address contemporary medical challenges. Continuous professional development programs should be implemented, with a strong focus on new medications, pharmacotherapy, drug toxicity, dosage, routes of administration, OTC drug selection, and the cost-effectiveness of treatments. The study strongly supports the regulation of the pharmacist profession as a recognized healthcare field. This regulation must include mandatory certification, licensing, and accreditation processes for pharmacists. Notably, around 25% of respondents believe that pharmacists should hold responsibilities comparable to those of physicians in patient treatment, as well-educated pharmacists can play a critical role in minimizing prescription errors and enhancing patient safety. Additionally, the establishment and expansion of residency programs for clinical pharmacists are essential to cultivate a highly qualified workforce capable of providing advanced pharmaceutical care in

clinical settings in Georgia. These measures are vital for elevating healthcare quality and ensuring patient safety.

Acknowledgments: Authors wish to Thank to Ministry of Education and Sciences of Georgia and Ministry of Education and Sciences of Armenia and Gratitude to Yerevan State Medical University and to Tbilisi State Medical University.

References:

1. Sulashvili N., Beglaryan M. Shashiashvili G. Medical Professional's job Satisfaction and pharmaceutical organization issues viewed by Drug Dispensers in Georgia. 8th Eurasian Multidisciplinary Forum, EMF 2018 (Tbilisi, 6-7 September 2018) Tbilisi, Georgia, 2018. P. 42-55.
2. National Matching Services. Summary results of the match for positions beginning in 2014. https://www.natmatch.com/ashprmp/ stats/2014applstats.html. Accessed March 24, 2014.
3. Hassell K. Centre for Pharmacy Workforce Studies (CPWS) briefing paper: GPhC register analysis2011.
4. Sulashvili N., Beglaryan M. Pharmacist's professional features and work gratification. Black sea scientific journal of academic research multidiscipline journal. 2016. Vol. 29. P. 62-68.
5. Sulashvili N., Beglaryan M. Pharmacists' professional features, viewed by the customer's (customer's) eyes in Georgia. Scientific-Practical Journal. Experimental and Clinical Medicine Tbilisi, Georgia. 2017. №4. P. 22-25.
6. Franklin B. D., van Mil J. W. Defining clinical pharmacy and pharmaceutical care. Pharm World Science. 2005; 27(3). P.137-144.
7. Sulashvili N. Peculiarities of professional and career improvement strategy for pharmacists' republic of Armenia ministry of education and science; Yerevan state medical university; Dissertation; Dissertation for the Scientific Degree of PhD in Pharmaceutical Sciences; On specialty 15.00.01 – Pharmacy; Yerevan – 2019. P. 1-175.
8. Sulashvili N. , Beglaryan M., Alavidze N. et al. Legal and regulatory scope, and identify the main challenges and opportunities of Georgian pharmacists. Bulletin of the medical institute after Mehrabyan. 2020. Vol. 9. P. 88-104.
9. Graf J, von den Driesch A, Koch K-C, Janssens U. Identification and characterization of errors and incidents in a medical intensive care unit. Acta Anaesthesiol Scand 2005;49(7):930—9.
10. Chatre C., Roubille F., Vernhet H., Jorgensen C., Pers Y-M. Cardiac complications attributed to chloroquine and hydroxychloroquine: a systematic review of the literature. Drug Saf 2018; 41(10):919—31.
11. García-Gil M., Velayos-Amo C. Hospital pharmacist experience in the intensive care unit: plan COVID. Farm Hosp. 2020;.44(7):32—5.
12. Shane P. Desselle, David P. Zgarrick, Greg Alston; Pharmacy Management: Essentials for All Practice Settings. Fourth Edition; 2016,pp. 329-347.
13. Sulashvili N. The Features of Professional Career Improvement Strategy and Job Satisfaction among pharmacists. Business-Engineering Journal. Business Engineering in Pharmacy. №2, 2014. Tbilisi, Georgia, pp. 195-199.
14. Sulashvili N., Kvizhinadze N., Maisuradze I. Pharmacist professional features in Georgia. // Conference of young scientists. Thesis collection. Georgian National Academy of Sciences. 18-19 May 2015. Tbilisi, Georgia, pp. 81-82.
15. Sulashvili N., M Beglaryan. Pharmacist mission gratification and profession improvement strategy. Black sea scientific journal of academic research conference newsletter. (Medicine, Pharmacy sciences). Volume 26. November 2015. Tbilisi, Georgia, pp. 10-12.
16. Sulashvili N., Beglaryan M. Pharmacist's professional features and work gratification. // Black sea scientific journal of academic research multidiscipline journal. (Medicine, Pharmacy sciences) Vol. 29. March-April 2016. Tbilisi, Georgia, pp. 62-68.

17. Sulashvili N., Beglaryan M. Pharmacists' professional features, viewed by the customer's (customer's) eyes in Georgia. Experimental and Clinical Medicine, Scientific-Practical Journal, №4, 2017. Tbilisi, Georgia, pp. 22-25.
18. Sulashvili, N. Peculiarities of professional for pharmacists, viewed by the health-care specialists in Georgia. Experimental and Clinical Medicine, Scientific-Practical Journal. №4 2017. Tbilisi, Georgia, pp.47-51.
19. Sulashvili N., Beglaryan M. Characteristics of pharmacist activity, viewed by the customer's//
20. International Science and Innovation Festival 2017. Conf.-es "Healthy Lifestyle-Scientific Evidence and Controversial issues" and "Innovation in Medicine" Tbilisi State Medical University. September 2017. Tbilisi, Georgia. pp. 30-31.
21. Sulashvili N., Beglaryan M. Professional features for employed pharmacy faculty students in Georgia. The New Armenian Medical Journal Supplement. YSMU Science Week 2017 Conference. November 27-December 1, Vol.11, №3, 2017, Yerevan, Armenia, p. 40.
22. Sulashvili N., Beglaryan M. Vocational peculiarities of young pharmacist professionals. International scientific journal Intellectual. №35, 2018. Tbilisi, Georgia. pp. 96-104.
23. Sulashvili N., Beglaryan M., Gogashvili M., Matoshvili M. Occupational particularities and strategy of Pharmacy faculty students. Experimental and Clinical Medicine, Scientific-Practical Journal. №3, 2018. Tbilisi, Georgia. pp. 15-24.
24. Sulashvili, N., Beglaryan M., Matoshvili M. Occupational features of pharmaceutical workers, viewed by the chief pharmacists. Caucasus Journal of Health Sciences and Public Health. Volume 2, Issue 2, June 2018.Tbilisi-Georgia. pp.56-61.
25. Sulashvili, N., Beglaryan M., Sulashvili M. Personal features, capabilities and skills of job adaptation for pharmacist specialists. Tbilisi David Agmashenebeli University Periodical Scientific Journal №13, 2018. Tbilisi, Georgia. pp. 231-236.
26. Sulashvili N., Beglaryan M., Kvijinadze N., Matoshvili M. Vocational training and activity of pharmacists in Georgia. Georgian Medical News. № 5 (278), May 2018. Tbilisi - New York. pp.199-206.
27. Michael Ira Smith, Albert I. Wertheimer, Jack E. Fincham. Pharmacy and the US Health Care System; Pharmaceutical Press. 4th Revised edition edition (April 15, 2013);pp. 402-426.
28. Sulashvili N., Beglaryan M. Pharmacist Occupational Features, Regulations Framework and Profession Enhancement Challenges of Pharmaceutics. Caucasus Journal of Health Sciences and Public Health. 2020; Vol. 4, Supplement 8, July 20-24; Pp. 31-36.
29. N. Kvizhinadze, D. Tophuria, N. Intskirveli, N. Sulashvili; Study of Factors Affecting on Population's Health Improvement.Caucasus Journal of Health Sciences and Public Health; E ISSN 2449-2450; ISSN 2449-2647; The University of Georgia Publishing House. www.caucasushealth.ge; Official Journal of the University of Georgia and Iv. Javakhishvili Tbilisi State University with Support of the Arctic University of Tromsø/Norway. Volume 4, Supplement 8, July 20-24, 2020; Pp 42-45.
30. Thomas R. Brown. Handbook of Institutional Pharmacy Practice // 4th (fourth) Edition; American Society of Health-System Pharmacists (December 1, 2006); pp. 65-71.
31. Sulashvili N., Beglaryan M. (2017). Characteristics of pharmacist activity as viewed by customers. In International Science and Innovation Festival 2017: Conferences "Healthy Lifestyle—Scientific Evidence and Controversial Issues" and "Innovation in Medicine" (pp. 30-31). Tbilisi State Medical University, Tbilisi, Georgia.
32. Sulashvili N., Beglaryan, M. (2017). Professional features for pharmacy faculty students in Georgia. The New Armenian Medical Journal. Supplement, 11(3), 40. YSMU Science Week 2017 Conference, Yerevan, Republic of Armenia.
33. Sulashvili N. (2019). Peculiarities of the professional and career improvement strategy for pharmacists. Abstract of the dissertation for the Ph.D. in Pharmaceutical Sciences (Specialty

15.00.01 – Pharmacy). Yerevan State Medical University After M. Heratsi, Ministry of Education and Science of the Republic of Armenia. Yerevan, 1-26.
34. Sulashvili N., Alavidze N., Abuladze N., Kvizhinadze N., Gogashvili M., Beglaryan M. (2020). Aspects of pharmaceutical professional and organizational issues in Georgia. In Materials of the 12th Scientific and Practical Internet Conference: Pharmacoeconomics in Ukraine, Conditions and Development Prospects (pp. 11-20). National University of Pharmacy, Kharkiv, Ukraine.
35. Sulashvili N. (2019). Peculiarities of the professional and career improvement strategy for pharmacists. Abstract of the dissertation for the Ph.D. in Pharmaceutical Sciences (Specialty 15.00.01 – Pharmacy). Specialized Council 026 "Theoretical Medicine" of the Supreme Certifying Committee of the Republic of Armenia, Yerevan State Medical University, 1-26.
36. Sulashvili N. (2019). Peculiarities of the professional and career improvement strategy for pharmacists. Dissertation for the Ph.D. in Pharmaceutical Sciences (Specialty 15.00.01 – Pharmacy). Ministry of Education and Science of the Republic of Armenia, Yerevan State Medical University, 1-175.
37. Sulashvili N., Beglaryan M., Zarnadze I., Zarnadze Sh., Alavidze N., Abuladze N. Kvizhinadze N. (2020). Vocational perspectives and the main professional opportunities and challenges for pharmacy faculty students in Georgia. In Materials of the V International Scientific and Practical Conference: Technological and Biopharmaceutical Aspects of Drug Development with Different Orientations of Action. Kharkiv, Ukraine. P. 35-51.
38. Sulashvili N., Chichoyan N., Gerzmava O., Tsintsadze T., Gabunia L., Kvizhinadze N., Okropiridze T., Beglaryan M. (2020). Organizational and regulatory framework: Identifying the main educational issues and potentials of young pharmacists in Georgia. In Materials of the V International Scientific and Practical Conference: Technological and Biopharmaceutical Aspects of Drug Development with Different Orientations of Action. Kharkiv, Ukraine. P. 51-69).
39. Finley P. R., Rens H. R., Pont J. T., et al. (2002). Impact of a collaborative pharmacy practice model on the treatment of depression in primary care. American Journal of Health-System Pharmacy. 59, 1518–1526.
40. Saari L. M., Judge T. A. Employee attitudes and job satisfaction. Hum Resour Manage. 2004; P. 65-69.
41. Locke E. A., What is job satisfaction? Organ Behav Hum Perform. 2009; 4; pp. 309–336.
42. Spector P. E., Job Satisfaction: Application, Assessment, Causes and Consequences. London: SAGE Publications. 2011; P. 78-82.
43. Liekens S., Vandael E., Roter D., Larson S., Smits T., Laekeman G., Foulon V. Impact of training on pharmacists' counseling of patients starting antidepressant therapy. Patient Educ. Couns. 2014; 94:110–115.
44. Rubio-Valera M., March Pujol M., Fernández A., Peñarrubia-María M.T., Travé P., López del Hoyo Y., Serrano-Blanco A. Evaluation of a pharmacist intervention on patients initiating pharmacological treatment for depression: A randomized controlled superiority trial. Eur. Neuropsychopharmacol. 2013; 23:1057–1066.
45. Arroll B., Goodyear-Smith F., Crengle S., Gunn J., Kerse N., Fishman T., Falloon K., Hatcher S. Validation of PHQ-2 and PHQ-9 to screen for major depression in the primary care population. Ann. Fam. Med. 2010; 8:348–353.
46. Ayorinde A.A., Porteous T., Sharma P. Screening for major diseases in community pharmacies: A systematic review. Int. J. Pharm. Pract. 2013; 21:349–361.
47. Elsever. Qualitative insights into job satisfaction and dissatisfaction with management among community and hospital pharmacists 2011; pp. 145-149.
48. Seston E., Hassell K., Ferguson J., Hann M. Exploring the relationship between pharmacists' job satisfaction, intention to quit the profession, and actual quitting // Res Social Adm Pharm. 2009; 5(2): pp. 121-32.

49. Cooper C. L., Rout U., Faragher B. Mental health, job satisfaction and job stress among pharmacists. BMJ. 2009; 298(6670); pp. 366-370.
50. Jain V., Priya R., Shubhi R., and others. Evaluation of job satisfaction and social identity of pharmacists. Ind J Pharm Prac. 2013; 6(1); pp. 47-58.
51. Hardigan P., Carvajal M. Job satisfaction among practicing pharmacists: a Rasch analysis. The Internet Journal of Allied Health Sciences and Practice. 2007; pp. 98-102.
52. Maio V., Goldfarb N. I., Hartmann C. W., Pharmacists' job satisfaction: variation by practice setting. P & T. 2004; P. 123-127.
53. Royal Pharmaceutical Society. Professional Standards for Hospital Pharmacy Services. London; 2014.
54. David Tipton; Professionalism, Work, and Clinical Responsibility in Pharmacy. Jones& Bartlett Learning; 1 edition (March 15, 2013); P. 205-269.
55. Seston E., Hassell K., Ferguson J., Hann M. Exploring therelationship between pharmacists' job satisfaction, intention to quit the profession, and actual quitting. Res Social Adm Pharm. 2009; P.52-59.
56. Hare S., Kraenow K. Depression screenings: Developing a model for use in a community pharmacy. J. Am. Pharm. Assoc. 2008; 48:46–51.
57. Knox E., Dopheide J., Wincor M., Han P. Depression screening in a university campus pharmacy: A pilot project. J. Am. Pharm. Assoc. 2006; 46: 502–506.
58. Rosser S., Frede S., Conrad W.F., Heaton P.C. Development, implementation, and evaluation of a pharmacist-conducted screening program for depression. J. Am. Pharm. Assoc. 2013; 53:22–29.

ЗМІСТ

РОЗДІЛ 1. КЛІНІКО-ФАРМАКОЛОГІЧНІ АСПЕКТИ ЗАСТОСУВАННЯ ЛІКАРСЬКИХ ЗАСОБІВ ……... 3

Міщенко О. Я., Бутко Я. О., Андріяненков О. В.
КЛІНІЧНА ФАРМАКОЛОГІЯ ТА
ПЕРСПЕКТИВНІ ТЕНДЕНЦІЇ РОЗРОБКИ ЛІКІВ ……... 4

Podolskyi I. M., Podolskyi M. I.
STEREOCHEMICAL ASPECTS OF PHARMACODYNAMICS
AND PHARMACOKINETICS OF MEDICINES …… 7

Otrishko I. A., Bezugla N. P., Vietrova K. V., Mesbahi H.
FACTORS CONTRIBUTING TO MEDICATION ADHERENCE
IN MOROCCAN ADULT ASTHMA PATIENTS:
CLINICAL AND PHARMACEUTICAL RESEARCH …… 9

Shchokina K. H.
COMPARATIVE CHARACTERISTICS OF MODERN GROUPS OF
DRUGS AFFECTING THE RENIN-ANGIOTENSIN-ALDOSTERONE
SYSTEM ……. 10

Bondariev Ye. V., Tymchenko Yu. V.
EVALUATION OF CLINICAL AND PHARMACEUTICAL EFFICACY
OF THE TRIAZOLE DERIVATIVES USE
IN MODERN GYNECOLOGICAL PRACTICE …….. 12

Berezniakov A., Bondariev Ye.
CLINICAL EFFICACY AND SAFETY OF THE ANTIHISTAMINIC
DRUG EBASTIN ……... 14

Крайдашенко О. О.
ФАРМАКОТЕРАПІЯ ДИСБАЛАНСУ СИСТЕМИ ПРОТЕАЗИ
АНТИПРОТЕАЗИ У ОСІБ ЗІ СТАБІЛЬНОЮ СТЕНОКАРДІЄЮ
НАПРУГИ В ПОЄДНАННІ З ХРОНІЧНИМ ОБСТРУКТИВНИМ
ЗАХВОРЮВАННЯМ ЛЕГЕНЬ ……... 16

Отрішко І. А., Вєтрова К. В., Безугла Н. П., Ковальчук А. О.
ВИВЧЕННЯ ОСОБЛИВОСТЕЙ ПОКАЗНИКІВ
ЯКОСТІ ЖИТТЯ ПАЦІЄНТІВ НА ФОНІ ЗАМІСНОЇ
ГОРМОНАЛЬНОЇ ТЕРАПІЇ ГІПОТИРЕОЗУ……. 17

Seniuk I. V., Kravchenko V. M., Riyad Qamouta
CLINICAL PHARMACOLOGY FROM THE PERSPECTIVE
OF THE WORLD HEALTH ORGANISATION ……... 18

Кахніашвілі А. С., Деримедвідь Л. В.
МЕТОДИКИ КОРЕКЦІЇ
ПОСТТРАВМАТИЧНОГО СТРЕСОВОГО РОЗЛАДУ…… 20

Кравченко В. М., Сенюк І. В., Галузінська Л. В.
КЛІНІЧНА ФАРМАКОЛОГІЯ НА ОСНОВІ ІНТЕРФЕРЕНЦІЇ РНК….. 22

Савельєва-Кулик Н. О.
ТЕРАПЕВТИЧНИЙ ПОТЕНЦІАЛ, БЕЗПЕКА І МІЖЛІКАРСЬКІ
ВЗАЄМОДІЇ ROSMARINUS OFFICINALIS L. ……. 24

Штробля А. Л., Гольцева С. С.
ОБІЗНАНІСТЬ ФАРМАЦЕВТІВ ЩОДО КЛІНІЧНОЇ ХАРАКТЕРИСТИКИ
ТРИВОЖНИХ РОЗЛАДІВ ТА НАДАННЯ ФАРМАЦЕВТИЧНОЇ
ОПІКИ ПРИ ЛІКУВАННІ ПОРУШЕНЬ ДІЯЛЬНОСТІ ЦНС …….. 26

Гриньків Я. О.
ТОПІРАМАТ ДЛЯ ПРОФІЛАКТИКИ МІГРЕНІ …….. 28

Карпінчик Л. С., Харченко Ю. В.
ВИКОРИСТАННЯ ІНФУЗІЙНИХ РОЗЧИНІВ ПРИ ГІПОКАЛІЄМІЇ
У ПАЦІЄНТІВ ЗАГАЛЬНОЇ ХІРУРГІЇ …….. 29

Хворост О. П., Опрошанська Т. В., Скребцова К. С.
АСОРТИМЕНТ МІОРЕЛАКСАНТІВ У ТЕРАПІЇ ЗАХВОРЮВАНЬ
ОПОРНО-РУХОВОГО АПАРАТУ …… 30

Tovchiga O. V., Inkielewicz-Stępniak I.
MECHANISMS ASSOCIATED WITH URIC ACID
IN NEUROPROTECTIVE STRATEGIES …… 31

Makovetskyi A., Zaychenko G.
VAGINAL DRUG DELIVERY SYSTEMS: INNOVATIONS THAT
CHANGE THE PHARMACOLOGICAL PROPERTIES OF DRUGS …… 33

Туранська Н. П., Вєтрова К. В., Отрішко І. А.
РОЛЬ ФАРМАЦЕВТА В ЗАБЕЗПЕЧЕННІ РАЦІОНАЛЬНОГО
ЛІКУВАННЯ ПАЦІЄНТІВ З АРТЕРІАЛЬНОЮ ГІПЕРТЕНЗІЄЮ …… 37

Білоусова Н. А.
ФУНКЦІОНАЛЬНЕ НАПОВНЕННЯ РОЛЕЙ ФАРМАЦЕВТІВ
ПРИ ПРОФІЛАКТИЦІ ІШЕМІЧНОЇ ХВОРОБИ СЕРЦЯ
НА ПЕРВИННОМУ РІВНІ МЕДИЧНОЇ ДОПОМОГИ ……..... 38

Сахарова Т. С.
ОПТИМІЗАЦІЯ СИМПТОМАТИЧНОГО
ЛІКУВАННЯ ТРИВОГИ У ПАЦІЄНТІВ З АРТЕРІАЛЬНОЮ 40
ГІПЕРТЕНЗІЄЮ В УМОВАХ АПТЕКИ ……

Ващенко О. О.
ДЕЯКІ АСПЕКТИ ЗАСТОСУВАННЯ
ШАМПУНІВ ІЗ КЕТОКОНАЗОЛОМ ПРИ ЛУПІ …….. 41

Катурова В. В.
ТРАВМИ, ЩО НАНЕСЕНІ ДОМАШНІМИ ТВАРИНАМИ ТА ЇХ 42
ЛІКУВАННЯ …….

Ривак Т. Б., Сувала Н. С.
ВИВЧЕННЯ ДУМКИ ПАЦІЄНТІВ І ФАРМАЦЕВТІВ ЩОДО
ЗАСТОСУВАННЯ РЕЦЕПТУРНИХ ЛІКАРСЬКИХ ЗАСОБІВ …….. 43

Bayraktar V. M.
TREATMENT OF HYPERESTROGENISM IN MEN OF FERTILE AGE
WITH TESTICULAR TESTOSTERONE DEFICIT ……. 45

Tarapata Michael, Kukhtenko Oleksandr
BEE-HERBAL PRODUCTS AND PROBIOTICS COMBINATION
FOR THERAPEUTIC ENHANCEMENT IN SKIN DISEASES …….. 48

Деримедвідь Л. В., Риженко І. М., Богуцька О. Є., Кузенков Р. В.
ПЕРЕВАГИ КОМБІНОВАНОЇ ТЕРАПІЇ ЦУКРОВОГО ДІАБЕТУ
КОМБІНАЦІЄЮ МЕТФОРМІНУ З ІНГІБІТОРАМИ
НАТРІЙ-ГЛЮКОЗНОГО КОТРАНСПОРТЕРА 2-ГО ТИПУ ……. 50

Севастьянова Т. В., Яшна П. О., Антонов В. С.
ВИБІР НЕСТЕРОЇДНИХ ПРОТИЗАПАЛЬНИХ ЗАСОБІВ
МІЖ ВІТЧИЗНЯНИМИ ТА ЗАКОРДОННИМИ ПРЕПАРАТАМИ ….. 52

Русанова А. І.
РОЛЬ РАНОЛАЗИНУ ЯК АНТИАРИТМІЧНОГО ПРЕПАРАТУ
В СУЧАСНІЙ МЕДИЧНІЙ ПРАКТИЦІ …… 59

Ткачова О. В., Белік Г. В.
ПОШУК ДОКАЗОВОЇ БАЗИ КЛІНІЧНОЇ ЕФЕКТИВНОСТІ
ПРОБІОТИКІВ ПРИ ЛІКУВАННІ ГОСТРОЇ ДІАРЕЇ …….. 62

Sulashvili Nodar, Gabunia Luiza, Beglaryan Margarita,
Gorgaslidze Nana, Ghambashidze Ketevan, Sulashvili Marika
THE SCIENTIFIC TALKS OF MANIFESTATION SOME TOPICAL
ISSUE ASPECTS OF FEATURES OF PHARMACOLOGY
OF THE ANTIARRHYTHMIC DRUGS IN GENERAL ……. 70

РОЗДІЛ 2. БЕЗПЕКА ФАРМАКОТЕРАПІЇ ТА СИСТЕМА ФАРМАКОНАГЛЯДУ ……..
85

Крайдашенко О. В., Косогор Ю. А., Каптур Л. М.
НЕБАЖАНІ ПОБІЧНІ РЕАКЦІЇ ЛІКАРСЬКИХ ЗАСОБІВ:
ФАРМАКОНАГЛЯД В ЗАПОРІЗЬКІЙ ОБЛАСТІ.
ПІДСУМКИ І ПІВРІЧЧЯ 2024 РОКУ ……. 86

Баюрка С. В., Карпушина С. А.
ЕПІДЕМІОЛОГІЯ ОТРУЄНЬ АНТИДЕПРЕСАНТАМИ ……. 88

Бут О. В., Хайтович М. В.
ФАРМАЦЕВТИЧНА ОПІКА В РАМКАХ РЕАБІЛІТАЦІЇ
ПАЦІЄНТІВ ІЗ ЗЛОЯКІСНИМИ НОВОУТВОРЕННЯМИ
ПІСЛЯ КУРСУ ХІМІОТЕРАПІЇ ……. 90

Яцкова Г. Ю.
ФАРМАЦЕВТИЧНА ПРОФІЛАКТИКА ПРИ ЗАСТОСУВАННІ
СТАТИНІВ ПРИ АРТЕРІАЛЬНІЙ ГІПЕРТЕНЗІЇ ……. 91

Козар В. В., Кондратенко К. К.
ПРОБЛЕМА ВПЛИВУ ЕЛЕКТРОННИХ СИГАРЕТ
НА ЗДОРОВ’Я ЛЮДИНИ ……... 93

Одинець І. Ю., Міщенко В. А., Пороша Н. С.,
Лисенко Л. С., Пушкар М. Б.
КЛІНІКА УРАЖЕНЬ НЕРВОВОЇ СИСТЕМИ ПРИ ГОСТРИХ
ОТРУЄННЯХ СИНТЕТИЧНИМИ КАНАБІНОЇДАМИ У ДІТЕЙ …….. 95

Кайдаш С. П., Слєсарчук В. Ю.
БЕЗПЕКА ВИКОРИСТАННЯ АНТИГІПЕРТЕНЗИВНИХ
ПРЕПАРАТІВ ПЕРШОГО РЯДУ ……... 97

Команюк Л. В., Паламар А.О.
ОСОБЛИВОСТІ ЗАСТОСУВАННЯ ЛІКІВ У ВАГІТНИХ ……. 99

Gabunia Luiza, Ghambashidze Ketevan, Ratiani Levan, Gorgaslidze Nana, Khetsuriani Shorena, Gamkrelidze Natia, Varazi Elena, Rukhadze Londa, Peikrishvili Nana, Sulashvili Nodar.
THE SCIENTIFIC TALKS OF MANIFESTATION OF TOPICAL ISSUES OF CHARACTERISTICS OF COGNITIVE-ENHANCING STIMULANT DRUGS ACCORDING THE SAFETY OF PHARMACOTHERAPY AND PHARMACOVIGILANCE SYSTEM IN GENERAL …… 101

РОЗДІЛ 3. ВІКОВІ, ГЕНДЕРНІ, ХРОНОФАРМАКОЛОГІЧНІ ТА ФАРМАКОГЕНЕТИЧНІ ЧИННИКИ РАЦІОНАЛЬНОГО ЗАСТОСУВАННЯ ЛІКАРСЬКИХ ЗАСОБІВ …….. 122

Крайдашенко О. В., Кремзер О. О., Самура Т. О.
ФАРМАКОГЕНЕТИЧНЕ ТЕСТУВАННЯ:
СУЧАСНИЙ СТАН ПРОБЛЕМИ ……... 123

Chopyak V. V., Fedorov Yu. V., Kovpak A. V.
GENDER ASPECTS OF PHARMACOTHERAPY …….. 124

Гришко Ю. М., Луценко Р. В.
РОЛЬ ВІКОВИХ, ГЕНДЕРНИХ, ХРОНОФАРМАКОЛОГІЧНИХ ТА ФАРМАКОГЕНЕТИЧНИХ ЧИННИКІВ У РАЦІОНАЛЬНІЙ ФАРМАКОТЕРАПІЇ …… 125

Файзуллін О. В., Халєєва О. Л., Бондарєв Є. В.
ДО ФАРМАКОГЕНЕТИКИ МЕТФОРМІНУ:
СУЧАСНИЙ СТАН ПРОБЛЕМИ ……... 127

Tsiklauri Mzia, Chitaia Guram, Kvirkvelia Nino, Sulashvili Nodar
THE SCIENTIFIC TALKS OF MANIFESTATION OF TOPICAL ISSUES ASPECTS OF CHARASTERICSTICS OF ASSESSMENT OF HEALTH CONDITION AND PHARMACOTHERAPY CHALLENGES OF COMPULSORILY RELOCATED CHILDREN FROM ABKHAZIA - WHICH IS THE INTEGRAL PART OF GEORGIA 132

РОЗДІЛ 4. АДМІНІСТРУВАННЯ АНТИМІКРОБНИХ ПРЕПАРАТІВ ЯК МУЛЬТИДИСЦИПЛІНАРНИЙ ПІДХІД ПРОТИДІЇ АНТИБІОТИКОРЕЗИСТЕНТНОСТІ …… 145

Паламар А. О., Круглик Я. Д.
ПИТАННЯ ПОДОЛАННЯ АНТИБІОТИКОРЕЗИСТЕНТНОСТІ
В УКРАЇНІ ……. 146

Кравченко В. М., Філімонова Н. І., Сенюк І. В.
ПЕРЕХРЕСТЯ АНТИБІОТИКОРЕЗИСТЕНТНОСТІ
ТА БІОСИНТЕЗУ АНТИБІОТИКІВ ……... 148

Шаповалова О. В., Кучма І. Ю.
ПАНДЕМІЯ COVID-19:
РОКИ ВПЛИВУ НА АНТИМІКРОБНУ РЕЗИСТЕНТНІСТЬ …….. 150

Воронкіна А. С., Дідківська А.. О., Хорунжий Д.. С.
АНАЛІЗ РИНКУ ДИТЯЧИХ АНТИМІКРОБНИХ ЛІКАРСЬКИХ
ЗАСОБІВ ДЛЯ ПЕРОРАЛЬНОГО ПРИЙОМУ:
РИЗИКИ АНТИБІОТИКОРЕЗИСТЕНТНОСТІ ……. 152

Kolomiets Y., Kharchenko Y.
THERAPY FOR MULTIDRUG-RESISTANT TUBERCULOSIS …….. 154

Паскевич М.. П., Жаботинська Н.. В.
ВИВЧЕННЯ РОЛІ ФАРМАЦЕВТА В ЗАСТОСУВАННІ
ПРОБІОТИКІВ ПРИ ПРОВЕДЕННІ АНТИБАКТЕРІАЛЬНОЇ ТЕРАПІЇ 155

Міщенко О. Я., Андріяненков О. В., Шаповалова О. В.
АНАЛІЗ ПРОБЛЕМНИХ ПИТАНЬ ПРОФЕСІЙНОЇ ДІЯЛЬНОСТІ
КЛІНІЧНИХ ФАРМАЦЕВТІВ У ВІДДІЛАХ ІНФЕКЦІЙНОГО
КОНТРОЛЮ ……. 156

Sulashvili Nodar, Robakidze Kakhaber, Beglaryan Margarita,
Gabunia Luiza, Gorgaslidze Nana, Alavidze Nato, Grigolia Lela,
Tadevosyan Ada (Adel), Khundzakishvili Veriko, Abuladze Nino,
Kvizhinadze Natia, Sulashvili Marika.
THE MANIFESTATION OF KEY ISSUE ASPECTS OF MULTIDISCI-
PLINARY STRATEGIES TO COMBAT ANTIBIOTIC RESISTANCE:
EMERGING THERAPEUTIC APPROACHES AND GLOBAL
CHALLENGES IN GENERAL …….. 168

РОЗДІЛ 5. ІНФОРМАЦІЙНІ ТЕХНОЛОГІЇ В КЛІНІЧНІЙ ФАРМАКОЛОГІЇ ТА КЛІНІЧНІЙ ФАРМАЦІЇ …….. 181

Назаркіна В. М., Кривенко Е. В.
ЦИФРОВІ ТЕХНОЛОГІЇ В ДОКАЗОВІЙ МЕДИЦИНІ:
ПІДТРИМКА ПРИЙНЯТТЯ РІШЕНЬ ЩОДО
РАЦІОНАЛЬНОГО ЗАСТОСУВАННЯ ЛІКІВ ……. 182

Подольський І. М., Назаркіна В. М., Бондарєв Є. В.
РЕПОЗИЦІОНУВАННЯ ЛІКАРСЬКИХ ЗАСОБІВ:
ЕВОЛЮЦІЯ ПІДХОДІВ …….. 187

РОЗДІЛ 6. ОРГАНІЗАЦІЙНО-УПРАВЛІНСЬКІ ПІДХОДИ ЗАБЕЗПЕЧЕННЯ РАЦІОНАЛЬНОГО ЗАСТОСУВАННЯ ЛІКАРСЬКИХ ЗАСОБІВ ……... 189

Зарічкова М. В., Мішина І. Ю.
ВДОСКОНАЛЕННЯ ПОСАДОВОЇ ІНСТРУКЦІЇ
ФАРМАЦЕВТА КЛІНІЧНОГО НА ОСНОВІ РЕЗУЛЬТАТІВ
ЕКСПЕРТНОГО ОЦІНЮВАННЯ ПРОФЕСІЙНОЇ ДІЯЛЬНОСТІ ……. 190

Братішко Ю. С., Толочко В. М., Зарічкова М. В., Артюх Т. О.,
Ковальська О. В.
АНАЛІЗ ЗАРУБІЖНОГО ТА ВІТЧИЗНЯНОГО ДОСВІДУ
ДЕРЖАВНОГО РЕГУЛЮВАННЯ СОЦІАЛЬНОЇ
ВІДПОВІДАЛЬНОСТІ ФАРМАЦЕВТИЧНОГО БІЗНЕСУ ……. 200

Назаркіна В. М., Іванова А. Д.
ПРОБЛЕМИ ТА ПЕРСПЕКТИВИ РЕЦЕПТУРНОГО ВІДПУСКУ
ЛІКАРСЬКИХ ЗАСОБІВ В УКРАЇНІ ……. 214

Залигіна Є. В.
АНАЛІЗ ВІДХИЛЕНЬ У ЛИСТАХ ЛІКАРСЬКИХ ПРИЗНАЧЕНЬ
У СТАЦІОНАРІ ХІРУРГІЧНОГО ПРОФІЛЮ
З МЕТОЮ УДОСКОНАЛЕННЯ СИСТЕМИ УПРАВЛІННЯ
ФАРМАЦЕВТИЧНОЮ СЛУЖБОЮ ЗОЗ …….. 216

Зіменковський А. Б., Шпортько М. О., Настюха Ю. С.
МОЖЛИВІСТЬ ІМПЛЕМЕНТАЦІЇ ПРАКТИКИ
МЕДИКАМЕНТОЗНОГО УЗГОДЖЕННЯ У ПРОФЕСІЙНУ
ДІЯЛЬНІСТЬ КЛІНІЧНИХ ФАРМАЦЕВТІВ В УКРАЇНІ …….. 217

Chehrynets A. A., Bondarieva I. V., Khaleq S.
ANALYSIS OF ETHICAL ASPECTS OF PHARMACISTS' WORK ……. 219

Герасимова О. О., Ткачова О. В.
ОЦІНКА СПОЖИВАННЯ В УКРАЇНІ ЛІКАРСЬКИХ ЗАСОБІВ, ЩО ПРИГНІЧУЮТЬ ПЕРИСТАЛЬТИКУ …… 220

Bondarieva I. V., Malyi V. V., Ifrad Yu.
STUDY OF CASES OF SUCCESSFUL USE OF SOCIAL MEDIA BY PHARMACEUTICAL COMPANIES ……. 221

Kvizhinadze N., Dughashvili N., Instkirveli N., Kiknavelidze N.
STRATEGIC MANAGEMENT ELEMENTS,
USED FOR GEORGIAN PHARMACEUTICAL SECTOR ……. 225

Shashiashvili N., Kvizhinadze N., Gaprindashvili M., Dughashvili N., Kiknavelidze N., Instkirveli N.
PECULIARITIES OF REFERENCE PRICING POLICY IN THE PHARMACEUTICAL SECTOR OF GEORGIA …….. 227

Kvizhinadze N., Intskirveli N., Tophuria D., Kiknavelidze N., Dughashvili N.
CHARACTERISTICS OF CUSTOMER BEHAVIOR
IN PHARMACY SERVICE IN GEORGIA …….. 229

Sulashvili Nodar, Okropiridze Tamar, Gorgaslidze Nana, Beglaryan Margarita, Gabunia Luiza, Ghambashidze Ketevan, Kvizhinadze Natia, Alavidze Nato, Khundzakishvili Veriko, Robakidze Kakhaber, Grigolia Lela, Sulashvili Marika
THE SCIENTIFIC DISCUSSION OF THE MANIFESTATION OF THE TOPICAL ISSUES ASPECTS OF CHARACTERISTICS OF SOME UP-TO-DATE CONSIDERATIONS REPERCUSSION ON PATIENTS' DENTAL HEALTH CARE AND PHARMACOTHERAPHY CHALLENGES IN GENERAL ……. 230

РОЗДІЛ 7. ОЦІНКА ТЕХНОЛОГІЙ ОХОРОНИ ЗДОРОВ'Я, ФАРМАКОЕКОНОМІЧНІ ДОСЛІДЖЕННЯ ТА ФОРМУЛЯРНА СИСТЕМА ЯК СКЛАДОВІ ЗАБЕЗПЕЧЕННЯ РАЦІОНАЛЬНОЇ ФАРМАКОТЕРАПІЇ …….. 250

Коляда Т. А.
ПРАВОВІ МЕХАНІЗМИ РЕАЛІЗАЦІЇ ДЕРЖАВНОЇ ПОЛІТИКИ

ЩОДО ЗАБЕЗПЕЧЕННЯ РАЦІОНАЛЬНОГО ВИКОРИСТАННЯ ЛІКАРСЬКИХ ЗАСОБІВ ……. 251

Баліцька О. П., Гайдай О. Д.
АНАЛІЗ МІНІМІЗАЦІЇ ВИТРАТ ІНГІБІТОРІВ ПРОТОННОЇ ПОМПИ, ЯКІ ЗАСТОСОВУВАЛИСЬ ДЛЯ ЛІКУВАННЯ ВИРАЗКОВОЇ ХВОРОБИ ДВАНАДЦЯТИПАЛОЇ КИШКИ В УМОВАХ СТАЦІОНАРУ ……... 253

Харченко Ю. В., Ткачова О. В.
АНАЛІЗ ВАРТОСТІ КУРСУ ЛІКУВАННЯ ДЕЗЛОРАТАДИНОМ ЗА МЕТОДОМ МІНІМІЗАЦІЇ ВИТРАТ ПРИ АЛЕРГІЧНОМУ РИНІТІ 254

Ткачова О. В., Процька В. В.
ОЦІНКА ВАРТОСТІ КУРСУ ЛІКУВАННЯ ВИРАЗКИ ШЛУНКУ ОМЕПРАЗОЛОМ ЗА МЕТОДОМ МІНІМІЗАЦІЇ ВИТРАТ …… 255

Ткачова О. В., Герасимова О. О., Яковлєва Л. В.
ДОСЛІДЖЕННЯ СПОЖИВАННЯ АНТИДЕПРЕСАНТІВ В УКРАЇНІ ПОРІВНЯНО З ДЕЯКИМИ КРАЇНАМИ ЄВРОПИ ……. 256

Fedotova M. S., Panfilova G. L.
COMPARATIVE ANALYSIS OF THE EFFECTIVENESS OF MEMANTINE AND ITS COMBINATIONS WITH ACETYLCHOLINESTERASE INHIBITORS IN PATIENTS WITH ALZHEIMER'S DISEASE ……. 257

Заболотня З. О., Заліська О. М., Калинюк Т. Г.
АНАЛІЗ ЕКСТЕМПОРАЛЬНИХ ЛІКАРСЬКИХ ЗАСОБІВ ПРИ ПОШИРЕНИХ ДЕРМАТОЛОГІЧНИХ ЗАХВОРЮВАННЯХ У США І В УКРАЇНІ ДЛЯ ОПТИМІЗАЦІЇ ФОРМУЛЯРНОГО ЗАБЕЗПЕЧЕННЯ …….. 258

Жадько С. В., Севрюков О. В., Бабічева Г. С., Чуфицький Є. О.
МАРКЕТИНГОВИЙ ТА ФАРМАКОЕКОНОМІЧНИЙ АНАЛІЗ ГІПОЛІПІДЕМІЧНИХ ЛІКАРСЬКИХ ЗАСОБІВ, ЩО МІСТЯТЬ СИМВАСТАТИН ……... 260

Адонкіна В. Ю., Міщенко О. Я., Грешко Ю. І.
АНАЛІЗ ЦЕНТРАЛІЗОВАНИХ ДЕРЖАВНИХ ЗАКУПІВЕЛЬ ЛІКАРСЬКИХ ЗАСОБІВ ДЛЯ ЕНДОКРИННОЇ ТЕРАПІЇ РАКУ МОЛОЧНОЇ ЗАЛОЗИ ……... 265

РОЗДІЛ 8. ТЕХНОЛОГІЧНІ (НАНОТЕХНОЛОГІЧНІ) АСПЕКТИ РОЗРОБКИ, ВПРОВАДЖЕННЯ ТА ВИКОРИСТАННЯ ЛІКАРСЬКИХ ЗАСОБІВ …….. 267

Сагайдак-Нікітюк Р. В., Губченко Т. Д., Шевченко В. О.
СТРАТЕГІЇ ПОКРАЩЕННЯ ДОСТАВЛЯННЯ ЛІКІВ В ОКО ……. 268

Іванова А. Д., Сліпченко Г. Д.
ПЕРСПЕКТИВА СТВОРЕННЯ НОВИХ ЗАСОБІВ
У ВИГЛЯДІ ЛАЙНЕРІВ …….. 270

Усенко Д. Л., Каплаушенко А. Г.
РОЗРОБКА МЕТОДИКИ ДЕТЕКТУВАННЯ КОМПЛЕКСУ
СУПУТНІХ ДОМІШОК НАТРІЙ 2-((4-АМІНО-5-(ТІОФЕН-2-
ІЛМЕТИЛ)-4*Н*-1,2,4-ТРІАЗОЛ-3-ІЛ)ТІО)АЦЕТАТУ
ЗА ДОПОМОГОЮ ВЕРХ-МС ……. 271

Хворост О. П., Опрошанська Т. В., Скребцова К. С.
ПЕРСПЕКТИВНІ ВИДИ ВІТЧИЗНЯНОЇ ЛІКАРСЬКОЇ РОСЛИННОЇ
СИРОВИНИ, ДЖЕРЕЛОМ ЯКОЇ Є РОСЛИНИ РОДИНИ *LAMIACEAE* 272

РОЗДІЛ 9. ДОКЛІНІЧНІ ТА КЛІНІЧНІ ДОСЛІДЖЕННЯ ЛІКАРСЬКИХ ЗАСОБІВ …… 273

Isachenko V., Ostashko V., Isachenko E.
NOVEL AND PROMISING EMBRYO CULTURE MEDIA
IN HUMAN IVF PROGRAMS ……. 274

Дембіцька Е. С., Цубанова Н. А., Семійон-Лучишин О. В.
ПЕРСПЕКТИВИ ЗАСТОСУВАННЯ КАТЕХІНІВ,
ЯК НОВИХ ТЕРАПЕВТИЧНИХ АГЕНТІВ
ДЛЯ ЛІКУВАННЯ ХРОНІЧНОЇ ВЕНОЗНОЇ НЕДОСТАТНОСТІ …… 275

Довбня Д. В., Каплаушенко А. Г.
ВИЗНАЧЕННЯ ФАРМАКОКІНЕТИЧНИХ ПАРАМЕТРІВ
ПОХІДНИХ 5-(2,4-, 3,4-ДИМЕТОКСИФЕНІЛ)-3*Н*-1,2,4-ТРІАЗОЛ-3-
ТІОНІВ ЗА ДОПОМОГОЮ ADME-АНАЛІЗУ ……. 276

Цубанова Н. А., Волощук Н. І., Застрижна М. Л.
ВИВЧЕННЯ РЕПАРАТИВНОЇ ДІЇ ЕКСТРАКТУ ТРАВИ
SAPONARIA OFFICINALIS НА МОДЕЛІ ЛІНІЙНИХ РІЗАНИХ РАН ... 277

Цубанова Н. А., Волощук Н. І., Галевич Г. Б.
ВИВЧЕННЯ ВПЛИВУ ЕКСТРАКТУ ТРАВИ *TAGETES PATULA L*
НА МОДЕЛІ СТРЕПТОЗОТОЦИНОВОГО ДІАБЕТУ …… 278

Білай І. М., Білай С. І., Білай А. І.
ВИВЧЕННЯ ФАРМАКОДИНАМІЧНИХ ЕФЕКТІВ
КВЕРЦЕТИНУ ……. 280

Цивунін В. В., Штриголь С. Ю., Северіна Г. І.
ЦЕРЕБРАЛЬНІ ПРОТИЗАПАЛЬНІ ВЛАСТИВОСТІ ДИГОКСИНУ
ЯК НЕОЧЕВИДНИЙ КОМПОНЕНТ ЙОГО ПРОТИСУДОМНОЇ ДІЇ ... 281

Верховодова Ю. В.
ВИВЧЕННЯ ЗМІН ГЕМАТОЛОГІЧНИХ ПОКАЗНИКІВ У ЩУРІВ В
УМОВАХ ЕКСПЕРИМЕНТАЛЬНОГО КОЛІТУ ПРИ ЛІКУВАННІ
МОДИФІКОВАНИМИ ФІТОЕКСТРАКТАМИ SALVIA OFFICINALIS 282

Тулуб І. О., Бурда Н. Є. , Фіра Л. С.
ВИВЧЕННЯ АНТИОКСИДАНТНОЇ АКТИВНОСТІ
ЕКСТРАКТІВ ЦИНІЇ ВИТОНЧЕНОЇ ……. 283

Ільїна С. К., Журавель І. О.
ДОСЛІДЖЕННЯ ПРОТИМІКРОБНОЇ АКТИВНОСТІ РОТИКІВ
САДОВИХ ТРАВИ ЕКСТРАКТУ ГУСТОГО ……... 284

Дрезналь Є. П., Кметь Т. І.
ВПЛИВ КАРБАЦЕТАМУ НА ГЛУТАТІОНОВИЙ ЛАНЦЮГ
ГІПОКАМПА СТАРИХ ЩУРІВ З ЕКСПЕРИМЕНТАЛЬНОЮ
ХВОРОБОЮ АЛЬЦГЕЙМЕРА …… .. 286

Андреєва І. Д., Осолодченко Т. П., Батрак О. А., Рябова І. С.
СТАБІЛЬНІСТЬ ПРОТИМІКРОБНОГО ЕФЕКТУ
ГЕЛЮ НА ОСНОВІ КОМПОЗИЦІЇ АЦИЛЬОВАНОГО НІЗИНУ
З ДИКЛОФЕНАКОМ НАТРІЯ ЩОДО КЛІНІЧНИХ
ШТАМІВ МІКРООРГАНІЗМІВ ……. 288

Осолодченко Т. П., Андреєва І. Д., Мартинов А. В., Завада Н. П.
ВИЗНАЧЕННЯ СТАБІЛЬНОСТІ ПРОТИМІКРОБНОГО ЕФЕКТУ
ГЕЛЮ НА ОСНОВІ КОМПОЗИЦІЇ СУКЦИЛЬОВАНОГО НІЗИНУ З
ДИКЛОФЕНАКОМ НАТРІЯ ЩОДО КЛІНІЧНИХ ШТАМІВ
МІКРООРГАНІЗМІВ …….. 289

Хільковець А. В., Білай І. М.
ДОСЛІДЖЕННЯ ГОСТРОЇ ТОКСИЧНОСТІ НОВИХ
ТІОФЕН ПОХІДНИХ 1,2,4-ТРИАЗОЛІВ …….. 290

Кушнір О. Ю., Кушнір О. О.
АКТИВНІСТЬ ЛАКТАТДЕГІДРОГЕНАЗИ В ПЕЧІНЦІ ЩУРІВ З
АЛОКСАНОВИМ ДІАБЕТОМ ЗА УМОВ УВЕДЕННЯ
МЕЛАТОНІНУ ……. 291

Іванченко Д. Г., Крісанова Н. В.
ДОСЛІДЖЕННЯ СПАЗМОЛІТИЧНИХ ТА ПРОТИЗАПАЛЬНИХ
ВЛАСТИВОСТЕЙ 8-АМІНО-1-БЕНЗИЛ-7-ЕТИЛ-3-
МЕТИЛКСАНТИНІВ ……... 292

Іванченко Д. Г., Рудько Н. П.
ПОШУК ГІПОЛІПІДЕМІЧНИХ ЗАСОБІВ СЕРЕД
8-АМІНО-7-(2-ГІДРОКСИ-2-ФЕНІЛЕТИЛ)-3-МЕТИЛКСАНТИНІВ ... 293

Шпичак О. С., Міщенко О. Я.
ВИВЧЕННЯ ПАРАМЕТРІВ ГОСТРОЇ ТОКСИЧНОСТІ РОСЛИННОЇ
СУБСТАНЦІЇ КАЛИНИ ЗВИЧАЙНОЇ ПЛОДІВ ЕКСТРАКТУ
РІДКОГО ……... 295

Хворост О. П., Опрошанська Т. В., Скребцова К. С., Горда А. О.
ПЕРСПЕКТИВНІ ВИДИ ВІТЧИЗНЯНОЇ ЛІКАРСЬКОЇ РОСЛИННОЇ
СИРОВИНИ, ЩО ЗАСТОСОВУЮТЬСЯ У ТЕРАПІЇ ЗАХВОРЮВАНЬ
ОПОРНО-РУХОВОГО АПАРАТУ ……. 297

Galuzinska L. V., Seniuk I. V.
MODERN ASPECTS OF COMPLEX THERAPY
FOR PATIENTS WITH PSORIASIS …… 298

Малоштан Л. М., Підгайна В. В., Яковлєва Л. В.
ВИВЧЕННЯ ПРОТИЗАПАЛЬНОЇ АКТИВНОСТІ
КРЕМУ «ПІРИСАЛІКС» ……. 299

Єрмоленко Т. І., Шаповал О. М.
ПЕРСПЕКТИВИ ВИКОРИСТАННЯ ЦИТОСТАТИКА
ЦИКЛОФОСФАМІДУ ДЛЯ ЕКСПЕРИМЕНТАЛЬНОГО
ВІДТВОРЕННЯ ВТОРИННОГО ІМУНОДЕФІЦИТУ У ЩУРІВ
ДИТЯЧОГО ВІКУ …… 300

Прижбило О. М., Кметь О. Г.
ПАТОГЕНЕТИЧНЕ ОБГРУНТУВАННЯ ЕФЕКТИВОСТІ
КАРБАЦЕТАМУ ПРИ МЕТАБОЛІЧНИХ ПОРУШЕННЯХ У ЩУРІВ . 302

Tovchiga O. V., Koiro O. O., Shtrygol' S. Yu., Stepanova S. I.
COMBINED USE OF THE SUBSTANCES OF HERBAL ORIGIN
WITH XANTHINE OXIDASE INHIBITORS:
EXAMPLE OF *AEGOPODIUM PODAGRARIA* L. ……. 304

Chankseliani Gocha, Girdaladze Avtandil, Gibradze Omar ,
Meshveliani Paata, Chelidze Kakha, Cheishvili Mirian, Kvernadze Ana,
Sulashvili Nodar
THE TOPICAL ISSUES ASPECTS OF CHARACTERISTICS OF
ENDOVASCULAR SURGERY AND PHARMACOLOGICAL
AND NON-PHARMACOLOGICAL TREATMENT CHALLENGES
FOR GASTROINTESTINAL AND DUODENAL ULCER BLEEDING
WITH BRIEF CASE REPORT ……. 307

РОЗДІЛ 10. МЕТОДИЧНІ АСПЕКТИ ВИКЛАДАННЯ ФАРМАКОЛОГІЇ, КЛІНІЧНОЇ ФАРМАКОЛОГІЇ ТА КЛІНІЧНОЇ ФАРМАЦІЇ В УКРАЇНІ ТА СВІТІ ... 323

Filipets N. D., Gerush O. V., Filipets O. O.
FORMATION OF THERAPEUTIC THINKING OF PHARMACIST
STUDENTS IN THE TEACHING OF PHARMACOLOGY
AND CLINICAL PHARMACY …… 324

Крайдашенко О. В., Свинтозельський О. О., Михайлик О. А.
ДОСВІД ЗАСТОСУВАННЯ ДИСТАНЦІЙНИХ ФОРМ НАВЧАННЯ
З АКЦЕНТОМ НА ФОРМУВАННЯ КЛІНІЧНОГО МИСЛЕННЯ
СУЧАСНОГО МАГІСТРА ФАРМАЦІЇ ……. 326

Панькевич О. Б., Громовик Б. П.
ФАРМАЦЕВТИЧНА БЕЗПЕКА
З ПОГЛЯДУ ФАРМАЦЕВТИЧНОЇ ТЕРМІНОСИСТЕМИ ……. 327

Слєсарчук В. Ю., Кайдаш С. П., Соколова К. В.
ПРОПОЗИЦІЇ ТА ОБҐРУНТУВАННЯ ВВЕДЕННЯ ПЕВНИХ
ВИБІРКОВИХ КОМПОНЕНТІВ ДО ОСНОВНИХ
В ОСВІТНЬО-ПРОФЕСІЙНІЙ ПРОГРАМІ СПЕЦІАЛЬНОСТІ
226 «ФАРМАЦІЯ, ПРОМИСЛОВА ФАРМАЦІЯ» ……... 328

Горчакова Н. О., Шумейко О. В., Клименко О. В.
МЕТОДИЧНІ ТА ТЕХНОЛОГІЧНІ АСПЕКТИ
ВИКЛАДАННЯ ФАРМАКОЛОГІЇ ……. 329

Кузнєцова В. Ю., Федосов А. І., Файзуллін О. В.
ПЕРСПЕКТИВИ РОЗРОБКИ ПРОФЕСІЙНОГО СТАНДАРТУ
«ПРОФЕСІОНАЛ З КЛІНІЧНИХ ДОСЛІДЖЕНЬ» …… 331

Артюх Т. О.
ДОСВІД КАФЕДРИ ОРГАНІЗАЦІЇ, ЕКОНОМІКИ ТА
УПРАВЛІННЯ ФАРМАЦІЄЮ ІПКСФ НФАУ
З ПРОВЕДЕННЯ РЕФЛЕКСІЇ ФАРМАЦЕВТІВ-ІНТЕРНІВ
ПІД ЧАС ДИСТАНЦІЙНОГО НАВЧАННЯ ……. 333

Бондарєв Є. В., Бутко Я. О., Міщенко О. Я., Файзуллін О. В.,
Тимченко Ю. В., Подольский І. М.
ДОСВІД ВИКОРИСТАННЯ ДИСТАНЦІЙНОЇ СКЛАДОВОЇ
ОЧНО-ДИСТАНЦІЙНОГО НАВЧАННЯ В УМОВАХ ВОЄННОГО
СТАНУ НА КАФЕДРІ КЛІНІЧНОЇ ФАРМАКОЛОГІЇ …… 335

Кишкан І. Г.
ПИТАННЯ БЕЗПЕКИ ЛІКІВ ТА БЕЗПЕЧНОЇ ФАРМАКОТЕРАПІЇ
ПРИ ВИВЧЕННІ ПОБІЧНОЇ ДІЇ ЛІКІВ СТУДЕНТАМИ
СПЕЦІАЛЬНОСТІ «ФАРМАЦІЯ, ПРОМИСЛОВА ФАРМАЦІЯ» ……. 341

Штриголь С. Ю., Жаботинська Н. В., Кіреєв І. В.
ПОРІВНЯЛЬНИЙ АНАЛІЗ УКРАЇНСЬКОГО ТА МІЖНАРОДНОГО
ДОСВІДУ ПІДГОТОВКИ КЛІНІЧНОГО ФАРМАЦЕВТА ……. 342

Sulashvili Nodar, Beglaryan Margarita, Gorgaslidze Nana, Gabunia Luiza, Kvizhinadze Natia, Alavidze Nato, Abuladze Nino, Gabunia Ketevani, Khetsuriani Shorena, Gamkrelidze Natia, Gumbaridze Lia, Rukhadze Londa, Pkhakadze Giorgi, de Jauregui Diego Rada Fernandez, Sulashvili Marika, Giorgobiani Marina, Okropiridze Tamar, Seniuk Igor, Grigolia Lela, Robakidze Kakhaber, Zarnadze Irine, Zarnadze Shalva (Davit)
SCIENTIFIC DISCOURSE OF ELEVATING THE ROLE
OF PHARMACISTS: A GLOBAL AND GEORGIAN PERSPECTIVES
ON CLINICAL RESPONSIBILITIES, PROFESSIONAL
ENHANCEMENT, JOB CONTENTMENT AND HEALTH CARE
IMPACT ……. 347

Наукове видання

АКТУАЛЬНІ ПИТАННЯ КЛІНІЧНОЇ ФАРМАКОЛОГІЇ ТА КЛІНІЧНОЇ ФАРМАЦІЇ (TOPICAL ISSUES OF CLINICAL PHARMACOLOGY AND CLINICAL PHARMACY)

Матеріали науково-практичної
Internet-конференції з міжнародною участю

29-30 жовтня 2024 року
м. Харків

Формат 60х84/16. Ум. друк. арк. 14,25. Тираж 100 пр. Зам. № 1121/10-24.

Національний фармацевтичний університет
вул. Григорія Сковороди, 53, м. Харків, 61002
Свідоцтво серії ДК № 3420 від 11.03.2009 р.

Надруковано з готового оригінал-макета у друкарні ФОП В. В. Петров
Єдиний державний реєстр юридичних осіб та фізичних осіб-підприємців.
Запис № 24800000000106167 від 08.01.2009 р.
61144, м. Харків, вул. Гв. Широнінців, 79в, к. 137, тел. (057) 78-17-137.
e-mail: bookfabrik@mail.ua